

A Proximal Sampling Algorithm

Jiaming Liang

Department of Computer Science
Yale University

September 28, 2022

SIAM Mathematics of Data Science, San Diego


Joint works with Yongxin Chen (Georgia Tech)

Travel made possible by NSF grant DMS-1757085


Introduction



Design and analysis of fast algorithms for **sampling** problems by leveraging tools from **optimization**.



(a) Optimization, $\min f(x)$



(b) Sampling, $\text{samp} \exp(-f(\bar{x}))$

Overview

- A proximal sampling algorithm for nonconvex, semi-smooth and composite potentials
- Improved complexity to sample from a distribution ε -close to the target distribution in KL, χ^2 and Rényi divergences
- Close interplay between sampling and optimization
Proximal point framework

Assumptions

Problem: sample from $\nu(x) \propto \exp(-f(x))$

(A1) f is semi-smooth, i.e., there exist $\alpha_i \in [0, 1]$ and $L_i > 0$, $i = 1, \dots, n$, s.t.

$$\|f'(u) - f'(v)\| \leq \sum_{i=1}^n L_{\alpha_i} \|u - v\|^{\alpha_i}, \quad \forall u, v \in \mathbb{R}^d;$$

Examples: $n = 1$

1) $\alpha_1 = 1$, smooth, 2) $\alpha_1 = 0$, nonsmooth, 3) $0 < \alpha_1 < 1$, weakly smooth

(A2) ν satisfies log-Sobolev inequality (LSI) or Poincaré inequality (PI).

$$\text{LSI: } H_\nu(\rho) \leq \frac{C_{LSI}}{2} J_\rho(\nu), \quad \text{PI: } \mathbb{E}_\nu[(\psi - \mathbb{E}_\nu[\psi])^2] \leq C_{PI} \mathbb{E}_\nu[\|\nabla \psi\|^2]$$

Observations: ν is **not** necessarily log-concave, f is **not** necessarily convex.

Comparison

Source	Complexity	Assumption	Metric
Chewi et al.	$\tilde{O}\left(\frac{C_{\text{PI}}^{1+1/\alpha} L_{\alpha}^{2/\alpha} d^{2+1/\alpha}}{\varepsilon^{1/\alpha}}\right)$	weakly smooth $\alpha > 0$, PI	Rényi
This work	$\tilde{O}\left(C_{\text{PI}} L_{\alpha}^{2/(1+\alpha)} d^2\right)$	semi-smooth, PI	Rényi

Table: Complexity bounds for sampling from non-convex semi-smooth potentials.

Source	Complexity	Assumption	Metric
Nguyen et al.	$\tilde{O}\left(C_{\text{LSI}}^{1+\max\{\frac{1}{\alpha_i}\}} \left[\frac{n \max\{L_{\alpha_i}^2\} d}{\varepsilon}\right]^{\max\{\frac{1}{\alpha_i}\}}\right)$	weakly smooth $\alpha_i > 0$, LSI	KL
This work	$\tilde{O}\left(C_{\text{LSI}} \sum_{i=1}^n L_{\alpha_i}^{2/(\alpha_i+1)} d\right)$	semi-smooth, LSI	KL
This work	$\tilde{O}\left(C_{\text{PI}} \sum_{i=1}^n L_{\alpha_i}^{2/(\alpha_i+1)} d\right)$	semi-smooth, PI	Rényi

Table: Complexity bounds for sampling from non-convex composite potentials.

Alternating Sampling Framework (ASF)

Joint distribution $\pi(x, y) \propto \exp[-f(x) - \frac{1}{2\eta}\|x - y\|^2]$

Algorithm 1 ASF (Shen, Tian and Lee 2021)

1. Sample $y_k \sim \pi^{Y|X}(y \mid x_k) \propto \exp[-\frac{1}{2\eta}\|x_k - y\|^2]$
 2. Sample $x_{k+1} \sim \pi^{X|Y}(x \mid y_k) \propto \exp[-f(x) - \frac{1}{2\eta}\|x - y_k\|^2]$
-

Restricted Gaussian Oracle (RGO)

Given y , sample from

$$\pi^{X|Y}(\cdot|y) \propto \exp\left(-f(\cdot) - \frac{1}{2\eta}\|\cdot - y\|^2\right).$$

Alternating Sampling Framework (ASF)

Joint distribution $\pi(x, y) \propto \exp[-f(x) - \frac{1}{2\eta}\|x - y\|^2]$

Algorithm 2 ASF (Shen, Tian and Lee 2021)

1. Sample $y_k \sim \pi^{Y|X}(y \mid x_k) \propto \exp[-\frac{1}{2\eta}\|x_k - y\|^2]$
 2. Sample $x_{k+1} \sim \pi^{X|Y}(x \mid y_k) \propto \exp[-f(x) - \frac{1}{2\eta}\|x - y_k\|^2]$
-

Restricted Gaussian Oracle (RGO)

Given y , sample from

$$\pi^{X|Y}(\cdot|y) \propto \exp\left(-f(\cdot) - \frac{1}{2\eta}\|\cdot - y\|^2\right).$$

Without an implementable and provable RGO, ASF is only conceptual.

Nontrivial

Proximal Point Framework (PPF)

Proximal point framework: constructs a sequence of proximal problems

$$x_{k+1} \leftarrow \text{prox}_{\eta f}(x_k) = \underset{x}{\operatorname{argmin}} \left\{ f(x) + \frac{1}{2\eta} \|x - x_k\|^2 \right\} \quad (*) \quad (1)$$

E.g., Chambolle-Pock for saddle point, ADMM for distributed optimization

Algorithm 3 PPF

1. $y_k \leftarrow \underset{x}{\operatorname{argmin}} \frac{1}{2\eta} \|x - x_k\|^2 = x_k$
 2. $x_{k+1} \leftarrow \underset{x}{\operatorname{argmin}} \left\{ f_{y_k}^{\eta}(x) := f(x) + \frac{1}{2\eta} \|x - y_k\|^2 \right\}$
-

ASF for sampling \longleftrightarrow PPF for optimization

RGO in sampling \longleftrightarrow proximal mapping in optimization


Relaxed Proximal Bundle Method (L. and Monteiro 2021)

f is convex and Lipschitz continuous (nonsmooth, $\alpha_1 = 0$). Subgradient method.

Approximately solve (1) by the cutting-plane method ([implementable](#))

$$z_j \leftarrow \text{prox}_{\eta f_j}(x_0) = \min_z \left\{ f_j(z) + \frac{1}{2\eta} \|z - z_0\|^2 \right\}, \quad z_0 = x_k$$

where $f_j(z) = \max\{f(w) + \langle f'(w), z - w \rangle : w \in \{z_0, z_1, \dots, z_{j-1}\}\}$



Complexities: PPF $\mathcal{O}(\varepsilon^{-1}) \times$ cutting-plane $\mathcal{O}(\varepsilon^{-1}) \implies$ total $\mathcal{O}(\varepsilon^{-2})$ **optimal**

[implementable and provable](#)

RGO Implementation

RGO: given y , sample from $\exp(-f_y^\eta(x))$

Algorithm 4 RGO Rejection Sampling

1. Compute an approximate stationary point w of f_y^η
2. Generate sample $X \sim \exp(-h_1(x))$
3. Generate sample $U \sim \mathcal{U}[0, 1]$
4. If

$$U \leq \frac{\exp(-f_y^\eta(X))}{\exp(-h_1(X))},$$

then accept/return X ; otherwise, reject X and go to step 2.

Proposal: $\exp(-h_1(x))$ where $h_1(x) \leq f_y^\eta(x)$

Rejection Sampling

$X \sim \pi^{X|Y}(\cdot|y)$ and

$$\begin{aligned}\mathbb{P}(X \text{ is accepted}) &= \mathbb{P}\left(U \leq \frac{\exp(-f_y^\eta(X))}{\exp(-h_1(X))}\right) \\ &= \frac{\int \exp(-f_y^\eta(x)) dx}{\int \exp(-h_1(x)) dx} \geq \frac{\int \exp(-h_2(x)) dx}{\int \exp(-h_1(x)) dx}\end{aligned}\quad (2)$$

Want to find h_1 and h_2 such that

i) sampling $\exp(-h_1(x))$ is easy,

ii) $h_1(x) \leq f_y^\eta(x) \leq h_2(x)$,

iii) (2) is bounded from below.

A Key Lemma

Consider $n = 1$, $\alpha \in [0, 1]$ and $L_\alpha > 0$

$$\|f'(u) - f'(v)\| \leq L_\alpha \|u - v\|^\alpha, \quad \forall u, v \in \mathbb{R}^d;$$

Lemma

Assume f is L_α -semi-smooth, then for $\delta > 0$ and every $u, v \in \mathbb{R}^d$, we have

$$|f(u) - f(v) - \langle f'(v), u - v \rangle| \leq \frac{M}{2} \|u - v\|^2 + \frac{(1 - \alpha)\delta}{2}, \quad M = \frac{L_\alpha^{\frac{2}{\alpha+1}}}{[(\alpha + 1)\delta]^{\frac{1-\alpha}{\alpha+1}}}.$$

Proof:

$$|f(u) - f(v) - \langle f'(v), u - v \rangle| \leq \frac{L_\alpha}{\alpha + 1} \|u - v\|^{\alpha+1}$$

Young's inequality $ab \leq \frac{a^p}{p} + \frac{b^q}{q}$, $\frac{1}{p} + \frac{1}{q} = 1$ with

$$a = \frac{L_\alpha}{(\alpha + 1)\delta^{\frac{1-\alpha}{2}}} \|u - v\|^{\alpha+1}, \quad b = \delta^{\frac{1-\alpha}{2}}, \quad p = \frac{2}{\alpha + 1}, \quad q = \frac{2}{1 - \alpha}.$$

Construction

Definition

A stationary point w^* of f_y^η is such that $f'(w^*) + \frac{1}{\eta}(w^* - y) = 0$.

Definition

An approximate stationary point w of f_y^η is s.t. $\|f'(w) + \frac{1}{\eta}(w - y)\| \leq \sqrt{Md}$.

$$h_1(x) := f(w) + \langle f'(w), x - w \rangle - \frac{M}{2} \|x - w\|^2 + \frac{1}{2\eta} \|x - y\|^2 - \frac{(1 - \alpha)\delta}{2},$$

$$h_2(x) := f(w^*) + \langle f'(w^*), x - w^* \rangle + \frac{M}{2} \|x - w^*\|^2 + \frac{1}{2\eta} \|x - y\|^2 + \frac{(1 - \alpha)\delta}{2}.$$

Answers: i) sampling $\exp(-h_1(x))$ is easy;
ii) verify $h_1(x) \leq f_y^\eta(x) \leq h_2(x)$ by the key lemma.

Remaining Questions

Q1. Rejection sampling complexity

$$[\mathbb{P}(X \text{ is accepted})]^{-1} \leq \frac{\int \exp(-h_1(x))dx}{\int \exp(-h_2(x))dx}$$

Q2. Optimization complexity to find an approx. stat. pt. w s.t.

$$\left\| f'(w) + \frac{1}{\eta}(w - y) \right\| \leq \sqrt{Md}$$

Answer to Q1 – RGO complexity

Proposition

Assume

$$\eta \leq \frac{1}{Md} = \frac{[(\alpha + 1)\delta]^{\frac{1-\alpha}{\alpha+1}}}{L_{\alpha+1}^{\frac{2}{\alpha+1}} d},$$

then the expected number of rejection steps in Algorithm 4 is at most $\exp\left(\frac{3(1-\alpha)\delta}{2} + 3\right)$.

Intuition: if η is small enough, h_1 and h_2 are convex quadratic functions, so

$$\frac{\int \exp(-h_1(x))dx}{\int \exp(-h_2(x))dx} \approx \left(\frac{1 + \eta M}{1 - \eta M}\right)^{d/2} \leq (1 + 4\eta M)^{d/2} \leq \left(1 + \frac{4}{d}\right)^{d/2} \leq e^2.$$

Answer to Q2 – Optimization complexity

Lemma

Let $f_y^\eta := f + \frac{1}{2\eta} \|\cdot - y\|^2$ and $(f_y^\eta)' := f' + \frac{1}{\eta}(\cdot - y)$, then for every $u, v \in \mathbb{R}^d$,

$$\begin{aligned} \frac{1}{2} \left(\frac{1}{\eta} - M \right) \|u - v\|^2 - \frac{(1 - \alpha)\delta}{2} &\leq f_y^\eta(u) - f_y^\eta(v) - \langle (f_y^\eta)'(v), u - v \rangle \\ &\leq \frac{1}{2} \left(\frac{1}{\eta} + M \right) \|u - v\|^2 + \frac{(1 - \alpha)\delta}{2}. \end{aligned}$$

f_y^η is nearly $(\eta^{-1} - M)$ -strongly convex and $(\eta^{-1} + M)$ -smooth

Proposition

Assume $\eta \leq \frac{1}{Md}$, then the iteration-complexity to find the approx. stat. pt. w s.t. $\|f'(w) + \frac{1}{\eta}(w - y)\| \leq \sqrt{Md}$ by Nesterov acceleration is $\tilde{O}(1)$.

$$\mu = \frac{1}{\eta} - M \approx M(d - 1), \quad L = \frac{1}{\eta} + M \approx M(d + 1), \quad \sqrt{L/\mu} \approx 1$$

RGO and ASF Complexities

Putting previous results together, we can implement RGO with $\tilde{\mathcal{O}}(1)$ subgradients of f and $\mathcal{O}(1)$ samples from Gaussian distribution in expectation.

Other ingredients for total complexity: **Convergence rate analysis of ASF**

Theorem (Chen, Chewi, Salim and Wibisono 2022)

If $\nu \propto \exp(-f)$ satisfies LSI with $C_{LSI} > 0$, then x_k of ASF $\sim \rho_k$, which satisfies

$$H_\nu(\rho_k) \leq \frac{H_\nu(\rho_0)}{\left(1 + \frac{\eta}{C_{LSI}}\right)^{2k}}.$$

Theorem (Chen, Chewi, Salim and Wibisono 2022)

If $\nu \propto \exp(-f)$ satisfies PI with $C_{PI} > 0$, then x_k of ASF $\sim \rho_k$, which satisfies

$$\chi_\nu^2(\rho_k) \leq \frac{\chi_\nu^2(\rho_0)}{\left(1 + \frac{\eta}{C_{PI}}\right)^{2k}}.$$

Main Result

Theorem

Suppose f is L_α -semi-smooth and ν satisfies PI. With $\eta \asymp 1/(L_\alpha^{\frac{2}{\alpha+1}} d)$, then ASF with RGO by rejection has complexity bound

$$\tilde{\mathcal{O}}\left(C_{\text{PI}} L_\alpha^{\frac{2}{\alpha+1}} d\right)$$

to achieve ε error to ν in terms of χ^2 divergence. Each iteration queries $\tilde{\mathcal{O}}(1)$ subgradients of f and generates $\mathcal{O}(1)$ samples in expectation from Gaussian distribution.

General Results – LSI

$$\|f'(u) - f'(v)\| \leq \sum_{i=1}^n L_{\alpha_i} \|u - v\|^{\alpha_i}, \quad \forall u, v \in \mathbb{R}^d;$$

Theorem

Suppose f is semi-smooth and ν satisfies LSI. With $\eta \asymp \left[\sum_{i=1}^n L_{\alpha_i}^{\frac{2}{\alpha_i+1}} d \right]^{-1}$, then ASF with RGO by rejection has complexity bound

$$\tilde{\mathcal{O}} \left(C_{\text{LSI}} \sum_{i=1}^n L_{\alpha_i}^{\frac{2}{\alpha_i+1}} d \right)$$

to achieve ε error to ν in terms of χ^2 divergence. Each iteration queries $\tilde{\mathcal{O}}(1)$ subgradients of f and generates $\mathcal{O}(1)$ samples in expectation from Gaussian distribution.

Theorem

Suppose f is semi-smooth and ν satisfies PI. With $\eta \asymp \left[\sum_{i=1}^n L_{\alpha_i}^{\frac{2}{\alpha_i+1}} d \right]^{-1}$, then ASF with RGO by rejection has complexity bound

$$\tilde{\mathcal{O}} \left(C_{\text{PI}} \sum_{i=1}^n L_{\alpha_i}^{\frac{2}{\alpha_i+1}} d \right)$$

to achieve ε error to ν in terms of χ^2 divergence. Each iteration queries $\tilde{\mathcal{O}}(1)$ subgradients of f and generates $\mathcal{O}(1)$ samples in expectation from Gaussian distribution.

Interpretation of Unadjusted Langevin Algorithm (ULA)

Algorithm 5 ASF

1. Sample $y_k \sim \pi^{Y|X}(y | x_k) \propto \exp[-\frac{1}{2\eta} \|x_k - y\|^2]$
 2. Sample $x_{k+1} \sim \pi^{X|Y}(x | y_k) \propto \exp[-f(x) - \frac{1}{2\eta} \|x - y_k\|^2]$
-

Algorithm 6 ULA

1. Sample $y_k \sim \pi^{Y|X}(y | x_k) \propto \exp[-\frac{1}{2\eta} \|x_k - y\|^2]$
 2. Sample $x_{k+1} \sim e^{-\langle \nabla f(y_k), x - y_k \rangle - \frac{1}{2\eta} \|x - y_k\|^2} \propto e^{-\frac{1}{2\eta} \|x - (y_k - \eta \nabla f(y_k))\|^2}$
-

$$\begin{aligned}x_{k+1} &= y_k - \eta \nabla f(y_k) + \sqrt{\eta} z_k, & z_k &\sim N(0, I), \\y_{k+1} &= x_{k+1} + \sqrt{\eta} z'_k, & z'_k &\sim N(0, I).\end{aligned}$$

$$\implies y_{k+1} = y_k - \eta \nabla f(y_k) + \sqrt{\eta}(z_k + z'_k) = y_k - \eta \nabla f(y_k) + \sqrt{2\eta} z, \quad z \sim N(0, I)$$

ULA can be viewed as ASF with RGO implemented without rejection

$$h_1(x) = f(y_k) + \langle f'(y_k), x - y_k \rangle + \frac{1}{2\eta} \|x - y_k\|^2 \leq f(x) + \frac{1}{2\eta} \|x - y_k\|^2 = f_{y_k}^\eta(x)$$

Conclusions

- A proximal sampling algorithm for $\nu \propto \exp(-f)$.
 f nonconvex, semi-smooth, composite. ν satisfies either LSI or PI.
- Total complexity $\tilde{\mathcal{O}}\left(C \sum_{i=1}^n L_{\alpha_i}^{\frac{2}{\alpha_i+1}} d\right)$ where $C = C_{\text{LSI}}$ or $C = C_{\text{PI}}$.

Each iteration takes $\tilde{\mathcal{O}}(1)$ subgradients of f and $\mathcal{O}(1)$ samples from Gaussian.

- Inspired by proximal point framework and proximal mapping.
Leverage tools from optimization to design and analyze sampling algorithms.
E.g., acceleration in sampling for weakly smooth potentials.

- Jiaming Liang and Yongxin Chen. A Proximal Algorithm for Sampling from Non-convex Potentials. 2022
- Yongxin Chen, Sinho Chewi, Adil Salim, and Andre Wibisono. Improved Analysis for a Proximal Algorithm for Sampling. COLT 2022
- Ruoqi Shen, Kevin Tian, and Yin Tat Lee. Structured Logconcave Sampling with a Restricted Gaussian Oracle. COLT 2021
- Dao Nguyen, Xin Dang, and Yixin Chen Unadjusted Langevin Algorithm for Non-convex Weakly Smooth Potentials. 2021
- Sinho Chewi, Murat A Erdogdu, Mufan Li, Ruoqi Shen, and Shunshi Zhang. Analysis of Langevin Monte Carlo from Poincare to Log-Sobolev. COLT 2022
- Jiaming Liang and Renato Monteiro. A Proximal Bundle Variant with Optimal Iteration-complexity for A Large Range of Prox Stepsizes. SIAM Journal of Optimization 2021

Thank you!