

P0: CytoSPACE assignments on native Xenium coordinates


Raw CytoSPACE families: every included cell is assigned




Families passing panel, seed, library and count checks



Raw reference groups: exploratory forced assignments




Fine groups passing all checks




Broad identities are supplied as input; subtype labels/fractions are not supplied. Native assignments are compulsory, not confidence-filtered.
Checks: same-stage reference-panel cross-library recoverability, 3 seeds, original counts, each reference library, expression fit and measured marker support.
Gray cells retain their original coordinates. Kcnd2 and Irx5 are unassayed; no measured electrical gradient is shown.

P7: CytoSPACE assignments on native Xenium coordinates


Raw CytoSPACE families: every included cell is assigned




Families passing panel, seed, library and count checks



Raw reference groups: exploratory forced assignments




Fine groups passing all checks



Broad identities are supplied as input; subtype labels/fractions are not supplied. Native assignments are compulsory, not confidence-filtered.
Checks: same-stage reference-panel cross-library recoverability, 3 seeds, original counts, each reference library, expression fit and measured marker support.
Gray cells retain their original coordinates. Kcnd2 and Irx5 are unassayed; no measured electrical gradient is shown.

P14: CytoSPACE assignments on native Xenium coordinates


Raw CytoSPACE families: every included cell is assigned




Families passing panel, seed, library and count checks



Raw reference groups: exploratory forced assignments





Fine groups passing all checks





Broad identities are supplied as input; subtype labels/fractions are not supplied. Native assignments are compulsory, not confidence-filtered.
Checks: same-stage reference-panel cross-library recoverability, 3 seeds, original counts, each reference library, expression fit and measured marker support.
Gray cells retain their original coordinates. Kcnd2 and Irx5 are unassayed; no measured electrical gradient is shown.

P21: CytoSPACE assignments on native Xenium coordinates










Broad identities are supplied as input; subtype labels/fractions are not supplied. Native assignments are compulsory, not confidence-filtered.
Checks: same-stage reference-panel cross-library recoverability, 3 seeds, original counts, each reference library, expression fit and measured marker support.
Gray cells retain their original coordinates. Kcnd2 and Irx5 are unassayed; no measured electrical gradient is shown.

Gene recovery when all listed genes are omitted from CytoSPACE matching



Evaluation is within the fixed ventricular input cohort. Broad input annotations are retained. Grid bins require ≥20 cells.
These are reference-derived predictions compared with measured Xenium genes; they do not validate unassayed Kcnd2 or Irx5.