

P0: individual spatial evidence maps for all 8 ventricular RNA groups

P0_M0 · Cycling




Proposed: 2,912; prioritized: 1,833
Measured supporting genes: Atad2, Top2a, Rad51b

P0_M1 · Ddc / metabolic transition




Proposed: 3,842; prioritized: 0
Measured supporting genes: Cacna1d, Angpt1, Ivns1abp

P0_M2 · Ncam1 / neonatal working




Proposed: 6,366; prioritized: 558
Measured supporting genes: Ncam1, Mapk10, Kcnq5

P0_M3 · Ankrd1 remodeling A




Proposed: 240; prioritized: 0
Measured supporting genes: Ankrd1, Creb5, Itga9

P0_M4 · Fatty-acid-use associated




Proposed: 6,016; prioritized: 0
Measured supporting genes: Eda, Dnmt3a, Cd36

P0_M5 · Conduction-program candidate




Proposed: 1,661; prioritized: 892
Measured supporting genes: Slit2, Sema3a, Thsd4

P0_M6 · Ankrd1 remodeling B




Proposed: 946; prioritized: 0
Measured supporting genes: Ankrd1, Creb5, Acta2

P0_M7 · DNA-maintenance associated



Proposed: 11; prioritized: 0
Measured supporting genes: Pola1, Rad51b, Hells



All original locations are retained. Colors show continuous group weight; black outlines mark prioritized candidate placements.
Measured markers corroborate labels but also contributed to matching. Low weights and absent outlines do not establish biological absence.