






Reviewed RNA families · residual CM/mural groups excluded · original UMAP coordinates
P0: 2,142 retained nuclei
P7: 2,608 retained nuclei
P14: 5,241 retained nuclei
P21: 4,337 retained nuclei
UMAP1
UMAP2
F0
F1
F2
F3
F4
F5
F6
F7
F8
Postn-associated
Col8a1-associated
Gfpt2-associated
Lsamp-associated
Kirrel3-associated
Unresolved fibroblast program


P0 · 1,591/2,319 profiles meet unchanged fine-label rules
Original supported families
Refitted supported families
Refitted proposals: include uncertain profiles
Y (µm)
X (µm)
500
1000
1500
2000
2500
3000
3500
Postn-associated
Col8a1-associated
Gfpt2-associated
Lsamp-associated
Kirrel3-associated
Unresolved fibroblast program


P7 · 5,504/8,564 profiles meet unchanged fine-label rules
Original supported families
Refitted supported families
Refitted proposals: include uncertain profiles
Y (µm)
X (µm)
500
1000
1500
2000
2500
3000
3500
4000
1000
2000
3000
4000
5000
Postn-associated
Col8a1-associated
Gfpt2-associated
Lsamp-associated
Kirrel3-associated
Unresolved fibroblast program


P14 · 1,343/9,824 profiles meet unchanged fine-label rules
Original supported families
Refitted supported families
Refitted proposals: include uncertain profiles
Y (µm)
X (µm)
1000
2000
3000
4000
5000
6000
Postn-associated
Col8a1-associated
Gfpt2-associated
Lsamp-associated
Kirrel3-associated
Unresolved fibroblast program


P21 · 1,497/13,083 profiles meet unchanged fine-label rules
Original supported families
Refitted supported families
Refitted proposals: include uncertain profiles
Y (µm)
X (µm)
1000
2000
3000
4000
5000
6000
Postn-associated
Col8a1-associated
Gfpt2-associated
Lsamp-associated
Kirrel3-associated
Unresolved fibroblast program


P21 · Reviewed family weights at original Xenium positions
Col8a1-associated
Lsamp-associated
Gfpt2-associated
Kirrel3-associated
0
1000
2000
3000
4000
5000
6000
1000
2000
3000
4000
5000
X (µm)
Reference similarity weight (not probability)
0.0
0.2
0.4
0.6
0.8
1.0