

P14_arterial_1: Peri-arterial



10 arterial EC + 7 SMC seeds
RNA points sampled to≤5,000 per marker group for display; all molecules retained in data.
Alternative: fixed original cell seeds, including anucleate profiles; boundary stain only. This is a sensitivity test, not validated replacement segmentation.

P14_arterial_2: Peri-arterial




8 arterial EC + 14 SMC seeds

RNA points sampled to≤5,000 per marker group for display; all molecules retained in data.

Alternative: fixed original cell seeds, including anucleate profiles; boundary stain only. This is a sensitivity test, not validated replacement segmentation.

P14_Endothelial_ordinary: CM–Endothelial interface




mnolakpo-1: Endothelial; nuclear area 34.6 µm²

Every displayed marker molecule retains its original position; no expression smoothing.

Alternative: fixed original cell seeds, including anucleate profiles; boundary stain only. This is a sensitivity test, not validated replacement segmentation.

P14_Endothelial_flagged: CM–Endothelial interface




ebpgjecn-1: Mixed-marker review; nuclear area 56.7 µm²

Every displayed marker molecule retains its original position; no expression smoothing.

Alternative: fixed original cell seeds, including anucleate profiles; boundary stain only. This is a sensitivity test, not validated replacement segmentation.

P14_Fibroblast_ordinary: CM–Fibroblast interface




llkjcbje-1: Fibroblast; nuclear area 53.8 µm²

Every displayed marker molecule retains its original position; no expression smoothing.

Alternative: fixed original cell seeds, including anucleate profiles; boundary stain only. This is a sensitivity test, not validated replacement segmentation.

P14_Fibroblast_flagged: CM–Fibroblast interface




kaeimpih-1: Mixed-marker review; nuclear area 8.4 µm²

Every displayed marker molecule retains its original position; no expression smoothing.

Alternative: fixed original cell seeds, including anucleate profiles; boundary stain only. This is a sensitivity test, not validated replacement segmentation.

P14_anucleate_CM_1: CM without measured nuclear area




ebdlgbbf-1: Cardiomyocyte; nuclear area not measured

Every displayed marker molecule retains its original position; no expression smoothing.

Alternative: fixed original cell seeds, including anucleate profiles; boundary stain only. This is a sensitivity test, not validated replacement segmentation.

P14_anucleate_CM_2: CM without measured nuclear area



dglmgbmm-1: Cardiomyocyte; nuclear area not measured

Every displayed marker molecule retains its original position; no expression smoothing.

Alternative: fixed original cell seeds, including anucleate profiles; boundary stain only. This is a sensitivity test, not validated replacement segmentation.