
P21_arterial_1: Peri-arterial




2 arterial EC + 8 SMC seeds

RNA points sampled to≤5,000 per marker group for display; all molecules retained in data.

Alternative: fixed original cell seeds, including anucleate profiles; boundary stain only. This is a sensitivity test, not validated replacement segmentation.

P21_arterial_2: Peri-arterial




2 arterial EC + 11 SMC seeds

RNA points sampled to≤5,000 per marker group for display; all molecules retained in data.

Alternative: fixed original cell seeds, including anucleate profiles; boundary stain only. This is a sensitivity test, not validated replacement segmentation.

P21_Endothelial_ordinary: CM–Endothelial interface




ieoplmmo-1: Endothelial; nuclear area 18.9 µm²

Every displayed marker molecule retains its original position; no expression smoothing.

Alternative: fixed original cell seeds, including anucleate profiles; boundary stain only. This is a sensitivity test, not validated replacement segmentation.

P21_Endothelial_flagged: CM–Endothelial interface




jhmmjcpa-1: Mixed-marker review; nuclear area 9.8 µm²

Every displayed marker molecule retains its original position; no expression smoothing.

Alternative: fixed original cell seeds, including anucleate profiles; boundary stain only. This is a sensitivity test, not validated replacement segmentation.

P21_Fibroblast_ordinary: CM–Fibroblast interface




njagdbbl-1: Fibroblast; nuclear area 33.5 µm²

Every displayed marker molecule retains its original position; no expression smoothing.

Alternative: fixed original cell seeds, including anucleate profiles; boundary stain only. This is a sensitivity test, not validated replacement segmentation.

P21_Fibroblast_flagged: CM–Fibroblast interface




bglabdbe-1: Mixed-marker review; nuclear area 33.1 µm²

Every displayed marker molecule retains its original position; no expression smoothing.

Alternative: fixed original cell seeds, including anucleate profiles; boundary stain only. This is a sensitivity test, not validated replacement segmentation.

P21_anucleate_CM_1: CM without measured nuclear area




akpncpcc-1: Cardiomyocyte; nuclear area not measured

Every displayed marker molecule retains its original position; no expression smoothing.

Alternative: fixed original cell seeds, including anucleate profiles; boundary stain only. This is a sensitivity test, not validated replacement segmentation.

P21_anucleate_CM_2: CM without measured nuclear area



alhefcgg-1: Cardiomyocyte; nuclear area not measured

Every displayed marker molecule retains its original position; no expression smoothing.

Alternative: fixed original cell seeds, including anucleate profiles; boundary stain only. This is a sensitivity test, not validated replacement segmentation.