

P7_arterial_1: Peri-arterial




4 arterial EC + 3 SMC seeds
RNA points sampled to≤5,000 per marker group for display; all molecules retained in data.
Alternative: fixed original cell seeds, including anucleate profiles; boundary stain only. This is a sensitivity test, not validated replacement segmentation.

P7_arterial_2: Peri-arterial



19 arterial EC + 20 SMC seeds
RNA points sampled to≤5,000 per marker group for display; all molecules retained in data.
Alternative: fixed original cell seeds, including anucleate profiles; boundary stain only. This is a sensitivity test, not validated replacement segmentation.

P7_Endothelial_ordinary: CM–Endothelial interface




kkiagnpi-1: Endothelial; nuclear area 50.1 µm²

Every displayed marker molecule retains its original position; no expression smoothing.

Alternative: fixed original cell seeds, including anucleate profiles; boundary stain only. This is a sensitivity test, not validated replacement segmentation.

P7_Endothelial_flagged: CM–Endothelial interface




dmkempck-1: Mixed-marker review; nuclear area 50.7 µm²

Every displayed marker molecule retains its original position; no expression smoothing.

Alternative: fixed original cell seeds, including anucleate profiles; boundary stain only. This is a sensitivity test, not validated replacement segmentation.

P7_Fibroblast_ordinary: CM–Fibroblast interface




ibomidji-1: Fibroblast; nuclear area 56.6 µm²

Every displayed marker molecule retains its original position; no expression smoothing.

Alternative: fixed original cell seeds, including anucleate profiles; boundary stain only. This is a sensitivity test, not validated replacement segmentation.

P7_Fibroblast_flagged: CM–Fibroblast interface




kbeappgd-1: Mixed-marker review; nuclear area 35.7 µm²

Every displayed marker molecule retains its original position; no expression smoothing.

Alternative: fixed original cell seeds, including anucleate profiles; boundary stain only. This is a sensitivity test, not validated replacement segmentation.

P7_anucleate_CM_1: CM without measured nuclear area




facpamoa-1: Cardiomyocyte; nuclear area not measured

Every displayed marker molecule retains its original position; no expression smoothing.

Alternative: fixed original cell seeds, including anucleate profiles; boundary stain only. This is a sensitivity test, not validated replacement segmentation.

P7_anucleate_CM_2: CM without measured nuclear area



ckoibenf-1: Cardiomyocyte; nuclear area not measured

Every displayed marker molecule retains its original position; no expression smoothing.

Alternative: fixed original cell seeds, including anucleate profiles; boundary stain only. This is a sensitivity test, not validated replacement segmentation.