Broad-identity stability depends on boundaries and measurement depth

Boundary/QV changes and count-matched controls

	Core >1 µm	Random: core1 depth	Core >2 µm	Random: core2 depth	Add unassigned ≤2 µm	Native Q30	Random: Q30 depth
P0 · CM (n=5864)	86%	86%	54%	56%	98%	92%	95%
P0 · EC (n=1312)	77%	73%	49%	40%	93%	78%	79%
P0 · Fib (n=491)	68%	62%	35%	28%	95%	64%	68%
P0 · SMC (n=35)	57%	46%	29%	23%	91%	57%	57%
P7 · CM (n=7025)	91%	91%	66%	66%	98%	94%	96%
P7 · EC (n=1541)	68%	57%	33%	19%	94%	71%	71%
P7 · Fib (n=470)	64%	59%	32%	24%	94%	56%	65%
P7 · SMC (n=44)	50%	48%	20%	9%	98%	52%	52%
P14 · CM (n=5032)	95%	95%	83%	83%	97%	98%	98%
P14 · EC (n=1284)	70%	62%	40%	27%	90%	81%	81%
P14 · Fib (n=437)	69%	65%	40%	36%	88%	78%	81%
P14 · SMC (n=122)	73%	70%	43%	38%	90%	71%	82%
P21 · CM (n=3212)	91%	92%	76%	76%	97%	97%	97%
P21 · EC (n=1136)	65%	56%	32%	21%	85%	79%	81%
P21 · Fib (n=661)	69%	69%	42%	36%	92%	84%	84%
P21 · SMC (n=71)	63%	63%	30%	25%	85%	79%	83%

Image-covered cells: boundary alternatives

	Native raster control	Centroid Voronoi	Membrane watershed
P0 · CM (n=3422)	99%	93%	86%
P0 · EC (n=833)	95%	87%	71%
P0 · Fib (n=341)	93%	87%	64%
P0 · SMC (n=25)	88%	84%	48%
P7 · CM (n=3393)	99%	95%	89%
P7 · EC (n=1009)	95%	83%	67%
P7 · Fib (n=294)	94%	78%	54%
P7 · SMC (n=38)	92%	76%	55%
P14 · CM (n=2076)	99%	95%	88%
P14 · EC (n=854)	94%	78%	60%
P14 · Fib (n=312)	94%	76%	55%
P14 · SMC (n=95)	93%	80%	63%
P21 · CM (n=1613)	98%	95%	86%
P21 · EC (n=695)	94%	64%	52%
P21 · Fib (n=415)	96%	80%	64%
P21 · SMC (n=46)	93%	72%	59%

Fraction retaining the original broad label

Label retention is not an accuracy estimate or segmentation-error rate. Lost labels include uncertainty and insufficient RNA, not only lineage switches.
Random controls sample original transcripts without replacement to exactly match each cell's eroded/Q30 transcript total (one fixed technical draw).
The two panels use different cohorts; image comparisons require adequate crop coverage. All original seeds, including anucleate profiles, are retained.