
P0_arterial_1: Peri-arterial




7 arterial EC + 7 SMC seeds

RNA points sampled to≤5,000 per marker group for display; all molecules retained in data.


Alternative: fixed original cell seeds, including anucleate profiles; boundary stain only. This is a sensitivity test, not validated replacement segmentation.

P0_arterial_2: Peri-arterial



11 arterial EC + 8 SMC seeds
RNA points sampled to≤5,000 per marker group for display; all molecules retained in data.
Alternative: fixed original cell seeds, including anucleate profiles; boundary stain only. This is a sensitivity test, not validated replacement segmentation.

P0_Endothelial_ordinary: CM–Endothelial interface




iknnhlmp-1: Endothelial; nuclear area 33.4 µm²

Every displayed marker molecule retains its original position; no expression smoothing.

Alternative: fixed original cell seeds, including anucleate profiles; boundary stain only. This is a sensitivity test, not validated replacement segmentation.

P0_Endothelial_flagged: CM–Endothelial interface




abnfoiao-1: Mixed-marker review; nuclear area 28.2 µm²

Every displayed marker molecule retains its original position; no expression smoothing.

Alternative: fixed original cell seeds, including anucleate profiles; boundary stain only. This is a sensitivity test, not validated replacement segmentation.

P0_Fibroblast_ordinary: CM–Fibroblast interface




ieajpahp-1: Fibroblast; nuclear area 36.6 µm²

Every displayed marker molecule retains its original position; no expression smoothing.

Alternative: fixed original cell seeds, including anucleate profiles; boundary stain only. This is a sensitivity test, not validated replacement segmentation.

P0_Fibroblast_flagged: CM–Fibroblast interface




mebhojgo-1: Mixed-marker review; nuclear area 17.6 µm²

Every displayed marker molecule retains its original position; no expression smoothing.

Alternative: fixed original cell seeds, including anucleate profiles; boundary stain only. This is a sensitivity test, not validated replacement segmentation.

P0_anucleate_CM_1: CM without measured nuclear area




bbcnoama-1: Cardiomyocyte; nuclear area not measured

Every displayed marker molecule retains its original position; no expression smoothing.

Alternative: fixed original cell seeds, including anucleate profiles; boundary stain only. This is a sensitivity test, not validated replacement segmentation.

P0_anucleate_CM_2: CM without measured nuclear area



bplledck-1: Cardiomyocyte; nuclear area not measured

Every displayed marker molecule retains its original position; no expression smoothing.

Alternative: fixed original cell seeds, including anucleate profiles; boundary stain only. This is a sensitivity test, not validated replacement segmentation.