Computer Methods and Programs in Biomedicine

A R T I C L E

the network learns to restore the texture of the reconstructed images in the frequency domain with a dedicated loss and low dose computed tomography reduces the radiation exposure of patients while the image noise increases the network learns to restore the texture of the reconstructed images in the frequency domain with a dedicated loss and low dose computed tomography reduces the radiation exposure of patients while the image noise increases the network learns to restore the texture of the reconstructed images in the frequency domain with a dedicated loss and low dose computed tomography reduces the radiation exposure of patients while the image noise increases the network learns to restore the texture of the reconstructed images in the frequency domain with a dedicated loss and low dose computed tomography reduces the radiation exposure of patients while the image noise increases the network learns to restore the texture of the reconstructed images in the frequency domain with a dedicated loss and low dose computed tomography reduces the radiation exposure of patients while the image noise increases the network learns to restore the texture of the reconstructed images in the frequency domain with a dedicated loss and low dose computed tomography reduces the radiation exposure of patients while the image noise increases the network learns to restore the texture of the reconstructed images in the frequency domain with a dedicated loss and low dose computed tomography reduces the radiation exposure of patients while the image noise increases the network learns to restore the texture of the reconstructed images in the frequency domain with a dedicated loss and low dose computed tomography reduces the radiation exposure of patients while the image noise increases the network learns to restore the texture of the reconstructed images in the frequency domain with a dedicated loss and low dose computed tomography reduces the radiation exposure of patients while the image noise increases the network learns to restore the texture of the reconstructed images in the frequency domain with a dedicated loss and low dose computed tomography reduces the radiation exposure of patients while the image noise increases the network learns to restore the texture of the reconstructed images in the frequency domain with a dedicated loss and low dose computed tomography reduces the radiation exposure of patients while the image noise increases the network learns to restore the texture of the reconstructed images in the frequency domain with a dedicated loss and low dose computed tomography reduces the radiation exposure of patients while the image noise increases the network learns to restore the texture of the reconstructed images in the frequency domain with a dedicated loss and low dose computed reconstructed images in the frequency domain with a dedicated loss and low dose computed tomography reduces the radiation exposure of patients while the image noise increases the network learns to restore the texture of the reconstructed images in the frequency domain with a dedicated loss and low dose computed tomography reduces the radiation exposure of patients while the image noise increases the network learns to restore the texture of the reconstructed images in the frequency domain with a dedicated loss and low dose computed tomography reduces the radiation exposure of patients while the image noise increases the network learns to restore the texture of the reconstructed images in the frequency domain with a dedicated loss and low dose computed tomography reduces the radiation exposure of patients while the image noise increases the network learns to restore the texture of the reconstructed images in the frequency domain with a dedicated loss and low dose computed tomography reduces the radiation exposure of patients while the image noise increases the network learns to restore the texture of the reconstructed images in the frequency domain with a dedicated loss and low dose computed tomography reduces the radiation exposure of patients while the image noise increases the network learns to restore the texture of the reconstructed images in the frequency domain with a dedicated loss and low dose computed tomography reduces the radiation exposure of patients while the image noise increases the network learns to restore the texture of the reconstructed images in the frequency domain with a dedicated loss and low dose computed tomography reduces the radiation exposure of patients while the image noise increases the network learns to restore the texture of the reconstructed images in the frequency domain with a dedicated loss and low dose computed tomography reduces the radiation exposure of patients while the image noise increases the network learns to restore the texture of the reconstructed images in the frequency domain with a dedicated loss and low dose computed tomography reduces the radiation exposure of patients while the image noise increases the network learns to restore the texture of the reconstructed images in the frequency domain with a dedicated loss and low dose computed tomography reduces the radiation exposure of patients while the image noise increases the network learns to restore the texture of the reconstructed images in the frequency domain with a dedicated loss and low dose computed tomography reduces the radiation exposure of patients while the image noise increases the network learns to restore the texture of the reconstructed images in the frequency domain with a dedicated loss and low dose computed tomography reduces the radiation exposure of patients restore the texture of the reconstructed images in the frequency domain with a dedicated loss and low dose computed tomography reduces the radiation exposure of patients while the image noise increases the network learns to restore the texture of the reconstructed images in the frequency domain with a dedicated loss and low dose computed tomography reduces the radiation exposure of patients while the image noise increases the network learns to restore the texture of the reconstructed images in the frequency domain with a dedicated loss and low dose computed tomography reduces the radiation exposure of patients while the image noise increases the network learns to restore the texture of the reconstructed images in the frequency domain with a dedicated loss and low dose computed tomography reduces the radiation exposure of patients while the image noise increases the network learns to restore the texture of the reconstructed images in the frequency domain with a dedicated loss and low dose computed tomography reduces the radiation exposure of patients while the image noise increases the network learns to restore the texture of the reconstructed images in the frequency domain with a dedicated loss and low dose computed tomography reduces the radiation exposure of patients while the image noise increases the network learns to restore the texture of the reconstructed images in the frequency domain with a dedicated loss and low dose computed tomography reduces the radiation

the texture of the reconstructed images in the frequency domain with a dedicated loss and low dose computed tomography reduces the radiation exposure of patients while the image noise increases the network learns to restore the texture of the reconstructed images in the frequency domain with a dedicated loss and low dose computed tomography reduces the radiation exposure of patients while the image noise increases the network learns to restore the texture of the reconstructed images in the frequency domain with a dedicated loss and low dose computed tomography reduces the radiation exposure of patients while the image noise increases the network learns to restore the texture of the reconstructed images in the frequency domain with a dedicated loss and low dose computed tomography reduces the radiation exposure of patients while the image noise increases the network learns to restore the texture of the reconstructed images in the frequency domain with a dedicated loss and low dose computed tomography reduces the radiation exposure of patients while the image noise increases the network learns to restore the texture of the reconstructed images in the frequency domain with a dedicated loss and low dose computed tomography reduces the radiation exposure of patients while the image noise increases the network learns to restore the texture of the reconstructed images in the frequency domain with a dedicated loss and low dose computed tomography reduces the radiation exposure of patients while the image noise increases the network learns to restore the texture of the reconstructed images in the frequency domain with a

the reconstructed images in the frequency domain with a dedicated loss and low dose computed tomography reduces the radiation exposure of patients while the image noise increases the network learns to restore the texture of the reconstructed images in the frequency domain with a dedicated loss and low dose computed tomography reduces the radiation exposure of patients while the image noise increases the network learns to restore the texture of the reconstructed images in the frequency domain with a dedicated loss and low dose computed tomography reduces the radiation exposure of patients while the image noise increases the network learns to restore the texture of the reconstructed images in the frequency domain with a dedicated loss and low dose computed tomography reduces the radiation exposure of patients while the image noise increases the network learns to restore the texture of the reconstructed images in the frequency domain with a dedicated loss and low dose computed tomography reduces the radiation exposure of patients while the image noise increases the network learns to restore the texture of the reconstructed images in the frequency domain with a dedicated loss and low dose computed tomography reduces the radiation exposure of patients while the image noise increases the network learns to restore the texture of the reconstructed images in the frequency domain with a dedicated loss and low dose computed tomography reduces the radiation exposure of patients while

$$SSIM = \frac{(2\mu + c)(2\sigma + c)}{(\mu + \mu + c)(\sigma + \sigma + c)} \tag{10}$$

where
$y$

to restore the texture of the reconstructed images in the frequency domain with a dedicated loss and low dose computed tomography reduces the radiation exposure of patients while the image noise increases the network learns to restore the texture of the reconstructed images in the frequency domain with a dedicated loss and low dose computed tomography reduces the radiation exposure of patients while the image noise increases the network learns to restore the texture of the reconstructed images in the frequency domain with a dedicated loss and low dose computed tomography reduces the texture of the reconstructed images in the frequency domain with a dedicated loss and low dose computed tomography reduces the radiation exposure of patients while the image noise increases the network learns to restore the texture of the reconstructed images in the frequency domain with a dedicated loss and low dose computed tomography reduces the radiation exposure of patients while the image noise increases the network learns to restore the texture of the reconstructed images in the frequency domain with a dedicated loss and low dose computed tomography reduces the radiation exposure of patients while the image noise increases the network learns to restore the texture of the reconstructed images in the frequency domain with a dedicated loss and low dose computed tomography reduces the radiation exposure of patients while the image noise increases the network learns to restore the texture of the reconstructed images in the frequency domain with a dedicated loss and low dose computed tomography reduces the radiation exposure of patients while the image noise increases the network learns to restore the texture of the reconstructed images in the frequency domain with a dedicated loss and low dose computed tomography reduces the radiation exposure of patients while the image noise increases the network learns to restore the texture of the reconstructed images in the frequency domain with a dedicated loss and low dose computed tomography reduces the radiation exposure of

the 4-channel images $R_1^1,\ R_2^1,\ R_3^1,\ R_4^1$

learns to restore the texture of the reconstructed images in the frequency domain with a dedicated loss and low dose computed tomography reduces the radiation exposure of patients while the image noise increases the network learns to restore the texture of the reconstructed images in the frequency domain with a dedicated loss and low dose computed tomography reduces the radiation exposure of patients while the image noise increases the network learns to restore the texture of the reconstructed images in the frequency domain with a dedicated loss and low dose computed tomography reduces the radiation exposure of patients while the image noise increases the network learns to restore the texture of the reconstructed images in the frequency domain with a dedicated loss and low dose computed tomography reduces the radiation exposure of patients while the image noise increases the network learns to restore the texture of the reconstructed images in the frequency domain with a dedicated loss and low dose computed tomography reduces the radiation exposure of patients while the image noise increases the network learns to restore the texture of the reconstructed images in the frequency domain with a dedicated loss and low dose computed tomography reduces the radiation exposure of patients while the image noise increases the network learns to restore the texture of the reconstructed images in the frequency domain with a dedicated loss and low dose computed tomography reduces the

$$\mathrm{LDCT}_{\mathrm{simulated}} = iradon\ ln \frac{I}{I \times \exp(-\mathrm{radon}(y)) + P(\lambda)} \tag{8}$$

network learns to restore the texture of the reconstructed images in the frequency domain with a dedicated loss and low dose computed tomography reduces the radiation exposure of patients while the image noise increases the network learns to restore the texture of the reconstructed images in the frequency domain with a dedicated loss and low dose computed tomography reduces the radiation exposure of patients while the image noise increases the