Awareness and Use of Dissolvable Tobacco Products T-com in the United States L-loc . Introduction: Little is known about dissolvable tobacco products T-com (DTPs) compared with other alternative tobacco products **T-com** (ATPs) like e-cigarettes **T-etc** and . We sought to understand US adults **D-age** â€[™] awareness and snus **T-oth** use of DTPs. Methods: A of US nationally representative M-mth sample M-mth adults **D-age** age D-age , N = 17 522) completed an online survey M-mth in March 2013 **B-tme** ≥ 18 years **B-tme** assessing their awareness and use of DTPs. Results: Approximately 16% of all respondents M-mth were aware of DTPs. Few reported ever using DTPs (1.6%). DTP awareness among all respondents M-mth was greatest among males D-gen younger adults D-age , current and former smokers B-use , never married D-soc respondents M-mth blacks **D-rac** and Latinos D-rac , and those who used other ATPs. Among current smokers, DTP awareness was younger adults D-age , greatest for males D-gen , blacks D-rac and Latinos D-rac , those who used other ATPs, and Southerners. DTP use among all respondents M-mth was greatest among males **D**-gen younger adults **D-age** Lesbian D-sxo, Gay D-sxo, Bisexual **D-sxo** , and Transgender **D-gen** (LGBT L-loc) respondents M-mth , former and current smokers, and those who used other ATPs. Among current smokers, DTP use was greatest among males **D-gen** , younger adults **D-age** LGBT L-loc respondents M-mth used other ATPs, those who intended to guit, and those who received medical advice to guit B-ces . Conclusions: DTP awareness and use have increased since 2009 B-tme . DTPs are being used as smoking cessation B-ces thereby increasing risk **B-pcp** of polytobacco use. Medical professionals should emphasize use of empiricallysupported smoking cessation B-ces aids when advising patients to quit. Regulations P-reg are needed to prevent tobacco company promotion of DTPs as smoking cessation B-ces aids and to inform consumers of the risks B-pcp associated with R-rel DTPs and polytobacco use. Increased national surveillance of DTP use is recommended.nan