Smoking Cessation B-ces Awareness and Utilization Among Lesbian **D-sxo** Gay **D-sxo** Bisexual **D-sxo** and Transgender D-gen Adults D-age : An Analysis of the 2009â€"2010 B-tme National Adult Tobacco Survey M-dat . Introduction: Each year, there are more than R-rel 480 000 deaths in the United States L-loc Lesbian **D-sxo** bisexual **D-sxo** LGBT L-loc attributed to smoking. gay **D-sxo** and transgender **D**-gen (adults **D**-age are a vulnerable population that smokes at higher rates **B**-prv than heterosexuals D-sxo Methods: We used data collected from the National Adult Tobacco Survey M-dat 2009â€"2010 **B-tme** , a large, nationally representative **M-mth** study using a randomized M-mth of US landline and , national sample M-mth LGBT L-loc cellular telephone listings, (N = 118590). We compared adults **D-age** heterosexual D-sxo to their counterparts with regard to exposure B-exp to advertisements P-mkt promoting smoking cessation **B-ces**, and awareness and use of tobacco treatment P-trt services, including quitlines P-trt smoking cessation B-ces classes, health professional counseling P-trt , nicotine replacement therapy P-trt , and medications P-trt Results: Fewer GBT men **D-gen** , compared to heterosexual **D-sxo** men **D-gen** , were aware of the quitline LGBT L-loc individuals have similar exposure B-exp P-trt . However, to tobacco cessation B-ces advertising P-mkt, as well as similar awareness of and use of evidence based cessation B-ces methods as compared to heterosexual **D-sxo** peers. Conclusions: The similarly of awareness and use of cessation **B-ces** support indicates a need for LGBT L-loc -specific efforts to reduce smoking disparities. Potential interventions Ptrt would include: improving awareness of, access to and acceptability of current cessation B-ces methods for LGBT L-loc patients, developing tailored cessation B-ces interventions P-trt, and denormalizing P-mkt smoking in LGBT L-loc community spaces.nan