CS3243 : Introduction to Artificial Intelligence

Tutorial 9

NUS School of Computing

April 3, 2023

Admin

► Project 3

- Basics of Probability
- Independence
- ► Conditional Independence
- ► Bayes' Rule

- Bayesian Networks
- Determining independence between two nodes given a Bayesian Network
- ▶ Why? Simplify/reduce the size of our probability calculations!


- Bayesian Networks
- Determining independence between two nodes given a Bayesian Network
- Why? Simplify/reduce the size of our probability calculations!
- ► How?
- Constructing a Bayesian network from cause and effect relationships "Domain knowledge" or simply specified by the question
- Factorising the joint probability distribution given the Bayesian network
- Computing joint probabilities (or other conditional probabilities) using the (smaller) CPTs and probability laws (example chain rule)

- ▶ A Bayesian Network is a probabilistic graphical model that represents conditional dependencies between random variables through a Directed Acyclic Graph (DAG)
- ▶ The graph consists of nodes (variables) and edges $(X \to Y)$ means that X directly influences Y)
- We also include conditional distribution for each node given its parents: $Pr\{X|parents(X)\}$
- Conditional distribution can be represented as a CPT
- ▶ The distribution over X for each combination of parent values:
- ▶ Given $X_1, X_2, ..., X_n$: $Pr\{X_1 \wedge X_2 \wedge ... \wedge X_n\} = \prod_i Pr\{X_i | parents(X_i)\} \rightarrow \mathsf{Chain}$ rule!

- Given
- ightharpoonup G : Good Grades
- ightharpoonup C: Good Communication Skills
- $Pr{G = 1} = 0.7$
- $Pr\{C=1\}=0.2$

G	C	$\Pr\{I=1\mid G,C\}$
1	1	0.9
1	0	0.5
0	1	0.5
0	0	0.1

► The network



▶ What is the probability that Alice, who has poor grades and communication skills, performs well on her interview?

- ▶ What is the probability that Alice, who has poor grades and communication skills, performs well on her interview?
- $ightharpoonup Pr\{I=1|G=0,C=0\}$

- ▶ What is the probability that Alice, who has poor grades and communication skills, performs well on her interview?
- $ightharpoonup Pr\{I=1|G=0,C=0\}$
- Quite straightforward right?
- 0.1 (from the table itself)

- ▶ What is the probability that Alice, who has poor grades and communication skills, performs well on her interview?
- $ightharpoonup Pr\{I=1|G=0,C=0\}$
- Quite straightforward right?
- ▶ 0.1 (from the table itself)
- Extra Practice
- ▶ What do you think is the value of $Pr\{I=1, G=0, C=0\}$?

What is the probability that Bob is a student with great communication skills, assuming we do not know anything about him?

- What is the probability that Bob is a student with great communication skills, assuming we do not know anything about him?
- ▶ $Pr\{C=1\}$

- What is the probability that Bob is a student with great communication skills, assuming we do not know anything about him?
- ▶ $Pr\{C=1\}$
- ▶ 0.2 (given)

What is the probability that a student has good communication skills, given that he or she has performed well in an interview? Are good communication skills independent of good performance in an interview?

- What is the probability that a student has good communication skills, given that he or she has performed well in an interview? Are good communication skills independent of good performance in an interview?
- $Pr\{C=1|I=1\}$

- What is the probability that a student has good communication skills, given that he or she has performed well in an interview? Are good communication skills independent of good performance in an interview?
- $Pr\{C=1|I=1\}$
- $Pr\{C=1,I=1\}$ $Pr\{I=1\}$
- Apply Bayes' Rule

- Given
- ▶ Pr[the kit shows positive | the patient is a carrier] = 0.998
- $ightharpoonup Pr[{\it the kit shows negative} \mid {\it the patient is not a carrier}] = 0.996$

- Given
- ▶ Pr[the kit shows positive | the patient is a carrier] = 0.998
- $ightharpoonup Pr[{\it the kit shows negative} \mid {\it the patient is not a carrier}] = 0.996$
- Let's define some variables
- $\blacktriangleright X/\neg X$: Positive/negative test result
- $ightharpoonup Y/\neg Y$: Patient is/isn't a carrier

- ▶ Based on the new variables, let's rewrite the probabilities
- $Pr{Y} = 0.02$
- $ightharpoonup Pr\{X|Y\} = 0.998$
- $Pr\{\neg X|\neg Y\} = 0.996$

▶ Given that a patient is tested to be positive using this kit, what is the posterior belief that he is not a carrier? Give your answer to 3 decimal places.

- Given that a patient is tested to be positive using this kit, what is the posterior belief that he is not a carrier? Give your answer to 3 decimal places.
- Posterior belief : Probability of the parameters θ given the evidence Z, in other terms, simply $Pr\{\theta|Z\}$
- In this question, we need to find posterior probability that patient is not a carrier given positive test results, ie $Pr\{\neg Y|X\}$

- Given that a patient is tested to be positive using this kit, what is the posterior belief that he is not a carrier? Give your answer to 3 decimal places.
- Posterior belief : Probability of the parameters θ given the evidence Z, in other terms, simply $Pr\{\theta|Z\}$
- In this question, we need to find posterior probability that patient is not a carrier given positive test results, ie $Pr\{\neg Y|X\}$
- Apply Bayes' Rule
- $ightharpoonup Pr\{\neg Y|X\}$

Suppose that the patient doesn't entirely trust the result offered by the first kit (perhaps because it has expired) and decides to use another test kit. If the patient is again tested to be positive using this second kit, what is the (updated) likelihood that he is not a carrier? You can assume conditional independence between results of different test kits given the patient's state of virus contraction. Give your answer to 4 decimal places.

- Suppose that the patient doesn't entirely trust the result offered by the first kit (perhaps because it has expired) and decides to use another test kit. If the patient is again tested to be positive using this second kit, what is the (updated) likelihood that he is not a carrier? You can assume conditional independence between results of different test kits given the patient's state of virus contraction. Give your answer to 4 decimal places.
- ▶ We need to find the probability that patient is not a carrier given two positive test results, ie $Pr\{\neg Y|X_1 \land X_2\}$
- ► Apply Bayes' Rule
- $Pr\{\neg Y \land X_1 \land X_2\} \over Pr\{X_1 \land X_2\}$

► Constructing a Bayesian Network from given CPT

$P(\text{Wet_Grass} \mid \text{Sprinkler} \land \text{Rain})$	0.95	
$P(\text{Wet_Grass} \mid \text{Sprinkler} \land \neg \text{Rain})$	0.9	
$P(\text{Wet_Grass} \mid \neg \text{Sprinkler} \land \text{Rain})$	0.8	
$P(\text{Wet_Grass} \mid \neg \text{Sprinkler} \land \neg \text{Rain})$	0.1	
P(Sprinkler Rainy_Season)		
P(Sprinkler ¬ Rainy_Season)		
P(Rain Rainy_Season)		
$P(Rain \mid \neg Rainy_Season)$		
P(Rainy_Season)		

- One key observation to reduce the number of nodes in the tree
- Observe the variables, Sprinkler and Rainy Season
- ► The conditional probabilities between them tell something about their relationship

- One key observation to reduce the number of nodes in the tree
- ▶ Observe the variables, Sprinkler and $Rainy\ Season$
- ► The conditional probabilities between them tell something about their relationship
- ▶ Given two random variables A and B, if $Pr\{A|B\} = 0$ and $Pr\{A|\neg B\} = 1$, A and B are actually complementing variables
- Let's look at the tree!

Thank you!

If you have any questions, please don't hesitate. Feel free to ask! We are here to learn together!