A0214561 M

Zhuang Jianning

TE6

$$\begin{bmatrix}
1 & 1 & 2 & -1 & | & 1 & | & 2 & -1 & | & 1 \\
1 & 0 & 0 & 2 & | & 1 & | & -7 & | & -1 & 2 & -1 & | & 1 \\
0 & 1 & 2 & -2 & | & 2 & | & 2 & | & 2 & | & 2
\end{bmatrix}$$

$$\begin{bmatrix}
2c+R_2 \\
-7 \\
0-1-2 \\
0 & 0 & 1
\end{bmatrix}
\begin{bmatrix}
1 & 2 & -1 & | & 1 \\
0 & -1-2 \\
0 & 0 & 0 & 1
\end{bmatrix}
\begin{bmatrix}
1 & 0 & 2 & | & 1 \\
0 & -1-2 \\
0 & 0 & 0 & 1
\end{bmatrix}
\begin{bmatrix}
1 & 0 & 0 & 2 \\
0 & -1-2 \\
0 & 0 & 0 & 1
\end{bmatrix}
\begin{bmatrix}
1 & 0 & 0 & 2 \\
0 & -1-2 \\
0 & 0 & 0 & 1
\end{bmatrix}$$

$$x_4 = 2$$

Let  $x_1 = 5$ ,  $5 \in \mathbb{R}^2$ 
 $-x_2 - 2x_3 + 3x_4 = 0 = 0$ 
 $x_1 + 2x_4 = 1 = 0$ 
 $x_1 = -3$ 

$$\begin{pmatrix} \chi_1 \\ \chi_2 \\ \chi_3 \\ \chi_4 \end{pmatrix} = \begin{pmatrix} -3 \\ 6 \\ 0 \\ 2 \end{pmatrix} + 5 \begin{pmatrix} 0 \\ -2 \\ 1 \\ 0 \end{pmatrix}, SEIR$$

$$44 = 0$$
  
Let  $x_3 = 5$ ,  $5 \in \mathbb{R}^2$   
 $-x_2 - 2x_3 - 13x_4 = -1 = 0$   $x_1 = -1$   
 $x_1 + 2x_4 = -1 = 0$   $x_1 = -1$ 

$$\begin{pmatrix} \chi_1 \\ \chi_2 \\ \chi_1 \end{pmatrix} = \begin{pmatrix} -1 \\ 1 \\ 0 \\ 0 \end{pmatrix} + 5 \begin{pmatrix} 0 \\ -1 \\ 1 \\ 0 \end{pmatrix} , SEIR$$

Zhuang Jianning

TEL

$$A^{-1} \begin{bmatrix} A & 1 & 1 \\ -2 & 2/3 & 1 \\ -3/2 & 1/3 & 1/2 \\ 1 & -1/3 & 0 \end{bmatrix}$$

$$\begin{bmatrix}
0 & 0 & 1 & 1 & 1 & 0 & 0 \\
0 & 0 & 1 & 1 & 1 & 0 & 0 \\
0 & 0 & 1/2 & 0 & 0 & 0 & 0 \\
0 & 0 & 1/2 & 0 & 0 & 0 & 0 \\
0 & 0 & 1/2 & 0 & 0 & 0 & 0 \\
0 & 0 & 1/2 & 0 & 0 & 0 & 0 \\
0 & 0 & 1/2 & 0 & 0 & 0 & 0 \\
0 & 0 & 0 & 0 & 0 & 0 & 0 \\
0 & 0 & 0 & 0 & 0 & 0 & 0 \\
0 & 0 & 0 & 0 & 0 & 0 & 0 \\
0 & 0 & 0 & 0 & 0 & 0 & 0 \\
0 & 0 & 0 & 0 & 0 & 0 & 0 \\
0 & 0 & 0 & 0 & 0 & 0 & 0 \\
0 & 0 & 0 & 0 & 0 & 0 & 0 \\
0 & 0 & 0 & 0 & 0 & 0 & 0 \\
0 & 0 & 0 & 0 & 0 & 0 & 0 \\
0 & 0 & 0 & 0 & 0 & 0 & 0 \\
0 & 0 & 0 & 0 & 0 & 0 & 0 \\
0 & 0 & 0 & 0 & 0 & 0 & 0 \\
0 & 0 & 0 & 0 & 0 & 0 & 0 \\
0 & 0 & 0 & 0 & 0 & 0 & 0 \\
0 & 0 & 0 & 0 & 0 & 0 & 0 \\
0 & 0 & 0 & 0 & 0 & 0 & 0 \\
0 & 0 & 0 & 0 & 0 & 0 & 0 \\
0 & 0 & 0 & 0 & 0 & 0 & 0 \\
0 & 0 & 0 & 0 & 0 & 0 & 0 \\
0 & 0 & 0 & 0 & 0 & 0 & 0 \\
0 & 0 & 0 & 0 & 0 & 0 & 0 \\
0 & 0 & 0 & 0 & 0 & 0 & 0 \\
0 & 0 & 0 & 0 & 0 & 0 & 0 \\
0 & 0 & 0 & 0 & 0 & 0 & 0 \\
0 & 0 & 0 & 0 & 0 & 0 & 0 \\
0 & 0 & 0 & 0 & 0 & 0 & 0 \\
0 & 0 & 0 & 0 & 0 & 0 & 0 \\
0 & 0 & 0 & 0 & 0 & 0 & 0 \\
0 & 0 & 0 & 0 & 0 & 0 & 0 \\
0 & 0 & 0 & 0 & 0 & 0 & 0 \\
0 & 0 & 0 & 0 & 0 & 0 & 0 \\
0 & 0 & 0 & 0 & 0 & 0 & 0 \\
0 & 0 & 0 & 0 & 0 & 0 & 0 \\
0 & 0 & 0 & 0 & 0 & 0 & 0 \\
0 & 0 & 0 & 0 & 0 & 0 & 0 \\
0 & 0 & 0 & 0 & 0 & 0 & 0 \\
0 & 0 & 0 & 0 & 0 & 0 & 0 \\
0 & 0 & 0 & 0 & 0 & 0 & 0 \\
0 & 0 & 0 & 0 & 0 & 0 & 0 \\
0 & 0 & 0 & 0 & 0 & 0 & 0 \\
0 & 0 & 0 & 0 & 0 & 0 & 0 \\
0 & 0 & 0 & 0 & 0 & 0 & 0 \\
0 & 0 & 0 & 0 & 0 & 0 & 0 \\
0 & 0 & 0 & 0 & 0 & 0 & 0 \\
0 & 0 & 0 & 0 & 0 & 0 & 0 \\
0 & 0 & 0 & 0 & 0 & 0 & 0 \\
0 & 0 & 0 & 0 & 0 & 0 & 0 \\
0 & 0 & 0 & 0 & 0 & 0 & 0 \\
0 & 0 & 0 & 0 & 0 & 0 & 0 \\
0 & 0 & 0 & 0 & 0 & 0 & 0 \\
0 & 0 & 0 & 0 & 0 & 0 & 0 \\
0 & 0 & 0 & 0 & 0 & 0 & 0 \\
0 & 0 & 0 & 0 & 0 & 0 & 0 \\
0 & 0 & 0 & 0 & 0 & 0 & 0 \\
0 & 0 & 0 & 0 & 0 & 0 & 0 \\
0 & 0 & 0 & 0 & 0 & 0 & 0 \\
0 & 0 & 0 & 0 & 0 & 0 & 0 \\
0 & 0 & 0 & 0 & 0 & 0 & 0 \\
0 & 0 & 0 & 0 & 0 & 0 & 0 \\
0 & 0 & 0 & 0 & 0 & 0 & 0 \\
0 & 0 & 0 & 0 & 0 & 0 & 0 \\
0 & 0 & 0 & 0 & 0 & 0 & 0 \\
0 & 0 & 0 & 0 & 0 & 0 & 0 \\
0 & 0 & 0 & 0 & 0 & 0 \\
0 & 0 & 0 & 0 & 0 & 0 \\
0 & 0 & 0 & 0 & 0 & 0 \\$$

Zhuang Tianning

TE6

2 c) Let 
$$(A) = det(E_1) det(E_2) det(E_3) det(E_4) det(E_6) det($$

$$det(A) = 1 \begin{vmatrix} 2 & 3 & 1 \\ 2 & 0 & 0 \\ 2 & 5 & 7 \end{vmatrix} - 2 \begin{vmatrix} 2 & 3 & 1 \\ 0 & 0 & 0 \\ 1 & 5 & 7 \end{vmatrix} + (-1) \begin{vmatrix} 2 & 2 & 1 \\ 0 & 2 & 0 \\ 12 & 7 \end{vmatrix}$$

$$- 0 \begin{vmatrix} 2 & 2 & 3 \\ 0 & 2 & 0 \\ 12 & 5 \end{vmatrix}$$

$$= 1\left(-2\left|\frac{31}{57}\right| + 0\left|\frac{21}{27}\right| - 0\left|\frac{23}{25}\right|\right) \xrightarrow{\text{expansion flowy}}$$

$$-2\left(-0\left|\frac{31}{57}\right| + 0\left|\frac{21}{17}\right| - 0\left|\frac{23}{15}\right|\right) /$$

$$-1\left(-0\left|\frac{21}{27}\right| + 2\left|\frac{21}{17}\right| - 0\left|\frac{22}{12}\right|\right) - 0$$

$$= 1 \left( -2 \times 16 \right) -2 \left( 0 \right) -1 \left( 2 \times 13 \right)$$

$$= -58$$

3b) As the determinant of A is nonzero, A is invertible and the rank of the matrix = 4. Hence, all the allumns are integer dent and the 4 integer dent allumns fill IR4 space, thus the linear system Ax = b has a unique solution for every b

As the reduced row echelon from of A is the identity matrix I4

[A | b ] -> [I | A-1b] so Ax=b is construct for all value of a.

3c) i) 
$$\det(\frac{1}{2}A^{7}) = \frac{1}{2}^{4} \det(A^{7})$$
  
=  $\frac{1}{2}^{4} \det(A)$   
=  $-\frac{58}{16} = -\frac{29}{8}$ 

ii) det 
$$(AB^{-1})$$
 = det  $(A)$  det  $(B^{-1})$   
= det  $(A)$  det  $(B^{-1})$   
=  $-58 \times \frac{1}{3}$   
=  $-\frac{58}{3}$ 

iii) det 
$$((313)^{-1}) = (3^4 \times det(13))^{-1}$$
  
=  $3^{-5}$   
=  $\frac{1}{243}$