

Math 115A Homework 4

Jiaping Zeng

5/30/2020

2. Let V be a finite dimensional inner product space over $\mathbb{F} = \mathbb{R}$ or \mathbb{C} .

(a) Fix $y \in V$ and suppose $\langle x, y \rangle = 0$ for all $x \in V$. Show that $y = 0$.

Answer: By contradiction. Suppose y is nonzero and choose $x = y \neq 0$, then $\langle x, y \rangle = 0 \implies \langle x, x \rangle = 0$ for $x \neq 0$. This contradicts axiom (iv) of inner product space, therefore the initial assumption is false and $y = 0$.

(b) Let $T : V \rightarrow V$ be a linear map such that $\langle T(x), T(y) \rangle = \langle x, y \rangle$ for all pairs $x, y \in V$ (we call such a map a *metric* map). Prove that T is an isomorphism.

Answer: Let $\dim V = n$ and $\beta = \{u_1, \dots, u_n\}$ orthogonal basis of V . Take two arbitrary vectors of β , u_x and u_y , we have $\langle u_x, u_y \rangle = \langle T(u_x), T(u_y) \rangle$ by definition of T . Since β is an orthonormal basis, $\langle u_x, u_y \rangle = 0$ and therefore $\langle T(u_x), T(u_y) \rangle = 0$, implying that $T(u_x)$ and $T(u_y)$ are orthogonal. Using this process on u_1, \dots, u_n yields the orthogonal set $\gamma = \{T(u_1), \dots, T(u_n)\}$. Since γ is automatically linearly independent and has dimension n , $T(\beta) = \gamma$ spans V , i.e. $\text{rank}(T) = n = \dim(V)$. Then by rank-nullity theorem T is an isomorphism.

(c) Find all metric maps $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ that have $\det T = 1$.

Answer: Let $x = \begin{pmatrix} x_1 \\ x_2 \end{pmatrix}$ and $y = \begin{pmatrix} y_1 \\ y_2 \end{pmatrix}$ be generic nonzero vectors of \mathbb{R}^2 . In addition, let

$[T] = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$. Then, $\det T = 1$ implies that $ad - bc = 1$. Using the definition of metric map, we also have $\langle T(x), T(y) \rangle = \langle x, y \rangle$. We can substitute T , x and y as follows:

$$\begin{aligned} \left\langle \begin{pmatrix} a & b \\ c & d \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix}, \begin{pmatrix} a & b \\ c & d \end{pmatrix} \begin{pmatrix} y_1 \\ y_2 \end{pmatrix} \right\rangle &= \left\langle \begin{pmatrix} x_1 \\ x_2 \end{pmatrix}, \begin{pmatrix} y_1 \\ y_2 \end{pmatrix} \right\rangle \\ \implies \left\langle \begin{pmatrix} ax_1 + bx_2 \\ cx_1 + dx_2 \end{pmatrix}, \begin{pmatrix} ay_1 + by_2 \\ cy_1 + dy_2 \end{pmatrix} \right\rangle &= x_1y_1 + x_2y_2 \\ \implies (ax_1 + bx_2)(ay_1 + by_2) + (cx_1 + dx_2)(cy_1 + dy_2) &= x_1y_1 + x_2y_2 \\ \implies (a^2 + c^2)x_1y_1 + (b^2 + d^2)x_2y_2 + (ab + cd)(x_1y_2 + x_2y_1) &= x_1y_1 + x_2y_2 \end{aligned}$$

By matching coefficients, we have the following system of equations:

$$a^2 + c^2 = 1, b^2 + d^2 = 1, ab + cd = 0, ad - bc = 1$$

which has solution

$$a = d, b = \pm\sqrt{1-a^2}, c = \mp\sqrt{1-a^2}$$

Then, the set of all $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ that satisfies the form $[T] = \begin{pmatrix} a & \pm\sqrt{1-a^2} \\ \mp\sqrt{1-a^2} & a \end{pmatrix}$ contains all metric maps with $\det T = 1$.

4. Let V be an inner product space and let $r : V \rightarrow V^*$ be the map $r(x) = \varphi_x := \langle x, - \rangle$. In class we showed that if V is finite dimensional then r is an isomorphism.

- (a) Assume that V is finite dimensional. Prove that r is injective.

Answer:

- (b) Let $V = \mathbb{R}[x]$ and let $W = \{(a_0, a_1, \dots) \mid a_i \in \mathbb{R}\}$ be the vector space of all infinite sequences. Show that the map $f : V^* \rightarrow W$ given by $f(\varphi) = (\varphi(x^n))_{n \geq 0}$.

Answer:

- (c) Use this to demonstrate that r is not necessarily surjective, i.e. find an element $\varphi \in V^*$ such that $\varphi \neq r(p)$ for any $p \in \mathbb{R}[x]$.

Answer:

5. Let V be a finite dimensional inner product space. For any $T : V \rightarrow V$ define $\check{T} : V^* \rightarrow V^*$ by $\check{T}(\phi) = \phi \circ T$. Furthermore for any $X : V^* \rightarrow V^*$ define $X^\perp : V \rightarrow V$ by $X^\perp = r^{-1} \circ X \circ r$. Prove that $T^* = \check{T}^\perp$.

Answer: