

RtEstim: Effective reproduction number estimation with trend filtering

Jiaping Liu^{a,1}, Zhenglun Cai^b, Paul Gustafson^a, and Daniel J. McDonald^a

^aDepartment of Statistics, The University of British Columbia

^bCentre for Health Evaluation and Outcome Sciences, The University of British Columbia

November 21, 2023

Abstract

To understand the transmissibility and spread of infectious diseases, epidemiologists turn to estimates of the effective reproduction number. While many approaches exist, their utility may be limited by the challenges of surveillance data collection. Arbitrary model assumptions that are unverifiable with data alone and sophisticated, though computationally inefficient, frameworks that are critical limitations for many existing approaches. We propose a discrete spline-based approach **RtEstim** that solves a convex optimization problem — Poisson trend filtering — using the proximal Newton method. It produces a locally adaptive estimator for effective reproduction number estimation with heterogeneous smoothness. **RtEstim** remains accurate even under some process misspecifications and is computationally efficient, even for large-scale data. The implementation is easily accessible in a lightweight R package [rtestim](#).

Keywords: tuning parameter selection | heterogeneous smoothness

Author summary

Effective reproduction number estimation faces many challenges, for example, data is usually collected from multiple sources and there are reporting delays, that cause many problems, such as missing values, underreporting and lack of standardization. Such poor conditions of data hinder the accurate estimation of effective reproduction number. Our motivation is to develop a model that returns accurate estimation, which is robust to violation of model misspecification, and a tool that is straightforward to

use and computationally efficient, even for large-scale data. Considering the temporal evolution of effective reproduction number for multiple degrees, we propose a convex optimization model with an ℓ_1 trend filtering penalty for univariate data. We solve our model using the proximal Newton method, which converges rapidly and is numerically stable. Our software, R package `RtEstim`, can return the output in seconds for univariate incidence sequences with length 300 using candidate sets of length fifty and ten-fold cross validation.

Contents

1	Introduction	3
2	Methods	6
2.1	Renewal model for incidence data	6
2.2	Poisson trend filtering estimator	8
2.3	Solving over a sequence of tuning parameters	10
2.4	Choosing a final λ	10
2.5	Approximate confidence bands	11
2.6	Bayesian perspective	11
3	Results	12
3.1	Simulation settings	12
3.2	Simulation results	14
3.3	Real-data results: Covid-19 incident cases in British Columbia	17
3.4	Real-data results: Pandemic influenza in Baltimore, Maryland, 1918	19
4	Discussion	19
	References	21

¹To whom correspondence should be addressed. E-mail: jiaping.liu@stat.ubc.ca

1 Introduction

The effective reproduction number (also called the instantaneous reproduction number) is a key quantity for understanding infectious disease dynamics including the potential size of a pandemic, the required stringency of prevention measures, and the efficacy of other controls. The effective reproduction number is defined to be the average number of secondary infections caused by a new primary infection that occurs at a specific time. Tracking the time series of this quantity is therefore useful for understanding whether or not future infections are likely to increase or decrease from the current state. Let $\mathcal{R}(t)$ denote the effective reproduction number at time t . Practically, as long as $\mathcal{R}(t) < 1$, infections will decline gradually, eventually resulting in a disease-free equilibrium, whereas when $\mathcal{R}(t) > 1$, infections will continue to increase, resulting in endemic equilibrium. While $\mathcal{R}(t)$ is fundamentally a continuous time quantity, it can be related to data only at discrete points in time $t = 1, \dots, n$. This sequence of effective reproduction numbers over time is not observable, but, nonetheless, is easily interpretable and retrospectively describes the course of an epidemic. Therefore, a number of procedures exist to estimate \mathcal{R}_t from different types of observed incidence data such as cases, deaths, or hospitalizations, while relying on various domain-specific assumptions. Importantly, accurate estimation of effective reproduction numbers relies heavily on the quality of the available data, and, due to the limitations of data collection, such as underreporting and lack of standardization, estimation methodologies rely on various assumptions to compensate. Because model assumptions may not be easily verifiable from data alone, it is also critical for any estimation procedure to be robust to model misspecification.

Many existing approaches for effective reproduction number estimation are Bayesian: they estimate the posterior distribution of \mathcal{R}_t conditional on the observations. One of the first such approaches is the software **EpiEstim** (Cori et al., 2020), described by Cori et al. (2013). This method is prospective, in that it uses only observations available up to time t in order to estimate \mathcal{R}_t for each $i = 1, \dots, t$. An advantage of **EpiEstim** is its straightforward statistical model: new incidence data follows the Poisson distribution conditional on past incidence combined with the conjugate gamma prior distribution for \mathcal{R}_t with fixed hyperparameters. Additionally, the serial interval distribution is fixed and known. For this reason, **EpiEstim** requires little domain expertise for use, and it is computationally fast. Thompson et al. (2019) modified this method to distinguish imported cases from local transmission and simultaneously estimate the serial interval distribution. Nash et al. (2023) further extended **EpiEstim** by using “reconstructed” daily incidence data to handle irregularly spaced observations. Abbott et al. (2020) proposed a Bayesian latent variable framework, **EpiNow2** (Abbott et al., 2023), which leverages incident cases, deaths or other available streams simultaneously along with allowing

additional delay distributions (incubation period and onset to reporting delays) in modelling. [Lison et al. \(2023\)](#) proposed an extension that handles missing data by imputation followed by a truncation adjustment. These modifications are intended to increase accuracy at the most recent (but most uncertain) timepoints, to aid policymakers. [Parag \(2021\)](#) also proposed a Bayesian approach, **EpiFilter** based on the (discretized) Kalman filter and smoother. **EpiFilter** also estimates the posterior of \mathcal{R}_t given a Gamma prior and Poisson distributed incident cases. Compared to **EpiEstim**, however, **EpiFilter** estimates \mathcal{R}_t retrospectively using all available incidence data both before and after time t , with the goal of being more robust in low-incidence cases. [Gressani et al. \(2022b\)](#) proposed a Bayesian P-splines approach, **EpiLPS**, that assumes negative Binomial distributed incidence. [Trevisin et al. \(2023\)](#) also proposed a Bayesian model estimated with particle filtering to incorporate spatial structures. Bayesian approaches estimate the posterior distribution of the effective reproduction numbers and possess the advantage that credible intervals may be easily computed. A limitation of many Bayesian approaches, however, is that they usually require more intensive computational routines, especially when observed data sequences are long or hierarchical structures are complex. Below, we compare our method to two of the more computationally efficient Bayesian models, **EpiEstim** and **EpiLPS**.

There are also frequentist approaches for \mathcal{R}_t estimation. [Abry et al. \(2020\)](#) proposed regularizing the smoothness of \mathcal{R}_t through penalized regression with second-order temporal regularization, additional spatial penalties, and with Poisson loss. [Pascal et al. \(2022\)](#) extended this procedure by introducing another penalty on outliers. [Pircalabelu \(2023\)](#) proposed a spline-based model relying on the assumption of exponential-family distributed incidence. [Ho et al. \(2023\)](#) estimates \mathcal{R}_t while monitoring the time-varying level of overdispersion. There are other spline-based approaches such as [Azmon et al. \(2014\)](#); [Gressani et al. \(2022a\)](#), autoregressive models with random effects ([Jin et al., 2023](#)) that are robust to low incidence cases, and generalized autoregressive moving average (GARMA) models ([Hettinger et al., 2023](#)) that are robust to measurement errors in incidence data.

We propose a retrospective effective reproduction number estimator called **RtEstim** that requires only incidence data. Our model makes the conditional Poisson assumption, similar to much of the prior work described above, but is empirically more robust to misspecification. This estimator is defined by a convex optimization problem with Poisson loss and ℓ_1 penalty on the temporal evolution of $\log(\mathcal{R}_t)$ to impose smoothness over time. As a result, **RtEstim** generates discrete splines, and the estimated curves (in log-space) appear to be piecewise polynomials of an order selected by the user. Importantly, The estimates are locally adaptive, meaning that different time ranges may possess heterogeneous smoothness. Because we penalize the logarithm of \mathcal{R}_t , we naturally accommodate the positivity requirement, can handle large or




Figure 1: A demonstration of effective reproduction number estimation by `RtEstim` and the corresponding fitted incidence cases for the Covid-19 epidemic in British Columbia, Canada during a period from March 28, 2020 to June 28, 2023. The blue curve in the top panel is the estimated piecewise quadratic \mathcal{R}_t and the gray ribbon is the corresponding 95% confidence band. The black curve in the bottom panel is the observed Covid-19 daily confirmed cases, and the orange curve is the fitted incidence corresponding to the estimated \mathcal{R}_t .

small incidence measurements, and are automatically (reasonably) robust to outliers without additional constraints. A small illustration using three years of Covid-19 case data in British Columbia is shown in Figure 1.

While our approach is straightforward and requires little domain knowledge for implementation, we also implement a number of refinements. We use a proximal Newton method to solve the convex optimization problem along with warm starts to produce estimates efficiently, typically in a matter of seconds, even for long sequences of data. In a number of simulated experiments, we show empirically that our approach is more accurate than existing methods at estimating the true effective reproduction numbers.

The manuscript proceeds as follows. We first introduce the methodology of `RtEstim` including the usage of renewal equation and the development of Poisson trend filtering estimator. We explain how this method could be interpreted from the Bayesian perspective,

connecting it to previous work in this context. We provide illustrative experiments comparing our estimator to `EpiEstim` and `EpiLPS`. We then apply our `RtEstim` on the Covid-19 pandemic incidence in British Columbia and the 1918 influenza pandemic in the United States. Finally, we conclude with a discussion of the advantages and limitations of our approach and describe practical considerations for effective reproduction number estimation.

2 Methods

2.1 Renewal model for incidence data

The effective reproduction number $\mathcal{R}(t)$ is defined to be the expected number of secondary infections at time t produced by a primary infection sometime in the past. To make this precise, denote the number of new infections at time t as $y(t)$. Then the total primary infectiousness can be written as $\eta(t) := \int_0^\infty p(i)y(t-i)\mathrm{d}i$, where $p(i)$ is the probability that a new secondary infection is the result primary infection which occurred i time units in the past. The reproduction number is then given as the value that equates

$$\mathbb{E}[y(t) \mid y(j), j < t] = \mathcal{R}(t)\eta(t) = \mathcal{R}(t) \int_0^\infty p(i)y(t-i)\mathrm{d}i, \quad (1)$$

otherwise known as the renewal equation. The period between primary and secondary infections is exactly the generation time of the disease, but given real data, observed at discrete times (say, daily) this delay distribution must be discretized into contiguous time intervals, say, $(0, 1], (1, 2], \dots$, which results in the sequence $\{p_i\}_1^\infty$ corresponding to observations y_t and resulting in the discretized version of Equation (1),

$$\mathbb{E}[y_t \mid y_1, \dots, y_{t-1}] = \mathcal{R}_t\eta_t = \mathcal{R}_t \sum_{i=0}^\infty p_i y_{t-i}. \quad (2)$$

Many approaches to estimating \mathcal{R}_t rely on Equation (2) as motivation for their procedures, among them, `EpiEstim` (Cori et al., 2013) and `EpiFilter` (Parag, 2021).

In most cases, it is safe to assume that infectiousness disappears beyond τ timepoints ($p_i = 0$ for $i > \tau$) so that the truncated integral of the generation interval distribution $\int_0^\tau p_i \mathrm{d}i = 1$. Generation time, however, is usually unobservable and tricky to estimate, so common practice is to approximate it by the serial interval: the period between the symptom onsets of primary and secondary infections. If the infectiousness profile after symptom onset is independent of the incubation period (the period from the time of infection to the time of symptom onset), then this approximation is justifiable: the serial interval distribution and

the generation interval distribution share the same mean. However, other properties may not be similarly shared, and, in general, the generation interval distribution is a convolution of the serial interval distribution with the distribution of the difference between independent draws from the delay distribution from infection to symptom onset. See, for example, [Gostic et al. \(2020\)](#) for a fuller discussion of the dangers of this approximation. Nonetheless, treating these as interchangeable is common ([Cori et al., 2013](#)) and beyond the scope of this work. Additionally, we assume that the generation interval (and, therefore, the serial interval), is constant over time t . That is, the probability $p(i)$ depends only on the gap between primary and secondary infections and not on the time t when the secondary infection occurs. For our methods, we will assume that the serial interval can be accurately estimated from auxiliary data (say by contact tracing, or previous epidemics) and we will take it as fixed, as is common in existing studies, e.g., [Abry et al. \(2020\)](#); [Cori et al. \(2013\)](#); [Pascal et al. \(2022\)](#).

The renewal equation in Equation (2) relates observable data streams (incident cases) occurring at different time points to the reproduction number given the serial interval. The fact that it depends only on the observed incidence counts makes it reasonable to estimate \mathcal{R}_t . However, this relationship obscures some difficulties in data collection. Diagnostic testing targets symptomatic individuals, omitting asymptomatic primary infections which can lead to future secondary infections. Testing practices, availability, and uptake can vary across space and time ([Hitchings et al., 2021](#); [Pitzer et al., 2021](#)). Finally, incident cases as reported to public health are subject to delays due to laboratory confirmation, test turnaround times, and eventual submission to public health ([Pellis et al., 2021](#)). For these reasons, reported cases are lagging indicators of the course of the pandemic. Furthermore, they do not represent the actual number of new infections that occur on a given day, as indicated by exposure to the pathogen. The assumptions described above (constant serial interval distribution, homogenous mixing, similar susceptibility and social behaviours, etc.) are therefore consequential. That said, Equation (2) also provides some comfort about deviations from these assumptions. If y_t is scaled by a constant (in time) describing the reporting ratio, then it will cancel from both sides. Similar arguments mean that even if such a scaling varies in time, as long as it varies slowly relative to the set of p_i that are larger than 0, Equation (2) will be a reasonably accurate approximation, so that \mathcal{R}_t can still be estimated well from reported incidence data. Finally, even a sudden change, say from c_1 for $i = 1, \dots, t_1$ to c_2 for $i > t_1$ would only result in large errors for t in the neighbourhood of t_1 (where the size of this neighbourhood is again determined by the effective support of $\{p_i\}$). This robustness to certain types of data reporting issues partially justifies using Equation (2) to calculate \mathcal{R}_t .

2.2 Poisson trend filtering estimator

We use the daily confirmed incident cases y_t on day t to estimate the observed infectious cases under the model that y_t given previous incident cases y_{t-1}, \dots, y_1 and a constant serial interval distribution follows a Poisson distribution with mean Λ_t . That is,

$$y_t \mid y_1, \dots, y_{t-1} \sim \text{Poisson}(\Lambda_t), \text{ where } \Lambda_t = \mathcal{R}_t \sum_{i=0}^{t-1} p_i y_{t-i} = \mathcal{R}_t \eta_t.$$

Given a history of n confirmed incidence counts $\mathbf{y} = (y_1, \dots, y_n)^\top$, our goal is to estimate \mathcal{R}_t . A natural approach is to maximize the likelihood, producing the MLE:

$$\begin{aligned} \hat{\mathcal{R}} &= \operatorname{argmax}_{\mathcal{R} \in \mathbb{R}_+^n} \mathbb{P}(\mathcal{R} \mid \mathbf{y}, \mathbf{p}) = \operatorname{argmax}_{\mathcal{R} \in \mathbb{R}_+^n} \prod_{t=1, \dots, n} \frac{e^{-\mathcal{R}_t \eta_t} (\mathcal{R}_t \eta_t)^{y_t}}{y_t!} \\ &= \operatorname{argmin}_{\mathcal{R} \in \mathbb{R}_+^n} \frac{1}{n} \sum_{t=1}^n \mathcal{R}_t \eta_t - y_t \log(\mathcal{R}_t \eta_t). \end{aligned} \tag{3}$$

This optimization problem, however, is easily seen to yield a one-to-one correspondence between the confirmed cases and the effective reproduction, i.e., $\hat{\mathcal{R}}_t = y_t / \eta_t$, so that the estimated sequence $\hat{\mathcal{R}}$ will have no significant graphical smoothness.

The MLE is an unbiased estimator of the true parameter \mathcal{R}_t , but unfortunately has high variance: changes in y_t result in proportional changes in $\hat{\mathcal{R}}_t$. To avoid this behaviour, and to match the intuition that $\mathcal{R}_t \approx \mathcal{R}_{t-1}$, we advocate enforcing smoothness of the effective reproduction numbers. This requirement will decrease the variance, and hopefully lead to more accurate estimation of \mathcal{R} , as long as the smoothness assumption is reasonable. Smoothness assumptions are common (see e.g., [Parag \(2021\)](#) or [Gostic et al. \(2020\)](#)), but the *type* of smoothness assumed is critical. [Cori et al. \(2020\)](#) imposes smoothness indirectly by estimating \mathcal{R}_t with moving windows of of past observations. The Kalman filter procedure of [Parag \(2021\)](#) would enforce in ℓ_2 -smoothness ($\int_0^n (\hat{\mathcal{R}}''(t))^2 dt < C$ for some C), although the algorithm results in $\hat{\mathcal{R}}$ taking values over a discrete grid. [Pascal et al. \(2022\)](#) produces piecewise-linear $\hat{\mathcal{R}}_t$, which turns out to closely related to a special case of our methodology. Smoother estimated curves will provide high-level information about the entire epidemic, obscuring small local changes in $\mathcal{R}(t)$, but may also remove the ability to detect large sudden changes, such as those resulting from lockdowns or other major containment policies.

We choose to implement smoothness by assuming that $\log(\mathcal{R}(t))$ is a piecewise polynomial of arbitrary degree. We specifically consider discrete splines with various degrees of continuity. For example, 0th-degree discrete splines are piecewise constant, the 1st-degree curves are piecewise linear, and 2nd-degree curves are piecewise quadratic. For $k \geq 1$, k^{th} -degree discrete

splines are continuous and have continuous discrete differences up to degree $k - 1$ at the knots. To achieve such smoothness, we regularize the size of changes between adjacent effective reproduction numbers. Because $\mathcal{R}_t > 0$, we explicitly penalize the divided differences (discrete derivatives) of neighbouring values of $\log(\mathcal{R}_t)$. To achieve this, we penalize the ℓ_1 norm of the divided differences, which introduces sparsity into the curvature, so that the estimates have heterogeneous smoothness in different periods of time. It is a more realistic setting compared to homogeneous smoothness created by the squared ℓ_2 norm. Using different orders of divided differences result in estimated effective reproduction numbers with different smoothness assumptions.

To enforce smoothness of $\hat{\mathcal{R}}_t$, we add a trend filtering penalty to Equation (4) (Kim et al., 2009; Sadhanala et al., 2022; Tibshirani, 2014, 2022). Let $\theta := \log(\mathcal{R}) \in \mathbb{R}^n$, so that $\Lambda_t = \eta_t \exp(\theta_t)$, and $\log(\eta_t \mathcal{R}_t) = \log(\eta_t) + \theta_t$. For evenly spaced incident case data, we write our estimator as the solution to the optimization problem

$$\hat{\mathcal{R}} = \exp(\hat{\theta}) \quad \text{where} \quad \hat{\theta} = \underset{\theta \in \mathbb{R}^n}{\operatorname{argmin}} \eta^\top \exp(\theta) - \mathbf{y}^\top \theta + \lambda \|D^{(k+1)}\theta\|_1, \quad (4)$$

where $\exp(\cdot)$ applies elementwise. Here, $D^{(k+1)} \in \mathbb{Z}^{(n-k-1) \times n}$ is the $(k+1)^{\text{th}}$ order divided difference matrix for any $k \in \{0, \dots, n-1\}$. $D^{(k+1)}$ is defined recursively as $D^{(k+1)} = D^{(1)}D^{(k)}$, where $D^{(1)} \in \{-1, 0, 1\}^{(n-k-1) \times (n-k)}$ is a sparse matrix with diagonal band:

$$D^{(1)} = \begin{pmatrix} -1 & 1 & & & \\ & -1 & 1 & & \\ & & \ddots & \ddots & \\ & & & -1 & 1 \end{pmatrix}.$$

The tuning parameter λ balances data fidelity with desired smoothness. When $\lambda = 0$, the problem in Equation (4) reduces to the MLE in Equation (3). Larger tuning parameters privilege the regularization term and yield smoother estimates. Finally, there exists λ_{\max} such that any $\lambda \geq \lambda_{\max}$ will result in $D^{(k+1)}\hat{\theta} = 0$ and $\hat{\theta}$ will be the Kullback-Leibler projection of \mathbf{y} onto the null space of $D^{(k+1)}$ (see Section 2.3).

For unevenly-spaced incidence data, the spacing between neighboring parameters varies with the time between observations, and thus, the divided differences must be adjusted by the times that the observations occur. Given observation times $\mathbf{x} = (x_1, \dots, x_n)^\top$, for $k \geq 1$, define a k^{th} -order diagonal matrix

$$X^{(k)} = \operatorname{diag} \left(\frac{k}{x_{k+1} - x_1}, \frac{k}{x_{k+2} - x_2}, \dots, \frac{k}{x_n - x_{n-k}} \right).$$

Letting $D^{(\mathbf{x},1)} := D^{(1)}$, then for $k \geq 1$, the $(k+1)^{\text{th}}$ -order divided difference matrix for unevenly spaced data can be created recursively by $D^{(\mathbf{x},k+1)} := D^{(1)}X^{(k)}D^{(\mathbf{x},k)}$. No adjustment is required for $k = 0$.

Due to the penalty structure, this estimator is locally adaptive, meaning that it can potentially capture local changes such as the initiation of control measures. [Abry et al. \(2020\)](#); [Pascal et al. \(2022\)](#) considered only the 2nd-order divided difference of \mathcal{R}_t rather than its logarithm. In comparison to their work, our estimator (1) allows for arbitrary degrees of temporal smoothness and (2) avoids the potential numerical issues of penalizing/estimating positive real values. Furthermore, as we will describe below, our procedure is computationally efficient for estimation over an entire sequence of penalty strengths λ and provides methods for choosing how smooth the final estimate should be.

2.3 Solving over a sequence of tuning parameters

We can solve the Poisson trend filtering estimator over an arbitrary sequence of λ that produces different levels of smoothness in the estimated curves. We consider a candidate set $\boldsymbol{\lambda} = \{\lambda_m\}_{m=1}^M$, which contains a strictly decreasing λ sequence.

Let $D := D^{(k+1)}$ for simplicity in the remainder of this section. As $\lambda \rightarrow \infty$, the penalty term $\lambda \|D\theta\|_1$ dominates the Poisson objective, so that minimizing the objective is asymptotically equivalent to minimizing the penalty term, which results in $\|D\theta\|_1 = 0$. In this case, the divided differences of θ with order $k+1$ is always 0, and thus, θ must lie in the null space of D , that is, $\theta \in \mathcal{N}(D)$. The same happens for any λ beyond this threshold, so define λ_{\max} to be the smallest λ that produces $\theta \in \mathcal{N}(D)$. It turns out that this value can be written explicitly as $\lambda_{\max} = \|(D^\dagger)^\top (\eta - y)\|_\infty$, where D^\dagger is the (left) generalized inverse of D satisfying $D^\dagger D = I$. Therefore, we use $\lambda_1 = \lambda_{\max}$ and then choose the minimum λ_M to be $r\lambda_{\max}$ for some $r \in (0, 1)$ (typically $r = 10^{-5}$). Given any $M \geq 3$, we generate a sequence of λ values to be equally spaced on the log-scale between λ_1 and λ_M .

2.4 Choosing a final λ

We estimate model accuracy over the candidate set through K -fold cross validation (CV) to choose the best tuning parameter. Specifically, we divide \mathbf{y} (except the first and last observations) roughly evenly and randomly into K folds, estimate \mathcal{R}_t for all $\boldsymbol{\lambda}$ leaving one fold out, and then predict the held-out observations. Model accuracy can be measured by multiple metrics such as mean squared error $\text{MSE}(\hat{y}, y) = n^{-1} \|\hat{y} - y\|_2^2$ or mean absolute error $\text{MAE}(\hat{y}, y) = n^{-1} \|\hat{y} - y\|_1$, but we prefer to use the deviance, to mimic the likelihood in [Equation 3](#): $D(y, \hat{y}) = \frac{1}{n} \sum_{i=1}^n 2(y_i \log(y_i) - y_i \log(\hat{y}_i) - y_i + \hat{y}_i)$, where we define $0 \log(0) =$

0.

To estimate the sequence efficiently, the model is fitted sequentially by visiting each component of $\boldsymbol{\lambda}$ in order. The estimates produced with a larger λ are used as the initial values (warm starts) for the next smaller λ . By solving through the entire sequence of tuning parameters, we have a better chance to achieve a better trade-off between bias and variance, and accordingly, improved accuracy relative to procedures examining one fixed value of λ . Note that for any K and any M , we will end up estimating the model $(k+1)M$ times rather than once.

2.5 Approximate confidence bands

We also provide empirical confidence bands of the estimators with approximate coverage guarantees. Consider the related estimator $\tilde{\mathcal{R}}_t$ defined as

$$\tilde{\mathcal{R}} = \exp(\tilde{\theta}) \quad \text{where} \quad \tilde{\theta} = \underset{\theta \in \mathbb{R}^n}{\operatorname{argmin}} \eta^\top \exp(\theta) - \mathbf{y}^\top \theta + \lambda \|D\theta\|_2^2.$$

Let $\tilde{\mathbf{y}} = \eta \circ \mathcal{R}$, and then it can be shown (for example, Theorem 2 in [Vaiteer et al., 2017](#)) that an estimator for $\operatorname{Var}(\tilde{\mathbf{y}})$ is given by $(\operatorname{diag}(\tilde{\mathbf{y}}^{-2}) + \lambda D^\top D)^\dagger$. Finally, an application of the delta method shows that $\operatorname{Var}(\tilde{\mathbf{y}}_t)/\eta_t^2$ is an estimator for $\operatorname{Var}(\tilde{\mathcal{R}}_t)$ for each $t = 1, \dots, n$. We therefore use $(\operatorname{diag}(\tilde{\mathbf{y}}^{-2}) + \lambda D^\top D)_t^\dagger/\eta_t^2$ as an estimator for $\operatorname{Var}(\hat{\mathcal{R}}_t)$. An approximate $(1-\alpha)\%$ confidence interval then can be written as $\hat{\mathcal{R}} \pm T(\alpha/2, n - \text{df})s$, where s_t is the square-root of $\operatorname{Var}(\hat{\mathcal{R}}_t)$ for each $t = 1, \dots, n$ and df is the number of changepoints in $\tilde{\theta}$ plus $k+1$.

2.6 Bayesian perspective

Unlike many other methods for \mathcal{R}_t estimation, our approach is frequentist rather than Bayesian. Nonetheless, it has a corresponding Bayesian interpretation: as a state-space model with Poisson observational noise, autoregressive transition equation of degree $k \geq 0$, e.g., $\theta_{t+1} = 2\theta_t - \theta_{t-1} + \varepsilon_{t+1}$ for $k = 1$, and Laplace transition noise $\varepsilon_{t+1} \sim \text{Laplace}(0, 1/\lambda)$. Compared to **EpiFilter** ([Parag, 2021](#)), another retrospective study of \mathcal{R}_t , we share same observational assumptions, but our approach has a different transition noise. **EpiFilter** estimates the posterior distribution of \mathcal{R}_t , and thus it can provide credible interval estimates as well. Our approach produces the maximum *a posteriori* estimate via an efficient convex optimization, omitting the need for MCMC sampling. But the associated confidence bands are created differently.

3 Results

Implementation of the our approach is provided in the R package `rtestim`. All experiments are run in R with version 4.3.1 on a MacBook with an Apple M1 Pro chip and 32GB RAM running under macOS Sonoma 14.0. The R packages used for simulation and real-data application are versions `EpiEstim` 2.2-4, `EpiLPS` 1.2.0, and `rtestim` 0.0.4.

3.1 Simulation settings

We simulate four scenarios of the time-varying effective reproduction numbers, intended to mimic different epidemics. The first two scenarios are simple cases that are rapidly controlled by intervention, where the graphical curves consist of one discontinuity and two segments. Scenario 1 has constant \mathcal{R}_t before and after an intervention, while Scenario 2 grows exponentially, then decays. The other two scenarios are more complicated, where more waves in the epidemics are involved. Scenario 3 has four linear segments with three discontinuities, which reflect the effect of an intervention, resurgence to rapid transmission, and finally suppression of pandemic. Scenario 4 involves sinusoidal waves throughout the epidemic. The first three scenarios and the last scenario are motivated by [Parag \(2021\)](#) and [Gressani et al. \(2022b\)](#) respectively. We name the four scenarios as (1) 2-segment constant, (2) 2-segment exponential curve, (3) 4-segment linear line, and (4) periodic curve respectively.

In all cases, the times of observation are regular, and consider epidemics of length $n = 300$. Specifically, in Scenario 1, $\mathcal{R}_t = 2, 0.8$ before and after $t = 70$. In Scenario 2, \mathcal{R}_t increases and decreases exponentially with rates 0.015, 0.005 pre and post $t = 50$. In Scenario 3, \mathcal{R}_t reduces from 2.5 to 2 linearly between $t \in [1, 60]$, falls to 0.8 at $t = 61$ and goes linearly down to 0.6 until $t = 110$, resurges to 1.7 at $t = 111$ and grows linearly back to 2 until $t = 150$, and then drops to 0.9 at $t = 151$ and descends to 0.5 until the end. In Scenario 4, \mathcal{R}_t is realization of the continuous, periodic curve generated by the function

$$\mathcal{R}(t) = 0.2 \left((\sin(\pi t/12) + 1) + (2 \sin(\pi t/6) + 2) + (3 \sin(\pi t/1.2) + 3) \right),$$

evaluated at equally spaced points $t \in [0, 10]$. These settings are illustrated in the left column of [Figure 2](#).

We assume that the serial interval follows Gamma distribution with fixed shapes and scales $(3, 3)$, $(2.5, 2.5)$, $(3.5, 3.5)$ and $(3.5, 3.5)$ for Scenarios 1–4 respectively. We consider all epidemics starting from $y_1 = 2$ cases and generating until timepoints $t = 300$. We compute the expected incidence Λ_t using the renewal equation, and generate the incidence samples from the Poisson distribution $y_t \sim \text{Pois}(\Lambda_t)$. To verify the performance of our model under the violation

of this distributional assumption of incidence, we also generate incidence samples using the negative Binomial distribution with dispersion size 5, i.e., $y_t \sim \text{NB}(\Lambda_t, \text{size} = 5)$. We generate 50 random samples for each setting of experiments, resulting in 400 total experiments. An example of each effective reproduction number scenario with a single corresponding Poisson and negative Binomial sample incidence sequences is displayed in [Figure 2](#).




Figure 2: The effective reproduction numbers (left column) and corresponding sample incident cases drawn from a Poisson (middle column) or negative Binomial (right column) distribution. The rows correspond to the four \mathcal{R}_t settings.

We compare `RtEstim` to `EpiEstim` and `EpiLPS`. Unfortunately, `EpiFilter` frequently fails to converge due to the large case counts in many simulations. `EpiEstim` estimates the posterior distribution of effective reproduction numbers given a Gamma prior and Poisson distributed incidence. It estimates the reproduction number over a sliding window, assuming the reproduction number is constant during the specific time window. A longer sliding window averages out more fluctuations, leading to smoother estimates, whereas, a shorter sliding window is more responsive to sudden spikes or declines. We tried the default, a weekly sliding window, as well as a monthly window. However, since neither considerably outperform the other across all scenarios, we defer the monthly case to the Appendix. `EpiLPS` is another Bayesian approach that estimates P-splines coupled with Laplace approximations of the conditional posterior based on the negative Binomial likelihood. For `RtEstim` on the four scenarios respectively, we estimate (1) piecewise constant $k = 0$, (2) piecewise linear & cubic $k = 1, 3$, (3) piecewise linear $k = 1$ and (4) piecewise cubic polynomials $k = 3$. In each case, we examine a grid of 50 λ s, selecting the best using 10-fold cross validation. For all models and problems, we use the generation interval distribution used to create the data.

To measure estimation accuracy, we compare $\hat{\mathcal{R}}$ to \mathcal{R} using the Kullback-Leibler (KL) divergence: a standard metric that measures the distance between two probability distributions. Since \mathcal{R}_t can be regarded as the expectation of Poisson distribution, we use the KL divergence for Poisson distributions (averaged across all coordinates) to measure the accuracy of the \mathcal{R}_t estimates

$$D_{KL}(\mathcal{R} \parallel \hat{\mathcal{R}}) = \sum_{t=1}^n w_t \left(\mathcal{R}_t \log \left(\frac{\mathcal{R}_t}{\hat{\mathcal{R}}_t} \right) + \hat{\mathcal{R}}_t - \mathcal{R}_t \right),$$

where $\mathcal{R} = \{\mathcal{R}_t\}_{t=1}^n$ and $w_t = \eta_t / \sum_t \eta_t$ is the rescaled total infectiousness. To fairly compare across methods, we drop the estimates during the first week because estimates from `EpiEstim` do not begin until $t = 8$ (using a weekly window). KL divergence is more appropriate for measuring accuracy because it connects directly to the Poisson likelihood used to generate the data, whereas standard measures like the mean-squared error correspond to Gaussian data. This has the effect of increasing the relative cost of mistakes when Λ_t is small. Other details of the experimental settings are deferred to the Appendix.

3.2 Simulation results

`RtEstim` overall outperforms `EpiEstim` and `EpiLPS` in the experimental study. [Figure 3](#) visualizes the KL divergence across the three models. Under both Poisson and negative Binomial distributions, `RtEstim` is easily the most accurate across Scenarios 1 and 3: the median of KL divergence is much lower and the boxes frequently fail to overlap indicating a better performance than the other two methods across all 50 simulations. The advantage is

less for the negative Binomial case compared to the Poisson case, but still obvious. **RtEstim** and **EpiLPS** have similar performance in Scenarios 2 and 4. For the Poisson case, **RtEstim** and **EpiLPS** both have very small KL scores, which are very close to 0. In Scenario 4, **RtEstim** is slightly better for Poisson and **EpiLPS** is better for negative Binomial, but the boxes largely overlap to each other. **EpiLPS** has a slightly lower median and a smaller IQR in Scenario 2 for the negative Binomial case. The two degrees of **RtEstim** used for Scenario 2 perform very close to each other for both distributions, which implies a low risk of model misspecification. We will examine a single realization of each experiment to investigate these global conclusions in more detail.




Figure 3: Boxplot of KL divergence between the estimated $\hat{\mathcal{R}}_t$ and the true \mathcal{R}_t across 50 random samples for each approach given Poisson incidence (*in top panels*) and negative Binomial incidence (*in bottom panels*) respectively. Outliers are excluded. Full visualization is deferred to the Appendix. A few **RtEstim** models for Scenario 2 run expired, since they reach the maximum iteration number, and thus, are excluded here.

Figure 4 shows 1 realization for the estimated reproduction numbers under the Poisson generative model for all 4 scenarios. Compared to **EpiEstim** and **EpiLPS**, which have rather severe difficulties at the beginning of the time series, **RtEstim** estimates are more accurate




Figure 4: Example of effective reproduction number estimation for Poisson incidence.

— they nearly overlap with the true values — without suffering from the edge problem. Scenario 2 is a difficult problem for all methods: the sharp increase at the end of the stage of exponential growth is difficult to capture. Although the truth is piecewise exponential are likely best represented with a piecewise cubic curve, the actual curvature is so gentle that linear estimation ($k = 1$) appears potentially reasonable. We, therefore, fit piecewise linear and cubic $\hat{\mathcal{R}}_t$ curves using `RtEstim` for Scenario 2 to evaluate model misspecification. However, both cases have difficulty recovering the acute rise in the growth phase. An explanation of such failure is that the model imposes discontinuity at the changepoint, which hinders the estimates from fitting the two discontinuous phases. Scenario 1 is the simplest case with only one knot and two constant segments. Besides the edge problem, `EpiEstim` and `EpiLPS` produce “smooth” estimated curves that are continuous at the changepoint, which results in large mistakes in that neighbourhood. Since the piecewise constant `RtEstim` estimator does not force any smoothness in \mathcal{R}_t , it easily captures the sharp decrease change.

To investigate the performance under the violation of the Poisson distributional assumption (of both `RtEstim` and `EpiEstim`), we also examine estimation accuracy with negative Binomial data. Figure 5 displays a realization, analogous to the previous case, for all methods and scenarios. `RtEstim` has more difficulty relative to the Poisson incidence setting, especially




Figure 5: Example of effective reproduction number estimation for negative Binomial incidence.

at the beginning of the outbreak. This is most pronounced in Scenario 4, where **RtEstim** is overly smooth, except in the last wave. In Scenario 2, **RtEstim** successfully captures the changepoint, but suffers from the same problem as in the Poisson setting. In Scenario 3, the piecewise linear **RtEstim** recovers the curvature of \mathcal{R}_t well, but is less accurate than in the Poisson incidence cases.

Finally, it is important to provide a brief comparison of the running times of three models across the 8 experimental settings. We find that almost all models across all experiments complete within 10 seconds. **RtEstim** generally takes the longest, likely due to a relatively large candidate set — 50 values of λ and 10 folds of cross validation — while other models run only a single time for a fixed setting of hyperparameters per experiment. Additional results on timing comparisons are deferred to the Appendix.

3.3 Real-data results: Covid-19 incident cases in British Columbia

We implement **RtEstim** on Covid-19 incident confirmed cases in British Columbia (B.C.) as reported on May 18, 2023 (visualized in Figure 6) by the B.C. Centre for Disease Control. The reported incident cases provide a snapshot of how testing recommendations and practices




Figure 6: Covid-19 daily confirmed incident cases between March 1st, 2020 and April 15th, 2023 in British Columbia, Canada.

have changed over the three years. We use the gamma distribution with shape 2.5 and scale 2.5 to approximate the serial interval function, which is empirically reasonable, since they are close to the parameters in a recent study (Lehtinen et al., 2021), which summarizes estimated parameters of the serial interval distributions of SARS-CoV-2 by different approaches.

Considering the first, second, and third polynomial degrees, the estimated effective reproduction numbers of Covid-19 in British Columbia (illustrated in Figure 7) are always less than 3 except at the very early stage, which means that one distinct infected individuals on average infects less than three other individuals in the population. Examining three different settings for k , the temporal evolution of $\hat{\mathcal{R}}$ (across all regularization levels λ) are similar near the highest peak around the end of 2021 before dropping shortly thereafter. Throughout the estimated curves, the peaks and troughs of the reproduction numbers precede the growth and decay cycles of confirmed cases, as expected. We also visualize 95% confidence bands for the point estimates at the smallest one of the “optimal” tuning parameters with lowest CV (i.e., KL divergence) scores in Figure 7.

The estimated reproduction numbers are relatively unstable before April 1st, 2022. The highest peak coincides with the emergence and global spread of the Omicron variant. The estimated reproduction numbers fall below 1 during two time periods — roughly from April 1st, 2021 to July 1st, 2021 and from January 1st, 2022 to April 1st, 2022. The first trough

coincides with the introduction of Covid-19 vaccines in British Columbia. The second trough, shortly after the greatest peak may be due to variety of factors resulting in the depletion of the susceptible population such as increased self-isolation in response to the peak and media and immunity incurred via recent infection. Since around April 1st, 2022, estimated reproduction numbers have remained relatively stable (fluctuating around 1) corresponding to low reported cases (though reporting behaviours have also changed significantly since the Omicron wave).

3.4 Real-data results: Pandemic influenza in Baltimore, Maryland, 1918

We also apply `RtEstim` to daily reported influenza cases in Baltimore, Maryland occurring during the world-wide pandemic of 1918 from September to November. The dataset (shown in [Figure 8](#)) is included in the `EpiEstim` R package. The 1918 influenza outbreak, caused by the H1N1 influenza A virus, was an unprecedentedly deadly, infecting almost one-third of the population across the world ([Taubenberger and Morens, 2006](#)). In [Figure 9](#), the CV-tuned piecewise cubic estimates best capture the growth at the beginning of the pandemic. It suggests that the pandemic grew rapidly over the first 30 days before declining below 1 after 50 days. However, it also suggests an increase toward the end of the period. With the currently available dataset, it is hard to tell if there is a next wave or a steady decline ahead. The CV-tuned piecewise constant and linear estimates in [Figure 9](#) both suggest a steady decline. It is supported by a couple of arguments. The influenza incidence reduce to 0 in the end. The \mathcal{R} estimates of `EpiEstim` with 1-week, 2-week and 4-week sliding windows in [Cori et al. \(2013\)](#) are all lower than the threshold 1 in their right tails.

4 Discussion

The `RtEstim` methodology provides a locally adaptive estimator using Poisson trend filtering on univariate data. It captures the heterogeneous smoothness of effective reproduction numbers given observed incidence data rather than resulting in global smoothness. This is a nonparametric regression model which can be written as a convex optimization (minimization) problem. Minimizing the distance (averaged KL divergence per coordinate) between the estimators and (functions of) observations guarantees data fidelity while the divided differences between pairs of neighbouring parameters imposes smoothness. The ℓ_1 -regularization results in sparsity of the divided differences, which leads to heterogeneous smoothness within certain periods of time.

The property of local adaptivity (heterogenous smoothness) is useful to automatically distinguish, for example, seasonal outbreaks from outbreaks driven by other factors (behavioural changes, foreign introduction, etc.). Given a well-chosen polynomial degree, for example, the growth rates can be quickly detected, alerting public health to implement policy changes. The effective reproduction numbers can be estimated retrospectively to examine the efficacy of such policies, whether they result in \mathcal{R}_t falling below 1 or the speed of their effects. The smoothness of \mathcal{R}_t curves (including the polynomial degrees and tuning parameters) should be chosen based on the purpose of the study in practice, e.g., epidemic forecasting may require a more wiggly curve that contains more fluctuation information, while retrospective studies that solely target on understanding of the pandemic may prefer a smoother curve with less important information smoothed out.

Our method **RtEstim** provides a natural way to deal with missing data, for example, on weekends and holidays or due to changes in reporting frequency. While solving the convex optimization problem, the edge lengths of the line graphs are adjusted, correctly penalizing the distance between irregularly spaced data. Computing the total primary infectiousness is also easily generalized to irregular reporting by modifying the discretization of the serial interval distribution. Additionally, because the ℓ_1 penalty introduces sparsity (operating like a median rather than a mean), this procedure is relatively insensitive to outliers compared to ℓ_2 regularization.

There are a number of limitations that may influence the quality of \mathcal{R}_t estimation. While our model is generic for incidence data, rather than tailored to any specific disease, it does assume that the generation interval is short relative to the period of data collection. More specialized methodologies would be required for diseases with long incubation periods such as HIV or Hepatitis. Our approach, does not explicitly model imported cases, nor distinguish between subpopulations that may have different mixing behaviour. While the Poisson distribution is common, it does not handle overdispersion (observation variance larger than the mean). The negative binomial distribution is a good alternative, but more difficult to estimate in this context. As described in the introduction section, justifying the expression for \mathcal{R} assumes that a relatively constant proportion of true infections is reported. However, if this proportion varies with time (say, due to change in surveillance practices and testing recommendations), the estimates may be biased over this window. A good example is that in early January 2021, during the height of the Omicron wave, British Columbia moved from testing all symptomatic individuals to testing only those in at-risk groups. The result was a sudden change that would render \mathcal{R}_t estimates on either side of this time point incommensurable.

As currently implemented, **RtEstim** uses a fixed serial interval throughout the period

of study, but as factors such as population immunity vary, the serial interval may vary as well (Nash et al., 2023). Another issue relates to the equating serial and generation intervals (also mentioned above). The serial interval distribution is generally wider than that of the generation interval, because the serial interval involves the convolution of two distributions, and is unlikely to actually follow a named distribution like Gamma, though it may be reasonably well approximated by one. Our implementation allows for an arbitrary distribution to be used, but requires the user to specify the discretization explicitly, requiring more nuanced knowledge than is typically available. Pushing this analysis further, to accommodate other types of incidence data (hospitalizations or deaths), a modified generation interval distribution would be necessary, and further assumptions would be required as well. Or else, one would first need to deconvolve deaths to infection onset before using our software.

Besides all the advantages and limitations of the methodology discussed above, our `RtEstim` estimator can be implemented easily through a lightweight R package `rtestim` and computed efficiently through the numerical algorithm, proximal Newton method developed in C++, especially for large-scale data. With accessible incidence cases, prespecified serial interval distribution, and chosen polynomial degrees, `RtEstim` is able to produce accurate estimation of effective reproduction numbers and provide tuning parameter selection through a path algorithm.

References

- Abbott, S., Funk, S., Hickson, J., Badr, H. S., Monticone, P., Ellis, P., Azam, J., Munday, J., Allen, J., Johnson, A., Pearson, C. A. B., actions user, Chapman, L., DeWitt, M., Bosse, N. and Meakin, S. (2023) epiforecasts/EpiNow2: 1.4.0 release. <https://doi.org/10.5281/zenodo.8380568>.
- Abbott, S., Hellewell, J., Thompson, R. N., Sherratt, K., Gibbs, H. P., Bosse, N. I., Munday, J. D., Meakin, S., Doughty, E. L., Chun, J. Y. et al. (2020) Estimating the time-varying reproduction number of SARS-CoV-2 using national and subnational case counts. *Wellcome Open Research*, **5**, 112. <https://doi.org/10.12688/wellcomeopenres.16006.2>.
- Abry, P., Pustelnik, N., Roux, S., Jensen, P., Flandrin, P., Gribonval, R., Lucas, C.-G., Guichard, É., Borgnat, P. and Garnier, N. (2020) Spatial and temporal regularization to estimate COVID-19 reproduction number $R(t)$: Promoting piecewise smoothness via convex optimization. *PLoS ONE*, **15**, e0237901. <https://doi.org/10.1371/journal.pone.0237901>.

- Azmon, A., Faes, C. and Hens, N. (2014) On the estimation of the reproduction number based on misreported epidemic data. *Statistics in Medicine*, **33**, 1176–1192. <https://doi.org/10.1002/sim.6015>.
- Cori, A., Cauchemez, S., Ferguson, N. M., Fraser, C., Dahlgvist, E., Demarsh, P. A., Jombart, T., Kamvar, Z. N., Lessler, J., Li, S. et al. (2020) EpiEstim: estimate time varying reproduction numbers from epidemic curves. Version 2.2-4. <https://cran.r-project.org/package=EpiEstim>.
- Cori, A., Ferguson, N. M., Fraser, C. and Cauchemez, S. (2013) A new framework and software to estimate time-varying reproduction numbers during epidemics. *American Journal of Epidemiology*, **178**, 1505–1512. <https://doi.org/10.1093/aje/kwt133>.
- Gostic, K. M., McGough, L., Baskerville, E. B., Abbott, S., Joshi, K., Tedijanto, C., Kahn, R., Niehus, R., Hay, J. A., De Salazar, P. M. et al. (2020) Practical considerations for measuring the effective reproductive number, R_t . *PLoS Computational Biology*, **16**, e1008409. <https://doi.org/10.1371/journal.pcbi.1008409>.
- Gressani, O., Faes, C. and Hens, N. (2022a) An approximate Bayesian approach for estimation of the instantaneous reproduction number under misreported epidemic data. *Biometrical Journal*, **65**, 2200024. <https://doi.org/10.1002/bimj.202200024>.
- Gressani, O., Wallinga, J., Althaus, C. L., Hens, N. and Faes, C. (2022b) Epilps: A fast and flexible Bayesian tool for estimation of the time-varying reproduction number. *PLoS Computational Biology*, **18**, e1010618. <https://doi.org/10.1371/journal.pcbi.1010618>.
- Hettinger, G., Rubin, D. and Huang, J. (2023) Estimating the instantaneous reproduction number with imperfect data: a method to account for case-reporting variation and serial interval uncertainty. *arXiv preprint arXiv:2302.12078*. <https://doi.org/10.48550/arXiv.2302.12078>.
- Hitchings, M. D., Dean, N. E., García-Carreras, B., Hladish, T. J., Huang, A. T., Yang, B. and Cummings, D. A. (2021) The usefulness of the test-positive proportion of severe acute respiratory syndrome coronavirus 2 as a surveillance tool. *American Journal of Epidemiology*, **190**, 1396–1405. <https://doi.org/10.1093/aje/kwab023>.
- Ho, F., Parag, K. V., Adam, D. C., Lau, E. H., Cowling, B. J. and Tsang, T. K. (2023) Accounting for the potential of overdispersion in estimation of the time-varying reproduction number. *Epidemiology*, **34**, 201–205. <https://doi.org/10.1097/ede.0000000000001563>.

- Jin, S., Dickens, B. L., Lim, J. T. and Cook, A. R. (2023) EpiMix: A novel method to estimate effective reproduction number. *Infectious Disease Modelling*, **8**, 704–716. <https://doi.org/10.1016/j.idm.2023.06.002>.
- Kim, S.-J., Koh, K., Boyd, S. and Gorinevsky, D. (2009) ℓ_1 trend filtering. *SIAM Review*, **51**, 339–360. <https://doi.org/10.1137/070690274>.
- Lehtinen, S., Ashcroft, P. and Bonhoeffer, S. (2021) On the relationship between serial interval, infectiousness profile and generation time. *Journal of the Royal Society Interface*, **18**, 20200756. <https://doi.org/10.1098/rsif.2020.0756>.
- Lison, A., Abbott, S., Huisman, J. and Stadler, T. (2023) Generative Bayesian modeling to nowcast the effective reproduction number from line list data with missing symptom onset dates. *arXiv preprint arXiv:2308.13262*. <https://doi.org/10.48550/arXiv.2308.13262>.
- Nash, R. K., Bhatt, S., Cori, A. and Nouvellet, P. (2023) Estimating the epidemic reproduction number from temporally aggregated incidence data: A statistical modelling approach and software tool. *PLoS Computational Biology*, **19**, e1011439. <https://doi.org/10.1371/journal.pcbi.1011439>.
- Parag, K. V. (2021) Improved estimation of time-varying reproduction numbers at low case incidence and between epidemic waves. *PLoS Computational Biology*, **17**, e1009347. <https://doi.org/10.1371/journal.pcbi.1009347>.
- Pascal, B., Abry, P., Pustelnik, N., Roux, S., Gribonval, R. and Flandrin, P. (2022) Nonsmooth convex optimization to estimate the Covid-19 reproduction number space-time evolution with robustness against low quality data. *IEEE Transactions on Signal Processing*, **70**, 2859–2868. <https://doi.org/10.1109/TSP.2022.3180926>.
- Pellis, L., Scarabel, F., Stage, H. B., Overton, C. E., Chappell, L. H., Fearon, E., Bennett, E., Lythgoe, K. A., House, T. A., Hall, I. et al. (2021) Challenges in control of COVID-19: short doubling time and long delay to effect of interventions. *Philosophical Transactions of the Royal Society B*, **376**, 20200264. <https://doi.org/10.1098/rstb.2020.0264>.
- Pircalabelu, E. (2023) A spline-based time-varying reproduction number for modelling epidemiological outbreaks. *Journal of the Royal Statistical Society Series C: Applied Statistics*, **72**, 688–702. <https://doi.org/10.1093/jrssc/qlad027>.
- Pitzer, V. E., Chitwood, M., Havumaki, J., Menzies, N. A., Perniciaro, S., Warren, J. L., Weinberger, D. M. and Cohen, T. (2021) The impact of changes in diagnostic testing

- practices on estimates of COVID-19 transmission in the United States. *American Journal of Epidemiology*, **190**, 1908–1917. <https://doi.org/10.1093/aje/kwab089>.
- Sadhanala, V., Bassett, R., Sharpnack, J. and McDonald, D. J. (2022) Exponential family trend filtering on lattices. *arXiv preprint arXiv:2209.09175*. <https://doi.org/10.48550/arXiv.2209.09175>.
- Taubenberger, J. K. and Morens, D. M. (2006) 1918 Influenza: the mother of all pandemics. *Emerging Infectious Diseases*, **17**, 69–79. <https://doi.org/10.3201%2Fid1201.050979>.
- Thompson, R. N., Stockwin, J. E., van Gaalen, R. D., Polonsky, J. A., Kamvar, Z. N., Demarsh, P. A., Dahlqvist, E., Li, S., Miguel, E., Jombart, T. et al. (2019) Improved inference of time-varying reproduction numbers during infectious disease outbreaks. *Epidemics*, **29**, 100356. <https://doi.org/10.1016/j.epidem.2019.100356>.
- Tibshirani, R. J. (2014) Adaptive piecewise polynomial estimation via trend filtering. *The Annals of Statistics*, **42**, 285–323. <https://doi.org/10.1214/13-AOS1189>.
- (2022) Divided differences, falling factorials, and discrete splines: Another look at trend filtering and related problems. *Foundations and Trends® in Machine Learning*, **15**, 694–846. <http://dx.doi.org/10.1561/22000000099>.
- Trevisin, C., Bertuzzo, E., Pasetto, D., Mari, L., Miccoli, S., Casagrandi, R., Gatto, M. and Rinaldo, A. (2023) Spatially explicit effective reproduction numbers from incidence and mobility data. *Proceedings of the National Academy of Sciences*, **120**, e2219816120. <https://doi.org/10.1073/pnas.2219816120>.
- Vaiter, S., Deledalle, C., Fadili, J., Peyré, G. and Dossal, C. (2017) The degrees of freedom of partly smooth regularizers. *Annals of the Institute of Statistical Mathematics*, **69**, 791–832. <https://doi.org/10.1007/s10463-016-0563-z>.




Figure 7: Estimated effective reproduction numbers for Covid19 daily confirmed counts between March 1st, 2020 and April 15th, 2023 in British Columbia, Canada. The left panels demonstrate estimates corresponding to 50 tuning parameters. The right panels display the CV-tuned estimates with 95% confidence intervals. The top, medium and bottom panels illustrate the estimated reproduction numbers (\mathcal{R}_t) using the Poisson trend filtering (in Equation (4)) with degrees $k = 1, 2, 3$ respectively.




Figure 8: Daily influenza incidence counts in Baltimore, Maryland between September and November in 1918.




Figure 9: Estimated effective reproduction numbers for influenza in Baltimore, Maryland in 1918. The left panels show estimates for all 50 tuning parameters under consideration. The right column displays the CV-tuned estimates with 95% confidence bands. The rows (top to bottom) show estimated reproduction numbers (\mathcal{R}_t) using the Poisson trend filtering (in Equation (4)) with degrees $k = 1, 2, 3$ respectively.