


1. (a) For the input point $[2,1]$ and $y = -1$, with weights 0.15 applied. For identity: The first layer returns with results $\begin{bmatrix} 0.2778125 & 0.2778125 \\ 0.13890625 & 0.13890625 \end{bmatrix}$ and layer two returns with results $\begin{bmatrix} 0.47529864 & 0.47529864 \end{bmatrix}$. For tanh: first layer w1: $\begin{bmatrix} 0.27569254 & 0.27569254 \end{bmatrix}$, second layer $\begin{bmatrix} 0.13784627 & 0.13784627 \end{bmatrix}$, $E_{in}(w)$ obtained was 0.3172898231266975
- (b) After peturbing the weights, I achieved the following: identity w1: $\begin{bmatrix} 0.27796641 & 0.27796641 \\ 0.1389832 & 0.1389832 \end{bmatrix}$ w2: $\begin{bmatrix} 0.47564328 & 0.47564328 \end{bmatrix}$ and tanh: w1: $\begin{bmatrix} 0.2758401 & 0.2758401 \end{bmatrix}$ w2: $\begin{bmatrix} 0.13792005 & 0.13792005 \end{bmatrix}$, $E_{in}(w)$ was 0.31737905364130986

As stated by the hint, after peturbing the weights, the results were relatively similar, only different in the final few digits, therefore we can conclude that our algorithm is correct.

2. FINISH

3. (a) We're given two two-dimensional inputs, $x_1 = (1,0)$ and $x_2 = (-1,0)$. The optimal separating hyperplane is simply a line. For SVM, we want to find the separator that maximizes the margin. Both of these lines depict a horizontal line with slope 0, the perpendicular bisector is the negative reciprocal of the line, which is a vertical line. A vertical line in $x = 0$ is our solution.
- (b) (i) They are the same point $z_1 = (1,0)$ and $z_2 = (-1,0)$
(ii) The same, $z = 0$ and it's a vertical line.



(c)

- (d) For $z(x) = \begin{bmatrix} x_1^3 - x_2 \\ x_1 x_2 \end{bmatrix}$, and $z(y) = \begin{bmatrix} y_1^3 - y_2 \\ y_1 y_2 \end{bmatrix}$, we can derive the following:

$$\begin{aligned}
 K(x, y) &= z(x) \cdot z(y) \\
 &= (x_1^3 - x_2)(y_1^3 - y_2) + (x_1 x_2)(y_1 y_2) \\
 &= x_1^3 y_1^3 - x_1^3 y_2 - x_2 y_1^3 + x_2 y_2 + x_1 x_2 y_1 y_2
 \end{aligned}$$

(e) $h(x) = \text{sign}(x_1^3 - x_2 + x_1x_2)$

4. FINISH