

# Supplementary Document of "A Knee Point Driven Evolutionary Algorithm for Multi-Objective Bilevel Optimization"

Jiaxin Chen, Jinliang Ding, *Senior Member, IEEE*, Ke Li, *Senior Member, IEEE*, Kay Chen Tan, *Fellow, IEEE*, and Tianyou Chai, *Life Fellow, IEEE*

## I. APPENDIX A

In this section, we first give a detailed description of the test problems proposed by us. Then, we give the characteristics and formulas of the test problems used in the comparison with the classical methods. After that, a practical problem is presented, including the background, process flow, and mathematical formula. Thereafter, we delineate the experimental setup of the algorithms.

### A. Benchmark Test Problems

We propose a series of multi-objective bilevel optimization test problems based on the bottom-up [1] and multi-objective bilevel optimization construction principles [2]. The proposed test problems (dubbed MBOP12 to MBOP15) are built on single-level test problems DEB2DK, DO2DK from [3], and DEB2DK2, DEB2DK1 from [4]. MBOP12 to MBOP15 present various difficulties to the multi-objective bilevel optimization, such as scalable, and discontinuity. The mathematical formulas and corresponding parameters are given in Table I.

The test problem MBOP12 is designed based on MBOP1 and DEB2DK, which imposes a more complicated lower-level Pareto front (PF) shape and upper-level optimal solution solving procedure. MBOP12 is a continuity, convexity, and scalable multi-objective bilevel test problem. For different upper-level decision variables, the lower-level Pareto-optimal solutions are located in  $f_1^L(\mathbf{x}) = q(x_1^L) \sin(\frac{\pi x_1^L}{2})$ ,  $f_2^L = q(x_1^L) \cos(\frac{\pi x_1^L}{2}) - x_1^U$ . For every  $i = 2, \dots, K$ ,  $x_i^L = x_i^U$ . Parameter  $\beta$  in  $q(x_1^L)$  controls the number of knee points in the lower level and is set to 4. For the upper-level problem, since the dominance relationship, the optimal solution is located on the lower left quarter circle with the center of  $(1 + r, 1 + r)$  and the radius of 1.

MBOP13 is a discontinuity problem with a difficult constraint, designed based on single-level DEB2DK2. For the lower-level problem,  $x_j^L = 0$ , for  $j = K + 1, \dots, K + L$ , and the Pareto-optimal solution is located in  $f_1(x) = r(x_1^L) \sin^3(\frac{\pi x_1^L}{2})$ ,  $f_2(x) = r(x_1^L) \cos(\frac{\pi x_1^L}{2})$ . But not all the solutions in the lower-level reaction set are in the inducible region because of upper-level constraint  $G(\mathbf{x}^U, \mathbf{x}^L)$ . We set the parameter controlling the number of knee points  $\beta$  as 2, and the upper-level problem has a discontinuous front, that is, the part where the PF of DEB2DK2 meets the constraint  $G(\mathbf{x}^U, \mathbf{x}^L)$ . Besides,  $x_j^L = 0$  for  $j = 2, \dots, K + L$ .

MBOP14 is designed according to the shape of DO2DK. In this problem, we add some difficulty to the solving procedure of the upper-level problem by setting the function  $T(t_1(\mathbf{x}), t_2(\mathbf{x}))$ . Through filtering some lower-level solutions, only part of the solutions can be the candidate for the upper-level optimal solutions. Thereby, examining the performance of the algorithm. For the upper-level PF,  $x_j^U = 0$ , for  $j = 2, \dots, K$ , and satisfies the function  $f_1(\mathbf{x}) = r(x_1^L) [\sin(\frac{\pi x_1^L}{2s+1} + (1 + \frac{2s-1}{2s+2})\pi) + 1]$ ,  $f_2(\mathbf{x}) = r(x_1^L) [\cos(\frac{\pi x_1^L}{2} + \pi) + 1]$ , in the part of  $z(x_1, x_2)$  region.

MBOP15 is a deceptive problem designed based on DEB2DK1. In this problem, we set  $\tau_1$  and  $\tau_2$  to positive numbers, which indicates that the solutions not in the feasible region may obtain a smaller upper-level function value than the true optimal solution. However, these solutions are not feasible according to the definition of bilevel optimization. This structure makes this problem difficult to solve. To prevent the algorithm from jumping into the undesired areas at the beginning, we use a progressive value  $T(x_1, x_2)$  to make the algorithm try to find a feasible solution. The upper- and lower-level optimal solutions are the intersection of the function ( $f_1(\mathbf{x}) = r(x_1^L) \sin \frac{\pi x_1^L}{2}$ ,  $f_2(\mathbf{x}) = r(x_1^L) \cos \frac{\pi x_1^L}{2}$ ) and region  $z_1(x_1, x_2)$ .

### B. Test Problems Used in Comparison to Classical Methods

To compare our proposed algorithm TKDEA with classical methods, we compare it with some classical methods. We first compare TKDEA with the classical methods proposed in [5] and [6] on the test problem in [6]. For convenience, the test problem and the two algorithms are dubbed Classic1, CA1, and CA2, respectively. CA1 is a classical method that is used to address linear multi-objective bilevel optimization problem (MBLOP) issues. The main idea of CA1 is to use a Pareto-filter scheme to find an approximated discrete representation of the efficient set. CA2 could solve the linear multi-objective bilevel problem where the constraint regions of the upper level and lower level be non-empty and compact. It uses fuzzy set theory and fuzzy programming to convert the linear multi-objective bilevel programming problem to a linear bilevel programming problem, then extends the  $K$ -th-best method to solve the final linear bilevel programming problem.

Then we compare TKDEA with the classical method (dubbed CA3) in [7]. We test the two algorithms on the test problem used in [7] (dubbed Classic2), which is also known as

TP3 [2], [8]. CA3 is a classical method that assumes some restrictions as bicriteria problems on both levels and a one-dimensional upper-level variable. It works on nonlinear non-convex MBLOPs. The detailed test problem formulations are given in Table II.

### C. Petroleum Refining Problem

To validate the effectiveness of our proposed algorithm, we also investigate the performance of TKDEA in a real-world engineering problem.

Petroleum refining is essential for crude oil application, which involves many complex technologies, such as refinery scheduling [9], operational optimization [10], and fast evaluation of physical properties [11]. This energy-intensive process emits a large chunk of greenhouse gases (e.g., carbon dioxide ( $\text{CO}_2$ )) while splitting crude oil. These greenhouse gases have a significant impact on the global climate and the environment. There are usually two DMs in this practical problem, i.e., the regulating authority and the petroleum refining enterprise. The regulating authority acts as a leader and aims to earn revenues through taxes and regulate the environmental damages. On the other hand, the petroleum refining enterprise reacts rationally to the decisions of the leader to maximize its profit and minimize its burnt fuel. The limited non-renewable crude oil resources and rising prices of burnt fuel are forcing the enterprises to achieve their objectives via operational optimization.

Petroleum refining is a huge engineering process that consumes large quantities of energy with many heavy equipments. It is unrealistic to measure all units, and we only study the main unit of petroleum refining, i.e., the crude oil distillation unit (CDU) consisting of an atmospheric furnace (AF) and an atmospheric distillation column (ADC).

After the crude oil is partially vaporized, the oil is heated by the AF and enters the ADC. The ADC consists of a main column, a condenser, three side-draw product strippers, and three pumparounds. The flashed feed vapors move upwards from the flash zone in the ADC, and the liquid moves downwards from the top to the bottom of the column due to gravity. The high-temperature gas and the low-temperature liquid meet at the tray for partial condensation and partial vaporization. The oil is cut off at different boiling points by this physical method. The top distillate product is naphtha, and the side-draw products from the three steam strippers are kerosene, light diesel, and atmospheric gas oil (AGO).

In terms of environmental preservation, heating and steam furnaces are one of the main sources of  $\text{CO}_2$ . The mathematical formulas for calculating the amount of fuel burnt and  $\text{CO}_2$  emission are taken from [12]. In this petroleum refining MBLOP, there are two assumptions. One is that the heat duty required by the furnace is proportional to the yield of the crude oil and product flow rate. The second is that the steam heat duty is proportional to the steam flow rate. The  $\text{CO}_2$  emission and the tax revenue of the regulating authority form the upper-level constraints. The lower-level constraints mean the overall product flow rate cannot exceed the oil flow rate, and the petroleum refining enterprise must obtain a certain income. This NP-hard problem is difficult to solve in a straightforward

way. The formulation of the petroleum refining problem is given in Table III. Some notations are used in the practical petroleum refining MBLOP. The summary of notations and their values are given in Table IV.

### D. Parameter Setting

The parameter setting of the experiment is given in Table V.  $N^U$  and  $N^L$  are the sizes of upper- and lower-level populations, respectively.  $T^U$  and  $T^L$  are the generations of upper- and lower-level problems, respectively. Since the complexity of BLEMO is  $O(N^U(2T^U+1)(T^L+1))$ , and TKDEA algorithm complexity is  $O(N^U(T^U+1)(T^L+1))$ . For the sake of fairness, the overall number of function evaluations should be kept as consistent as possible. Thus, the generation of the upper-level problem of BLEMO is set to be  $T^U/2$ . To eliminate any side effects brought the non-Pareto-optimal solutions or deceptive solutions, the solutions dominate the PF at the upper-level are removed from the HV calculation. In particular, the ideal and nadir points of each benchmark test problem are given in the seventh and eighth column of Table V.

The dimensions of upper- and lower-level decision variables are listed in the sixth column of Table V in the form of  $(n_u, n_\ell)$ , where  $n_u$  and  $n_\ell$  represent the dimension of decision variables at the upper level and lower level. For MBOP1–MBOP3 and MBOP12,  $K$  is set to be 3. For MBOP4, MBOP5, MBOP6, MBOP7, MBOP13, and MBOP15, two parameters control the dimension of the variable, where  $K$  is set to be 3, and  $L$  is set to be 2. For MBOP14,  $K$  is set to be 5. For the scalable MBOP9 test problem, the values of  $K$  are set to 2, 6, and 14, dubbed MBOP9-K2, MBOP9-K6, and MBOP9-K14, respectively. The dimensions of the upper-level decision variables  $n_u$  of MBOP13 to MBOP15 are set to be 5, 4, 4.

The parameter settings of Classic1 and Classic2 could be found in the last line but two and the last line but one of Table V, and the other parameter settings of the classical methods are the same as those in the original paper. Moreover, the parameter setting of the practical problem is given in the last line of Table V.

## II. APPENDIX B

This section verifies the effectiveness of TKDEA. First, the effectiveness of the TKDEA is investigated on the MBOP. Next, to verify the validity of the lower-level algorithm, we use ablation study for comparison. Finally, sensitivity study is performed to analyze the effect of the parameter.

### A. Comparisons with the peer algorithms

The statistical comparison results of IGD, HV, and IGD+ values, based on the Wilcoxon signed-rank test, are given in Table VII. From these results, we can see that TKDEA obtains the better results. In the following paragraph, we analyze the performance instance by instance.

MBOP1 to MBOP7 are test problems characterized by multi-modalities and scalable. TKDEA has achieved the best results on almost all the test problems. As for MBOP8, BLEMO obtains better IGD and HV values while TKDEA obtains a

better IGD+ value. Indeed, only some critical solutions at the lower level will respond to the optimal solution of the upper level. TKDEA may not find these solutions, resulting in the performance of TKDEA is not promising. MBOP9 is a difficult test problem. If the algorithm fails to find the lower-level Pareto-optimal set, these infeasible solutions are likely to dominate the true Pareto-optimal solutions at the upper level. For the MBOP9 on different lower-level variable dimensions, TKDEA is very competitive among all the dimensions and have a significantly better performance than the others. In terms of MBOP10, our proposed algorithm obtains the second-best performance. MBOP11 is a practical problem that lacks real PF, and we can see that TKDEA is significantly better than the other two algorithms.

Next, we analyze the numerical solution of MBOP11. MBLOP is a strongly NP-hard problem, so it is difficult to find the optimal bilevel solution. Only one numerical solution of MBOP11 is given in the original paper [13], i.e.,  $(\mathbf{x}^U = (146.2955, 28.9394), \mathbf{x}^L = (0, 67.9318, 0))$ . This numerical solution provides a good guide for subsequent research. However, we notice that some nondominated solutions are not discovered in most existing papers. In this paper, we give four nondominated solutions. Their variables are as follows: A:  $(\mathbf{x}^U = (0.5600, 107.5164), \mathbf{x}^L = (0, 0, 23.2238))$ , B:  $(\mathbf{x}^U = (16.6666, 100), \mathbf{x}^L = (0, 0, 29.6665))$ , C:  $(\mathbf{x}^U = (92.9000, 64.4244), \mathbf{x}^L = (0, 0, 60.1598))$ , and D:  $(\mathbf{x}^U = (146.1023, 29.0719), \mathbf{x}^L = (0, 67.6585, 0.2501))$ .  $\mathbf{x}^L$  is the approximate optimal lower-level decision variable we could find in the case of a fixed upper-level decision variable. It's worth noting that these solutions may not be the true bilevel optimal solutions (due to the strongly NP-hard of MBLOP), but these solutions are not dominated by the solutions obtained in the existing paper. The corresponding upper- and lower-level objective vectors could be found in Table VI.

MBOP12 to MBOP15 are test problems that contain scalable, deceptive, continuous, and discontinuous properties. And we can clearly see that TKDEA has much better performance than the comparison algorithms in terms of IGD, HV, and IGD+, demonstrating its effectiveness in tackling these test problems.

Moreover, Table VIII shows the comparisons of the overall, upper-level, and lower-level function evaluations (dubbed FE, ULFE, LLFE) achieved by three multi-objective bilevel optimization algorithms. The distribution and variance of IGD, HV and, IGD+ are given in Fig. 1, Fig. 2 and Fig. 3, respectively. Note that the IGD+ values of MBOP10 and MBOP11 are not available due to the lack of their real PFs. It can be seen from the above results that TKDEA achieves better performance than the other two algorithms.

### B. Comparisons with classical methods

Due to Classic1 and Classic2 lacking real PFs, we only calculate the HV values. Table IX and Table X give the detailed HV values of the compared algorithms. As can be seen from these results, TKDEA has an obvious advantage over CA1 and CA2 on Classic1 and obtains a better HV value on Classic2 than CA3.

### C. Ablation Study

To validate the competitiveness of the proposed lower-level algorithm, we use ablation study to investigate the effect of the lower-level algorithm.

1) *Effectiveness of the lower-level algorithm*: We first use ablation study to compare the proposed lower-level algorithm with several single-level constrained multi-objective algorithms (i.e., TKDEA, C-TAEA, I-DBEA, and DMOEA- $\epsilon$ C). The codes for C-TAEA, I-DBEA, and DMOEA- $\epsilon$ C are available on the PlatEMO [14]. All the algorithms are combined with the multi-objective bilevel optimization solving procedure introduced in Section III-A. The corresponding multi-objective bilevel algorithms are dubbed as B- $\mathbf{x}$ , where  $\mathbf{x}$  is either C-TAEA, I-DBEA, or DMOEA- $\epsilon$ C. Under the same solving procedure and generation settings, the number of function evaluations of the rest algorithms is the same as TKDEA. The statistical comparison results of IGD, IGD+, and HV values, based on the Wilcoxon signed-rank test, are presented in Table XI.

From these results, we can see that TKDEA significantly outperforms the other algorithms on most of the test problems. Especially in MBOP9 of different lower-level decision variable dimensions, TKDEA shows a competitive performance compared with other algorithms. For MBOP2, B-DMOEA- $\epsilon$ C achieves the best performance, while TKDEA does not perform very well. The reason may be that the upper-level population on MBOP2 test problem is easier trapped in local optimum due to the multi-modality of upper-level optimization problem. Thus causing TKDEA to fall into local optimum. According to the Wilcoxon signed-rank test, TKDEA obtains the best results on 51 and similar results on 29 out of 105 test problems. Moreover, the  $A_{12}$  effect size of four algorithms is given in Fig. 4. According to the  $A_{12}$  effect size of four algorithms, and the large effect sizes are 37%, 49%, and 54%, respectively. Moreover, we apply the Scott-Knott test to sort the performance of four algorithms. The Scott-Knott result is presented in Fig. 5. And we can clearly see from the result that TKDEA obtains first rank. After that, to observe the results intuitively, the data distributions of the IGD, HV, and IGD+ are given as violin plots and box plots in Fig. 6, Fig. 7, and Fig. 8, respectively. From these results, we can clearly see that our proposed algorithm has more centralized data distributions and obtains better results. Thus, we can conclude that TKDEA is competitive in dealing with the MBLOP.

2) *Effectiveness of two stages*: The above empirical study has shown that the proposed lower-level algorithm has better performance than the selected constrained multi-objective algorithms. The lower-level algorithm of TKDEA is composed of two stages. We empirically investigate the performance of TKDEA w.r.t. the variant TKDEA-S1 only uses the first-stage search step, and the variant TKDEA-S2 only uses the second-stage search step.

From the statistical comparison results of IGD, HV, and IGD+ results,  $A_{12}$  effect size results are shown in Fig. 9. We have witnessed a performance degradation when ablating any of the stages. And the sum of the better  $A_{12}$  effect size is 63% for TKDEA-S1 and 74% for TKDEA-S2. In addition, to facilitate an overall comparison, we further apply




Fig. 1. Violin plots and box plots of IGD achieved by three multi-objective bilevel optimization algorithms (The part corresponding to the dashed line is the enlarged plot).




Fig. 2. Violin plots and box plots of HV achieved by three multi-objective bilevel optimization algorithms (The part corresponding to the dashed line is the enlarged plot).




Fig. 3. Violin plots and box plots of IGD+ achieved by three multi-objective bilevel optimization algorithms (The part corresponding to the dashed line is the enlarged plot).




Fig. 4. Percentage of  $A_{12}$  effect size of three single-level algorithms combined with the same upper-level solving procedure compared with TKDEA.




Fig. 5. Violin plots and box plots of Scott-Knott test ranks achieved by four algorithms (the smaller rank is, the better performance achieved).

the Scott-Knott test to sort the performance of the compared algorithms, which is given in Fig. 10. TKDEA has shown the best performance, TKDEA-S1, TKDEA-S2, OMOPSO-BL, and BLEMO obtain second, third, fourth, and fifth ranks. Moreover, the upper-level nondominated solutions of three algorithms on MBOP-K14 is given in Fig. 11. From the figure, we can see that TKDEA can obtain more uniform solutions than

TKDEA-S1 and TKDEA-S2. From these results, we further confirm that TKDEA is better than TKDEA-S1 and TKDEA-S2 in the corresponding comparison.

#### D. Sensitivity Study

In our proposed algorithm, there is a switching parameter  $\alpha$  in the lower-level algorithm. To study the impact of parameter, we choose TKDEA as the baseline and empirically investigate the performance under different  $\alpha = \{0.1, 0.3, 0.5, 0.7, 0.9\}$  settings. Fig. 12, Fig. 13, and Fig. 14 give the box plots of IGD, HV, and IGD+. Two rows represent MBOP1 to MBOP3, MBOP6, and MBOP7, MBOP8 to MBOP9 (with different lower-level decision variable dimensions,  $K = 2, 6, 14$ ). From these figures, we can see that different  $\alpha$  results in different performances. In this paper, we set the parameter  $\alpha$  to 0.3.

#### REFERENCES

- [1] K. Deb, L. Thiele, M. Laumanns, and E. Zitzler, "Scalable test problems for evolutionary multiobjective optimization," in *Evolutionary Multiobjective Optimization*, ser. Advanced Information and Knowledge Processing, A. Abraham, L. C. Jain, and R. R. Goldberg, Eds. Springer, 2005, pp. 105–145.
- [2] K. Deb and A. Sinha, "An efficient and accurate solution methodology for bilevel multi-objective programming problems using a hybrid evolutionary-local-search algorithm," *Evol. Comput.*, vol. 18, no. 3, pp. 403–449, 2010.
- [3] J. Branke, K. Deb, H. Dierolf, and M. Osswald, "Finding knees in multi-objective optimization," in *PPSN'04: Proc. of the 8th International Conference on Parallel Problem Solving from Nature*, ser. Lecture Notes in Computer Science, vol. 3242. Springer, 2004, pp. 722–731.
- [4] L. Rachmawati and D. Srinivasan, "Multiobjective evolutionary algorithm with controllable focus on the knees of the pareto front," *IEEE Trans. Evol. Comput.*, vol. 13, no. 4, pp. 810–824, 2009.
- [5] C. O. Pieume, P. Marcotte, L. P. Fotso, and P. Siarry, "Solving bilevel linear multiobjective programming problems," *Am. J. Oper. Res.*, vol. 1, no. 4, pp. 214–219, 2011.




Fig. 6. Violin plots and box plots of IGD achieved by four algorithms (The part corresponding to the dashed line is the enlarged plot).




Fig. 7. Violin plots and box plots of HV achieved by four algorithms (The part corresponding to the dashed line is the enlarged plot).

- [6] M. H. Farahi and E. Ansari, "A new approach to solve multi-objective linear bilevel programming problems," *J. Math. Comput. Sci.*, vol. 1, no. 4, pp. 313–320, 2010.
- [7] G. Eichfelder, "Multiobjective bilevel optimization," *Mathematical Programming*, vol. 123, no. 2, pp. 419–449, 2010.
- [8] —, *Solving nonlinear multiobjective bilevel optimization problems with coupled upper level constraints*, 2007.
- [9] Y. Chen, J. Ding, and T. Chai, "A knowledge transfer based scheduling algorithm for large-scale refinery production," *IEEE Trans. Ind. Informatics*, vol. 18, no. 2, pp. 869–879, 2022.
- [10] Q. Chen, J. Ding, T. Chai, and Q. Pan, "Evolutionary optimization under uncertainty: The strategies to handle varied constraints for fluid catalytic cracking operation," *IEEE Trans. Cybern.*, vol. 52, no. 4, pp. 2249–2262, 2022.
- [11] J. Lu, J. Ding, C. Liu, and T. Chai, "Hierarchical-bayesian-based sparse stochastic configuration networks for construction of prediction intervals," *IEEE Trans. Neural Networks Learn. Syst.*, pp. 1–12, 2021.
- [12] M. Gadalla, Ž. Olujić, M. Jobson, and R. Smith, "Estimation and reduction of co2 emissions from crude oil distillation units," *Energy*, vol. 31, no. 13, pp. 2398–2408, 2006.
- [13] G. Zhang, J. Lu, and T. S. Dillon, "Decentralized multi-objective bilevel decision making with fuzzy demands," *Knowl. Based Syst.*, vol. 20, no. 5, pp. 495–507, 2007.
- [14] Y. Tian, R. Cheng, X. Zhang, and Y. Jin, "Platemo: A MATLAB platform for evolutionary multi-objective optimization," *IEEE Comput. Intell. Mag.*, vol. 12, no. 4, pp. 73–87, 2017.




Fig. 8. Violin plots and box plots of IGD+ achieved by four algorithms (The part corresponding to the dashed line is the enlarged plot).




Fig. 9. Percentage of  $A_{12}$  effect size of two algorithms.




Fig. 10. Violin plots and box plots of Scott-Knott test ranks achieved by five algorithms (the smaller rank is, the better performance achieved).




Fig. 11. Two stages of effectiveness on MBOP11-K14




Fig. 12. Collected comparisons of IGD when using different  $\alpha$  settings.




Fig. 13. Collected comparisons of HV when using different  $\alpha$  settings.



TABLE I  
PROPOSED TEST PROBLEM

Test problem	Function
MBOP12	$\min_{\mathbf{x}^U \in \Omega^U, \mathbf{x}^L \in \Omega^L} \mathbf{F}(\mathbf{x}^U, \mathbf{x}^L) = \begin{pmatrix} f_1(\mathbf{x}) = (1 + r - \cos(\alpha\pi x_1^U)) + \sum_{j=2}^K (x_j^U - \frac{j-1}{2})^2 + \tau \sum_{i=2}^K (x_i^L - x_i^U)^2 - \\ r \cos(\gamma \frac{\pi}{2} \frac{q(x_1^L) \sin \frac{\pi x_1^L}{2} - q(x_1^L) \cos \frac{\pi x_1^L}{2}}{x_1^U}) \\ f_2(\mathbf{x}) = (1 + r - \sin(\alpha\pi x_1^U)) + \sum_{j=2}^K (x_j^U - \frac{j-1}{2})^2 + \tau \sum_{i=2}^K (x_i^L - x_i^U)^2 + \\ r \sin(\gamma \frac{\pi}{2} \frac{q(x_1^L) \sin \frac{\pi x_1^L}{2} - q(x_1^L) \cos \frac{\pi x_1^L}{2}}{x_1^U}) \end{pmatrix}$ $\text{subject to } \mathbf{x}^L \in \underset{\mathbf{x}^L \in \Omega^L}{\text{argmin}} \begin{pmatrix} f_1^L(\mathbf{x}) = q(x_1^L) \sin \frac{\pi x_1^L}{2} + \sum_{i=2}^K (x_i^L - x_i^U)^2 + \sum_{i=2}^K 10(1 - \cos(\frac{\pi}{K}(x_i^L - x_j^U))) \\ f_2^L(\mathbf{x}) = q(x_1^L) \cos \frac{\pi x_1^L}{2} - x_1^U + \sum_{i=2}^K 10  \sin(\frac{\pi}{K}(x_i^L - x_j^U))  \end{pmatrix}$ $q(x_1^L) = 5 + 10(x_1^L - 0.5)^2 + \frac{2^{s/2}}{\beta} \cos(2\beta\pi x_1^L)$ $1 \leq x_1^U \leq 4, \quad 0 \leq x_1^L \leq 1$ $K \leq x_i^L, x_i^U \leq K, \text{ for } i = 2, \dots, K$ $r = 0.1, \alpha = 1, \tau = 1, \gamma = 1, \beta = 4, s = 0$
MBOP13	$\min_{\mathbf{x}^U \in \Omega^U, \mathbf{x}^L \in \Omega^L} \mathbf{F}(\mathbf{x}^U, \mathbf{x}^L) = \begin{pmatrix} f_1(\mathbf{x}) = l(\mathbf{x}^U) r(x_1^L) \sin^3(\frac{\pi x_1^L}{2})(1 + \sum_{j=2}^K (x_j^L)^2) \\ f_2(\mathbf{x}) = l(\mathbf{x}^U) r(x_1^L) \cos(\frac{\pi x_1^L}{2})(1 + \sum_{j=2}^K (x_j^L)^2) \end{pmatrix}$ $\text{subject to } \mathbf{x}^L \in \underset{\mathbf{x}^L \in \Omega^L}{\text{argmin}} \begin{pmatrix} f_1^L(\mathbf{x}) = l(\mathbf{x}^U) r(x_1^L) \sin^3(\frac{\pi x_1^L}{2})(1 + \sum_{j=K+1}^{K+L} (x_j^L)^2) \\ f_2^L(\mathbf{x}) = l(\mathbf{x}^U) r(x_1^L) \cos(\frac{\pi x_1^L}{2})(1 + \sum_{j=K+1}^{K+L} (x_j^L)^2) \end{pmatrix}$ $l(\mathbf{x}^U) = 1 + \frac{9}{n_u - 1} \sum_{i=1}^{n_u} x_i^U$ $r(x_1^L) = 2.5 + 10(x_1^L - 0.5)^2 + \frac{2^{s/2}}{\beta} \cos(2\beta\pi x_1^L)$ $G(\mathbf{x}^U, \mathbf{x}^L) = f_1(\mathbf{x}) + 0.8f_2(\mathbf{x}) \leq 2.8$ $0 \leq x_i^U, x_j^L \leq 1, \text{ for } i = 1, \dots, n_u, j = 1, \dots, K + L$ $s = 1, \beta = 2$
MBOP14	$\min_{\mathbf{x}^U \in \Omega^U, \mathbf{x}^L \in \Omega^L} \mathbf{F}(\mathbf{x}^U, \mathbf{x}^L) = \begin{pmatrix} f_1(\mathbf{x}) = t_1(\mathbf{x}) + \tau T(t_1(\mathbf{x}), t_2(\mathbf{x})) + \alpha \sum_{j=2}^K (x_j^L)^2 \\ f_2(\mathbf{x}) = t_2(\mathbf{x}) + \tau T(t_1(\mathbf{x}), t_2(\mathbf{x})) + \alpha \sum_{j=2}^K (x_j^L)^2 \end{pmatrix}$ $\text{subject to } \mathbf{x}^L \in \underset{\mathbf{x}^L \in \Omega^L}{\text{argmin}} \begin{pmatrix} f_1^L(\mathbf{x}) = t_1(\mathbf{x}) + \alpha(1 + \sum_{j=2}^K (x_j^L)^2) \\ f_2^L(\mathbf{x}) = t_2(\mathbf{x}) + \alpha(1 + \sum_{j=2}^K (x_j^L)^2) \end{pmatrix}$ $t_1(\mathbf{x}) = l(\mathbf{x}^U) r(x_1^L) [\sin(\frac{\pi x_1^L}{2^{s+1}}) + (1 + \frac{2^{s-1}}{2^{s+2}})\pi] + 1$ $t_2(\mathbf{x}) = l(\mathbf{x}^U) r(x_1^L) (\cos(\frac{\pi x_1^L}{2}) + \pi) + 1$ $l(\mathbf{x}^U) = 1 + \frac{9}{n_u - 1} \sum_{i=1}^{n_u} x_i^U$ $r(x_1^L) = 5 + 10(x_1^L - 0.5)^2 + \frac{1}{\beta} \cos(2\beta\pi x_1^L) \times 2^{\frac{s}{2}}$ $T(x_1, x_2) = \begin{cases} 0, & \text{if } (x_1, x_2) \text{ satisfy } z(x_1, x_2) \\ x_1 + x_2, & \text{otherwise} \end{cases}$ $z(x_1, x_2) = \{(x_1, x_2) \mid (x_1 \in [0.7, 0.83], x_2 \in [4.3, 5]) \cup (x_1 \in [1.05, 1.172], x_2 \in [1.96, 2.44]) \cup \\ (x_1 \in [1.65, 1.77], x_2 \in [0.8, 1.13]) \cup (x_1 \in [2.86, 3.1], x_2 \in [0.142, 0.3])\}$ $0 \leq x_i^U, x_j^L \leq 1, \text{ for } i = 1, \dots, n_u, j = 1, \dots, K$ $\tau = 1, \alpha = 1, s = 1, \beta = 4$
MBOP15	$\min_{\mathbf{x}^U \in \Omega^U, \mathbf{x}^L \in \Omega^L} \mathbf{F}(\mathbf{x}^U, \mathbf{x}^L) = \begin{pmatrix} f_1(\mathbf{x}) = t_1(\mathbf{x})(1 + \sum_{j=2}^K (x_j^L)^2) - \tau_1 T(t_1(\mathbf{x}), t_2(\mathbf{x})) \\ f_2(\mathbf{x}) = t_2(\mathbf{x})(1 + \sum_{j=2}^K (x_j^L)^2) - \tau_1 T(t_1(\mathbf{x}), t_2(\mathbf{x})) \end{pmatrix}$ $\text{subject to } \mathbf{x}^L \in \underset{\mathbf{x}^L \in \Omega^L}{\text{argmin}} \begin{pmatrix} f_1^L(\mathbf{x}) = (t_1(\mathbf{x}) + \tau_2(1 + T(t_1(\mathbf{x}), t_2(\mathbf{x}))))(1 + \sum_{j=K+1}^{K+L} (x_j^L)^2) \\ f_2^L(\mathbf{x}) = (t_2(\mathbf{x}) + \tau_2(1 + T(t_1(\mathbf{x}), t_2(\mathbf{x}))))(1 + \sum_{j=K+1}^{K+L} (x_j^L)^2) \end{pmatrix}$ $t_1(\mathbf{x}) = l(\mathbf{x}^U) r(x_1^L) \sin \frac{\pi x_1^L}{2}, t_2(\mathbf{x}) = l(\mathbf{x}^U) r(x_1^L) \cos \frac{\pi x_1^L}{2}$ $r(x_1^L) = 5 + 10(x_1^L - 0.5)^2 + \frac{1}{\beta} \cos(2\beta\pi x_1^L) 2^{\frac{s}{2}}$ $T(x_1, x_2) = \begin{cases} 0, & \text{if } (x_1, x_2) \text{ satisfy } z_1(x_1, x_2) \\ 0.5, & \text{if } (x_1, x_2) \text{ satisfy } z_2(x_1, x_2) \\ 1, & \text{otherwise} \end{cases}$ $z_1(x_1, x_2) = \{(x_1, x_2) \mid (x_1 \in [1.13, 1.67], x_2 \in [5.72, 6]) \cup (x_1 \in [2.59, 3], x_2 \in [3.88, 4.15]) \cup \\ (x_1 \in [3.88, 4.24], x_2 \in [2.57, 2.915]) \cup (x_1 \in [5.72, 5.98], x_2 \in [1.15, 1.7])\}$ $z_2(x_1, x_2) = \{(x_1, x_2) \mid (x_1 \in [1.13, 1.67], x_2 \notin [5.72, 6]) \cup (x_1 \in [2.59, 3], x_2 \notin [3.88, 4.15]) \cup \\ (x_1 \in [3.88, 4.24], x_2 \notin [2.57, 2.915]) \cup (x_1 \in [5.72, 5.98], x_2 \notin [1.15, 1.7])\}$ $l(\mathbf{x}^U) = 1 + \sum_{i=1}^{n_u} (x_i^U)^{0.1}$ $\mathbf{G}(\mathbf{x}^U, \mathbf{x}^L) = f_1(\mathbf{x}) + f_2(\mathbf{x}) \leq 8$ $0 \leq x_i^U, x_j^L \leq 1, \text{ for } i = 1, \dots, n_u, j = 1, \dots, K + L$ $\tau_1 = 1, \tau_2 = 5, \beta = 4, s = 1$

TABLE II  
CLASSICAL TEST PROBLEM

Test problem	Function
Classic1	$\max_{\mathbf{x}^U \in \Omega^U, \mathbf{x}^L \in \Omega^L} \mathbf{F}(\mathbf{x}^U, \mathbf{x}^L) = \begin{pmatrix} f_1(\mathbf{x}) = x_1^U + 2x_2^U \\ f_2(\mathbf{x}) = 3x_1^U + x_2^U \end{pmatrix}$ $\text{subject to } \mathbf{x}^L \in \operatorname{argmax}_{\mathbf{x}^L \in \Omega^L} \begin{pmatrix} f_1^L(\mathbf{x}) = x_1^L + 3x_2^L \\ f_2^L(\mathbf{x}) = 2x_1^L + x_2^L \end{pmatrix}$ $\mathbf{G}(\mathbf{x}^U, \mathbf{x}^L) = \{(\mathbf{x}_1^U, \mathbf{x}_2^U) \in \mathbb{R}^2 \mid \mathbf{x}_1^U + \mathbf{x}_2^U \leq 3, x_1^U, x_2^U \geq 0\}$ $\mathbf{g}(\mathbf{x}^U, \mathbf{x}^L) = \{(\mathbf{x}_1^U, \mathbf{x}_2^U, \mathbf{x}_1^L, \mathbf{x}_2^L) \in \mathbb{R}^4 \mid -x_1^U + x_1^L + x_2^L \leq 6, -x_2^U + x_1^L \leq 3, x_1^U + x_2^U + x_2^L \leq 8, x_1^L, x_2^L \geq 0\}$
Classic2	$\min_{\mathbf{x}^U \in \Omega^U, \mathbf{x}^L \in \Omega^L} \mathbf{F}(\mathbf{x}^U, \mathbf{x}^L) = \begin{pmatrix} f_1(\mathbf{x}) = x_1^L + (x_2^L)^2 + x^U + \sin^2(x_1^L + x^U) \\ f_2(\mathbf{x}) = \cos(x_2^L) \cdot (0.1 + x^U) \cdot (\exp(-\frac{x_1^L}{0.1 + x_2^L})) \end{pmatrix}$ $\text{subject to } \mathbf{x}^L \in \operatorname{argmin}_{\mathbf{x}^L \in \Omega^L} \begin{pmatrix} f_1^L(\mathbf{x}) = \frac{(x_1^L - 2)^2 + (x_2^L - 1)^2}{4} + \frac{x_2^L x^U + (5 - x^U)^2}{16} + \sin(\frac{x_2^L}{10}) \\ f_2^L(\mathbf{x}) = \frac{(x_1^L)^2 + (x_2^L - 6)^4 - 2x_1^L x^U - (5 - x^U)^2}{80} \end{pmatrix}$ $\mathbf{G}(\mathbf{x}^U, \mathbf{x}^L) = \{x^U \in \mathbb{R}^1 \mid 0 \leq x^U \leq 10\}$ $\mathbf{g}(\mathbf{x}^U, \mathbf{x}^L) = \{(x^U, x_1^L, x_2^L) \in \mathbb{R}^3 \mid (x_1^L)^2 - x_2^L \leq 0, 5(x_1^L)^2 + x_2^L \leq 10, x_2^L - (5 - x^U/6) \leq 0, x_1^L \geq 0\}$

TABLE III  
MATHEMATICAL DEFINITIONS OF THE PETROLEUM REFINING MBLOP

Petroleum Refining MBLOP	Function
Objective	$\min_{\tau \in \Omega^U, F^c, \mathbf{F}^p, F^s \in \Omega^L} \mathbf{F}(\tau, F^c, \mathbf{F}^p, F^s) = \begin{pmatrix} f_1(\mathbf{x}) = Q^{\text{fue}} \zeta \\ f_2(\mathbf{x}) = -\sum_{i=1}^{n^p} \tau_i F_i^p \end{pmatrix}$ $\text{subject to } (F^c, \mathbf{F}^p, F^s) \in \operatorname{argmin}$ $\left( f_1^L(\mathbf{x}) = -\sum_{i=1}^{n^p} C_i^p F_i^p + C^c F^c + C^s F^s + \left( \sum_{i=1}^{n^p} \tau_i F_i^p \right) \& f_2^L(\mathbf{x}) = Q^{\text{fue}} \right)$
Constraint	$\text{Upper Level: } f_1(\mathbf{x}) \geq 0, f_2(\mathbf{x}) \leq 0$ $\text{Lower Level: } F^c - \sum_{i=1}^{n^p} F_i^p \geq 0, f_2^L(\mathbf{x}) \geq 0$
Variable	$\text{Upper Level: } 1.0 \leq \tau_1 \leq 1.4, \quad 0.6 \leq \tau_2 \leq 1.0, \quad 0.6 \leq \tau_3 \leq 1.0, \quad 0.6 \leq \tau_4 \leq 1.0$ $\text{Lower Level: } 10.1 \leq F_1^p \leq 10.21, \quad 30.01 \leq F_2^p \leq 30.93 \quad 59.6 \leq F_3^p \leq 60.5,$ $26.84 \leq F_4^p \leq 31.52, \quad 448.52 \leq F^c \leq 458.45, \quad 0.35 \leq F^s \leq 0.36$
Related Calculations	$Q^{\text{fue}} = Q^{\text{fst}} + Q^{\text{fur}} = \frac{Q^{\text{pst}}}{\lambda^p} (h^p - h^w) \frac{T^{\text{ftb}} - T^0}{T^{\text{ftb}} - T^s} + \frac{Q^{\text{pfu}}}{\eta^f}$ $\eta^f = \frac{T^{\text{ftf}} - T^s}{T^{\text{ftf}} - T^0}$ $\zeta = \left( \frac{\alpha}{\phi} \right) \left( \frac{C\%}{100} \right)$ $Q^{\text{pst}} = \beta F^s$ $Q^{\text{pfu}} = \gamma \left( F^c + \sum_{i=1}^{n^p} F_i^p \right)$

TABLE IV  
SUMMARY OF NOTATIONS USED IN THE PRACTICAL PETROLEUM REFINING MULTI-OBJECTIVE BILEVEL PROBLEM

Parameter	Description	Value
$\zeta$	the relevant fuel factor	—
$n^p$	the number of oil products	4
$\lambda^p, h^p$	the latent heat and enthalpy of steam	1933.19, 2791.73
$Q^{\text{pst}}$	the heat duty required by steam heating	—
$Q^{\text{pfu}}$	the heat duty required by furnace	—
$h^w$	the enthalpy of the boiler feed water at 100°C	419
$T^{\text{ftb}}$	the flame temperature of the boiler flue gas	1800
$T^0$	the environmental temperature	25
$T^s$	the stack temperature of exhaust gas	160
$T^{\text{ftf}}$	the theoretical flame temperature of the furnace	1800
$\eta^f$	the theoretical efficiency of the furnace	—
$\alpha$	the ratio of molar masses of $\text{CO}_2$ and C	3.67
C%	carbon content	0.865
$\phi$	the net heating value of fuel with a carbon content of C%	3971
$\tau_i$	per unit tax imposed on the oil product $i$ . ( $i = 1, \dots, n^p$ )	—
$F^c, F^s, F_i^p$	the flow rate of crude oil, steam, product $i$ . ( $i = 1, \dots, n^p$ )	—
$C^c, C^s, C_i^p$	the prices of crude, steam, product $i$ . ( $i = 1, \dots, n^p$ )	79.6, 0.14, 103.5, 92.7, 99.0, 96.6
$Q^{\text{fue}}$	the amount of fuel burnt for the crude oil distillation process	—
$Q^{\text{fst}}, Q^{\text{fur}}$	the amount of fuel burnt for the steam boilers and furnace	—
$\gamma$	the ratio of the heat duty required by the furnace and the yield of the crude oil and product flow rate	1
$\beta$	the ratio of the steam heat duty and the steam flow rate	1

TABLE V  
PARAMETER SETTING

Test problem	$N^U$	$T^U$	$N^L$	$T^L$	Variable dimension	Ideal point	Nadir point
MBOP1	400	200	40	40	( $K, K$ )	(0, 0)	(1.1, 1.1)
MBOP2	400	200	40	40	( $K, K$ )	(-0.2388, -0.8378)	(0.8090, 0.0138)
MBOP3	400	200	40	40	( $K, K$ )	(-0.1882, -0.2427)	(1.4, 1)
MBOP4	400	100	40	40	(1, $K+L$ )	(0, 0)	(1, 2)
MBOP5	400	100	40	40	(1, $K+L$ )	(0, 0)	(1, 1.8)
MBOP6	400	100	40	40	(1, $K+L$ )	(0, 0)	(1, 2)
MBOP7	400	100	40	40	(1, $K+L$ )	(0, 0)	(1, 1.8)
MBOP8	400	200	40	40	(1, 2)	(-2, -1)	(-1, 0)
MBOP9	300	100	60	80	(1, $K$ )	(0.5, 0)	(1, 0.5)
MBOP10	600	200	60	40	(1, 2)	(23, 2)	(2, -9)
MBOP11	400	200	40	100	(2, 3)	(-300, -2000)	(-1200, -300)
MBOP12	400	200	40	40	( $K, K$ )	(0, 0)	(1.1, 1.1)
MBOP13	400	200	40	40	( $n_u, K + L$ )	(0.0436, 0.9)	(2.073, 3.433)
MBOP14	400	100	40	40	( $n_u, K$ )	(0.7052, 0.1427)	(3.0957, 4.9934)
MBOP15	400	200	40	60	( $n_u, K + L$ )	(1.1340, 1.1534)	(5.971, 5.9958)
Classic1	200	50	40	40	(2, 2)	(-6, -9)	(-3, -3)
Classic2	200	50	40	40	(1, 2)	(23, 2)	(2, -9)
Petroleum Refining	400	50	40	40	(4, 6)	( $4.9 \times 10^{-4}$ , -138)	( $5.1 \times 10^{-4}$ , -80)

TABLE VI  
THE NONDOMINATED SOLUTIONS OF MBOP11

	$\mathbf{x}^U$	$\mathbf{x}^L$	$f_1(\mathbf{x})$	$f_2(\mathbf{x})$	$f_1^L(\mathbf{x})$	$f_2^L(\mathbf{x})$
Solution given in [13]	(146.2955, 28.9394)	(0, 67.9318, 0)	474.6819	1850.0609	1030.5456	1469.0532
Solution A	(0.5600, 107.5164)	(0, 0, 23.2238)	1037.8790	312.9680	833.1288	526.3208
Solution B	(16.6666, 100)	(0, 0, 29.6665)	1005.6661	468.6654	903.9984	618.6656
Solution C	(92.9000, 64.4244)	(0, 0, 60.1598)	853.1990	1205.5880	1239.4248	1055.7368
Solution D	(146.1023, 29.0719)	(0, 67.6585, 0.2501)	476.1582	1847.6744	1031.4754	1467.5113

TABLE VII  
MEDIAN AND INTERQUARTILE RANGE OF IGD, HV AND IGD+ ON MBOP TEST PROBLEMS.

Test Problem	TKDEA			BLEMO			OMOPSO-BL		
	IGD	HV	IGD+	IGD	HV	IGD+	IGD	HV	IGD+
MBOP1	<b>5.85E-2</b>	<b>7.21E-1</b>	<b>5.66E-2</b>	1.77E+0 <sup>†</sup>	0.00E+0 <sup>†</sup>	1.75E+0 <sup>†</sup>	1.87E-1 <sup>†</sup>	5.37E-1 <sup>†</sup>	1.70E-1 <sup>†</sup>
	<b>1.73E-3</b>	<b>2.81E-3</b>	<b>1.35E-3</b>	2.31E+0	2.18E-2	2.32E+0	8.71E-2	9.67E-2	9.95E-2
MBOP2	<b>4.87E-2</b>	<b>5.12E-1</b>	<b>4.85E-2</b>	3.48E+0 <sup>†</sup>	0.00E+0 <sup>†</sup>	3.48E+0 <sup>†</sup>	3.40E-1 <sup>†</sup>	1.40E-1 <sup>†</sup>	3.40E-1 <sup>†</sup>
	<b>7.94E-3</b>	<b>1.64E-2</b>	<b>8.23E-3</b>	4.15E+0	0.00E+0	4.17E+0	1.86E-1	9.35E-2	1.92E-1
MBOP3	<b>1.57E-1</b>	<b>4.12E-1</b>	<b>1.54E-1</b>	8.16E-1 <sup>†</sup>	6.77E-2 <sup>†</sup>	7.91E-1 <sup>†</sup>	1.51E+0 <sup>†</sup>	0.00E+0 <sup>†</sup>	1.51E+0 <sup>†</sup>
	<b>8.87E-2</b>	<b>9.42E-2</b>	<b>8.15E-2</b>	6.18E-1	1.34E-1	6.32E-1	3.89E-1	0.00E+0	3.89E-1
MBOP4	2.06E-1	2.53E-1	1.89E-1	1.81E-1 $\approx$	3.47E-1 $\ddagger$	1.65E-1 $\ddagger$	<b>3.32E-2<math>\ddagger</math></b>	<b>4.61E-1<math>\ddagger</math></b>	<b>3.06E-2<math>\ddagger</math></b>
	7.02E-2	6.45E-2	6.57E-2	9.24E-2	4.90E-2	6.31E-2	<b>8.45E-3</b>	<b>8.04E-3</b>	<b>7.39E-3</b>
MBOP5	3.70E-1	1.16E-1	3.24E-1	3.96E-1 $\approx$	1.83E-1 $\ddagger$	3.05E-1 $\approx$	<b>2.27E-1<math>\ddagger</math></b>	<b>2.61E-1<math>\ddagger</math></b>	<b>2.06E-1<math>\ddagger</math></b>
	6.51E-2	4.27E-2	5.35E-2	1.44E-1	2.12E-2	6.26E-2	<b>9.65E-3</b>	<b>9.12E-3</b>	<b>8.98E-3</b>
MBOP6	<b>1.49E-1</b>	3.22E-1	<b>1.35E-1</b>	1.89E-1 $\approx$	<b>3.52E-1<math>\approx</math></b>	1.45E-1 $\approx$	3.11E-1 <sup>†</sup>	2.28E-1 <sup>†</sup>	3.09E-1 <sup>†</sup>
	<b>3.45E-2</b>	4.00E-2	<b>3.28E-2</b>	1.31E-1	<b>7.50E-2</b>	8.02E-2	4.64E-2	3.22E-2	4.68E-2
MBOP7	<b>2.81E-1</b>	1.87E-1	<b>2.64E-1</b>	3.65E-1 <sup>†</sup>	<b>2.00E-1<math>\approx</math></b>	2.88E-1 <sup>†</sup>	6.46E-1 <sup>†</sup>	7.26E-3 <sup>†</sup>	6.28E-1 <sup>†</sup>
	<b>2.67E-2</b>	3.30E-2	<b>2.85E-2</b>	1.81E-1	<b>4.72E-2</b>	7.86E-2	1.63E-1	4.11E-2	1.50E-1
MBOP8	9.94E-2	7.60E-3	<b>1.11E-3</b>	<b>4.94E-2<math>\ddagger</math></b>	<b>2.14E-1<math>\ddagger</math></b>	1.13E-2 <sup>†</sup>	8.75E-2 <sup>†</sup>	9.54E-3 $\approx$	1.35E-3 $\approx$
	1.33E-2	4.05E-2	<b>1.88E-3</b>	<b>2.68E-2</b>	<b>5.45E-2</b>	1.13E-2	5.79E-3	2.12E-2	8.11E-4
MBOP9-K2	<b>4.05E-3</b>	<b>8.18E-1</b>	<b>2.35E-3</b>	1.47E-1 <sup>†</sup>	2.89E-1 <sup>†</sup>	8.18E-2 <sup>†</sup>	1.93E-2 <sup>†</sup>	7.29E-1 <sup>†</sup>	5.68E-3 <sup>†</sup>
	<b>6.54E-4</b>	<b>2.81E-3</b>	<b>3.73E-4</b>	1.12E-1	5.38E-1	1.10E-1	1.09E-2	7.78E-2	3.38E-3
MBOP9-K6	<b>6.31E-3</b>	<b>8.09E-1</b>	<b>2.99E-3</b>	1.30E-1 <sup>†</sup>	5.06E-1 <sup>†</sup>	7.29E-2 <sup>†</sup>	2.79E-2 <sup>†</sup>	7.08E-1 <sup>†</sup>	5.69E-3 <sup>†</sup>
	<b>1.38E-3</b>	<b>5.82E-3</b>	<b>7.83E-4</b>	1.33E-1	3.85E-1	7.47E-2	1.59E-2	1.57E-1	5.14E-3
MBOP9-K14	<b>1.85E-2</b>	<b>7.76E-1</b>	8.16E-3	1.78E-1 <sup>†</sup>	4.09E-1 <sup>†</sup>	1.32E-1 <sup>†</sup>	2.91E-2 <sup>†</sup>	7.54E-1 <sup>†</sup>	<b>6.31E-3<math>\ddagger</math></b>
	<b>7.00E-3</b>	<b>2.73E-2</b>	2.50E-3	3.10E-1	5.24E-1	3.16E-1	1.40E-2	6.24E-2	<b>2.91E-3</b>
MBOP10	-	4.11E-1	-	-	3.85E-1 <sup>†</sup>	-	-	<b>5.08E-1<math>\ddagger</math></b>	-
	-	1.58E-2	-	-	9.29E-3	-	-	<b>8.85E-3</b>	-
MBOP11	-	<b>7.19E-1</b>	-	-	3.44E-1 <sup>†</sup>	-	-	6.82E-1 <sup>†</sup>	-
	-	<b>3.72E-1</b>	-	-	2.94E-1	-	-	1.22E-1	-
MBOP12	<b>4.63E-3</b>	<b>7.79E-1</b>	<b>4.04E-3</b>	1.25E+0 <sup>†</sup>	0.00E+0 <sup>†</sup>	1.21E+0 <sup>†</sup>	1.79E-1 <sup>†</sup>	5.23E-1 <sup>†</sup>	1.70E-1 <sup>†</sup>
	<b>1.15E-3</b>	<b>1.36E-3</b>	<b>9.38E-4</b>	1.06E+0	3.74E-2	1.09E+0	4.37E-2	7.86E-2	5.95E-2
MBOP13	<b>8.10E-2</b>	<b>6.18E-1</b>	<b>3.28E-2</b>	9.09E-1 <sup>†</sup>	2.36E-1 <sup>†</sup>	9.04E-1 <sup>†</sup>	7.32E+0 <sup>†</sup>	0.00E+0 <sup>†</sup>	7.30E+0 <sup>†</sup>
	<b>1.89E-2</b>	<b>5.64E-3</b>	<b>6.43E-3</b>	1.25E+0	4.45E-1	1.27E+0	2.26E+0	0.00E+0	2.29E+0
MBOP14	<b>8.74E-1</b>	<b>6.75E-1</b>	<b>2.21E-1</b>	1.59E+1 <sup>†</sup>	0.00E+0 <sup>†</sup>	1.59E+1 <sup>†</sup>	1.11E+1 <sup>†</sup>	0.00E+0 <sup>†</sup>	1.11E+1 <sup>†</sup>
	<b>1.06E-3</b>	<b>5.99E-5</b>	<b>5.62E-5</b>	1.14E+1	0.00E+0	1.14E+1	3.58E+0	0.00E+0	3.58E+0
MBOP15	<b>3.16E-2</b>	<b>4.44E-1</b>	<b>2.56E-2</b>	8.75E-2 $\approx$	4.25E-1 $\approx$	8.55E-2 $\approx$	1.36E+1 <sup>†</sup>	0.00E+0 <sup>†</sup>	1.36E+1 <sup>†</sup>
	<b>8.64E-3</b>	<b>1.72E-3</b>	<b>5.80E-3</b>	-	-	-	5.96E-1	0.00E+0	5.96E-1
	$\dagger/\approx/\ddagger$			10/4/1	11/3/3	11/3/1	13/0/2	13/1/3	11/1/3

<sup>†</sup> denotes the performance of TKDEA is significantly better than the other peers according to the Wilcoxon signed-rank test at a 0.05 significance level while  $\ddagger$  denotes the opposite case, and  $\approx$  denotes that there is no significant difference between the two algorithms.

The IGD and IGD+ values of MBOP10 and MBOP11 are not available due to the lack of their true PF.

TABLE VIII  
MEAN AND STANDARD DEVIATION OF THE NUMBER OF FUNCTION EVALUATIONS ON MBOP TEST PROBLEMS

Test Problem	TKDEA/ BLEMO			OMOPSO-BL		
	FE	ULFE	LLFE	FE	ULFE	LLFE
MBOP1	3296400(0)	80400(0)	3216000(0)	3531820(17902)	162620(437)	3369200(17465)
MBOP2	3296400(0)	80400(0)	3216000(0)	3517306(13753)	162266(335)	3355040(13418)
MBOP3	3296400(0)	80400(0)	3216000(0)	3856294(40817)	170534(996)	3685760(39821)
MBOP4	1656400(0)	40400(0)	1616000(0)	2143950(63960)	90720(1560)	2053200(62400)
MBOP5	1656400(0)	40400(0)	1616000(0)	1996320(54120)	87120(1320)	1909200(52800)
MBOP6	1656400(0)	40400(0)	1616000(0)	1928588(27599)	85468(673)	1843120(26926)
MBOP7	1656400(0)	40400(0)	1616000(0)	2049620(37622)	88420(918)	1961200(36704)
MBOP8	3296400(0)	80400(0)	3216000(0)	3700904(43292)	166744(1056)	3534160(42236)
MBOP9-K2	2454300(0)	30300(0)	2424000(0)	2857530(54651)	64830(675)	2792700(53976)
MBOP9-K6	2454300(0)	30300(0)	2424000(0)	2874297(56666)	65037(700)	2809260(55966)
MBOP9-K14	2454300(0)	30300(0)	2424000(0)	2907588(49672)	65448(613)	2842140(49059)
MBOP10	4944600(0)	120600(0)	4824000(0)	5827860(87945)	256860(2145)	5571000(85800)
MBOP11	8120400(0)	80400(0)	8040000(0)	8802360(116150)	165960(1150)	8636400(115000)
MBOP12	3296400(0)	80400(0)	3216000(0)	3524932(15429)	162452(376)	3362480(15053)
MBOP13	1656400(0)	40400(0)	1616000(0)	1789024(14516)	82064(354)	1706960(14162)
MBOP14	1648200(0)	40200(0)	1608000(0)	1754430(5311)	81030(130)	1673400(5182)
MBOP15	828200(0)	20200(0)	808000(0)	1043178(24482)	44658(597)	998520(23885)




Fig. 14. Collected comparisons of IGD+ when using different  $\alpha$  settings.

TABLE IX  
MEDIAN AND IQR OF HV ON CLASSICAL1

HV	TKDEA	CA1	CA2
Classic1	<b>4.84E-1</b> <b>(9.73E-3)</b>	4.08E-1 <sup>†</sup> (0.00E+0)	2.50E-1 <sup>†</sup> (0.00E+0)

<sup>†</sup> denotes the performance of TKDEA is significantly better than the other peers according to the Wilcoxon signed-rank test at a 0.05 significance level.

TABLE X  
MEDIAN AND IQR OF HV ON CLASSICAL2

HV	TKDEA	CA3
Classic2	<b>5.11E-1</b> <b>(1.20E-2)</b>	4.61E-1 <sup>†</sup> (0.00E+0)

<sup>†</sup> denotes the performance of TKDEA is significantly better than the other peers according to the Wilcoxon signed-rank test at a 0.05 significance level.

TABLE XI  
MEDIAN AND INTERQUARTILE RANGE OF IGD, HV AND IGD+ ON MBOP TEST PROBLEMS.

Test Problem	Indicator	TKDEA	B-C-TAEA	B-I-DBEA	B-DMOEA-εC
MBOP1	IGD	<b>5.85E-2(1.73E-3)</b>	5.97E-2(3.98E-3)†	5.95E-2(4.74E-3)≈	6.01E-2(5.36E-3)†
	HV	<b>7.21E-1(2.81E-3)</b>	7.19E-1(3.34E-3)≈	7.17E-1(4.52E-3)†	7.20E-1(6.81E-3)≈
	IGD+	5.66E-2(1.35E-3)	5.73E-2(2.12E-3)≈	5.73E-2(4.36E-3)≈	<b>5.63E-2(3.94E-3)≈</b>
MBOP2	IGD	4.87E-2(7.94E-3)	<b>2.54E-2(5.24E-3)‡</b>	3.39E-2(7.48E-3)‡	2.87E-2(6.45E-3)‡
	HV	5.12E-1(1.64E-2)	<b>5.57E-1(1.02E-2)‡</b>	5.42E-1(1.30E-2)‡	5.53E-1(1.10E-2)‡
	IGD+	4.85E-2(8.23E-3)	<b>2.51E-2(5.25E-3)‡</b>	3.35E-2(7.40E-3)‡	2.82E-2(6.40E-3)‡
MBOP3	IGD	<b>1.57E-1(8.87E-2)</b>	2.45E-1(3.12E-1)†	2.35E-1(2.07E-1)†	NaN†
	HV	<b>4.12E-1(9.42E-2)</b>	3.71E-1(2.57E-1)†	3.38E-1(1.23E-1)†	NaN†
	IGD+	<b>1.54E-1(8.15E-2)</b>	2.33E-1(3.06E-1)†	2.33E-1(1.77E-1)†	NaN†
MBOP4	IGD	2.06E-1(7.02E-2)	1.63E-1(2.46E-2)‡	<b>1.50E-1(4.02E-2)‡</b>	2.00E-1(5.38E-2)≈
	HV	2.53E-1(6.45E-2)	3.08E-1(4.48E-2)‡	<b>3.16E-1(3.72E-2)‡</b>	2.67E-1(6.01E-2)≈
	IGD+	1.89E-1(6.57E-2)	1.52E-1(3.16E-2)‡	<b>1.41E-1(3.62E-2)‡</b>	1.86E-1(4.77E-2)≈
MBOP5	IGD	3.70E-1(6.51E-2)	3.16E-1(4.13E-2)‡	<b>3.07E-1(3.00E-2)‡</b>	3.53E-1(5.25E-2)≈
	HV	1.13E-1(4.27E-2)	1.51E-1(4.04E-2)‡	<b>1.59E-1(2.98E-2)‡</b>	1.21E-1(4.19E-2)≈
	IGD+	3.24E-1(5.35E-2)	2.92E-1(3.69E-2)≈	<b>2.86E-1(2.93E-2)‡</b>	3.21E-1(3.92E-2)≈
MBOP6	IGD	1.49E-1(3.45E-2)	<b>1.42E-1(2.60E-2)≈</b>	1.61E-1(3.87E-2)≈	1.84E-1(4.77E-2)†
	HV	3.22E-1(4.00E-2)	3.22E-1(3.49E-2)≈	<b>3.29E-1(5.46E-2)≈</b>	2.85E-1(3.79E-2)†
	IGD+	1.35E-1(3.28E-2)	<b>1.34E-1(2.66E-2)≈</b>	1.47E-1(3.66E-2)≈	1.73E-1(4.40E-2)†
MBOP7	IGD	<b>2.81E-1(2.67E-2)</b>	2.95E-1(3.02E-2)≈	3.06E-1(5.73E-2)†	3.48E-1(4.21E-2)†
	HV	<b>1.87E-1(3.30E-2)</b>	1.64E-1(3.47E-2)†	1.80E-1(2.66E-2)≈	1.30E-1(3.78E-2)‡
	IGD+	<b>2.64E-1(2.85E-2)</b>	2.84E-1(3.17E-2)≈	2.85E-1(4.83E-2)†	3.14E-1(3.37E-2)†
MBOP8	IGD	<b>9.94E-2(1.33E-2)</b>	1.09E-1(1.25E-2)≈	1.14E-1(3.93E-3)†	8.18E-1(3.95E-1)†
	HV	<b>7.60E-3(4.05E-2)</b>	0.00E+0(1.12E-3)†	0.00E+0(1.68E-3)†	0.00E+0(0.00E+0)≈
	IGD+	1.11E-3(1.88E-3)	2.75E-4(2.01E-3)‡	<b>2.67E-5(4.61E-5)‡</b>	5.70E-1(4.27E-1)†
MBOP9-K2	IGD	<b>4.05E-3(6.54E-4)</b>	4.31E-3(1.01E-3)†	6.97E-3(1.64E-3)†	4.70E-3(1.58E-3)†
	HV	<b>8.18E-1(2.81E-3)</b>	8.16E-1(5.01E-3)†	8.05E-1(6.75E-3)†	8.17E-1(4.60E-3)≈
	IGD+	<b>2.35E-3(3.73E-4)</b>	2.47E-3(4.27E-4)≈	3.22E-3(5.07E-4)†	2.62E-3(5.63E-4)†
MBOP9-K6	IGD	<b>6.31E-3(1.38E-3)</b>	1.07E-2(2.22E-3)†	3.31E-2(2.53E-2)†	2.93E-2(3.29E-2)≈
	HV	<b>8.09E-1(5.82E-3)</b>	7.78E-1(4.11E-2)†	7.08E-1(1.61E-1)†	7.45E-1(6.88E-2)†
	IGD+	<b>2.99E-3(7.83E-4)</b>	3.73E-3(1.14E-3)†	1.11E-2(5.03E-3)†	6.20E-3(5.57E-3)†
MBOP9-K14	IGD	<b>1.85E-2(7.00E-3)</b>	3.75E-2(1.33E-2)†	2.30E-1(1.27E-1)†	5.26E-2(2.57E-2)†
	HV	<b>7.76E-1(2.73E-2)</b>	6.96E-1(4.38E-2)†	1.44E-1(2.01E-1)†	6.87E-1(5.89E-2)†
	IGD+	<b>8.16E-3(2.50E-3)</b>	1.83E-2(7.60E-3)†	2.18E-1(1.41E-1)†	2.48E-2(1.20E-2)†
MBOP10	IGD	-	-	-	-
	HV	4.11E-1(1.58E-2)	<b>5.51E-1(3.21E-3)‡</b>	5.19E-1(1.66E-2)‡	1.08E-1(4.74E-2)≈
	IGD+	-	-	-	-
MBOP11	IGD	-	-	-	-
	HV	7.19E-1(3.72E-1)	6.11E-1(3.24E-1)†	5.59E-1(1.63E-1)†	<b>7.28E-1(8.40E-2)≈</b>
	IGD+	-	-	-	-
†/≈/‡			15/10/10	18/6/11	18/13/4

† denotes the performance of TKDEA is significantly better than the other peers according to the Wilcoxon signed-rank test at a 0.05 significance level while ‡ denotes the opposite case, and ≈ denotes that there is no significant difference between the two algorithms.

The IGD and IGD+ values of MBOP10 and MBOP11 are not available due to the lack of their true PF.