

1 **Patterned progression of gut microbiota**
2 **predisposes preterm infants to necrotizing**
3 **enterocolitis and late onset sepsis: pilot**
4 **data from a non-Western population**

5 **Jiayi Liu¹, Jianhua Sun², Yuqing Li³, Yi Feng⁴, Liya Pan⁵, Zhoulonglong**
6 **Xie⁶, Zhilong Yan⁷, Jianhua Zhao⁸, and Li Hong⁹**

7 ¹**Department of Clinical Nutrition, Shanghai Children's Medical Center, School of**
8 **Medicine Shanghai Jiao Tong University, Shanghai, China**

9 ²**Department of Clinical Nutrition, Shanghai Children's Medical Center, School of**
10 **Medicine Shanghai Jiao Tong University, Shanghai, China**

11 ³**Department of Clinical Nutrition, Shanghai Children's Medical Center, School of**
12 **Medicine Shanghai Jiao Tong University, Shanghai, China**

13 ⁴**Department of Clinical Nutrition, Shanghai Children's Medical Center, School of**
14 **Medicine Shanghai Jiao Tong University, Shanghai, China**

15 ⁵**Department of Clinical Nutrition, Shanghai Children's Medical Center, School of**
16 **Medicine Shanghai Jiao Tong University, Shanghai, China**

17 ⁶**Department of Clinical Nutrition, Shanghai Children's Medical Center, School of**
18 **Medicine Shanghai Jiao Tong University, Shanghai, China**

19 ⁷**Department of Clinical Nutrition, Shanghai Children's Medical Center, School of**
20 **Medicine Shanghai Jiao Tong University, Shanghai, China**

21 ⁸**Shanghai Majorbio Bio-Pharm Technology Co., Ltd, Shanghai, China**

22 ⁹**Department of Clinical Nutrition, Shanghai Children's Medical Center, School of**
23 **Medicine Shanghai Jiao Tong University, Shanghai, China**

24 Corresponding author:

25 Li Hong⁹

26 Email address: hongli@scmc.com.cn

27 **ABSTRACT**

Background and Objectives

Intestinal microbiota dysbiosis might predispose preterm infants to necrotizing enterocolitis(NEC) and late onset sepsis(LOS). In this observational prospective study, we aimed to profile and compare postpartum microbiota progression patterns in non-Western preterm patients with either condition.

Methods

We enrolled preterm infants with gestational age less than 33 weeks and birth weight more than 950g, from July 2013 to December 2014. We began fecal sample collection from the the first stool after birth and prospectively collected until discharge. Bacterial V3 V4 region of 16s rRNA genes from each stool sample were amplified, sequenced and analyzed.

Results

A total of 192 fecal samples from 24 patients were studied, of whom four developed NEC, three LOS; the remaining 17 were used as controls. [The post-partum gut microbiota colonization started to diverge among NEC, LOS and their matched control groups, from the second week after birth. Microbiota of the LOS infants was the least diversified (Shannon index=1.66), while that of the control group was the most diversified(Shannon index=0.88, $p=0.01$). Potentially pathogenic genus *Enterococcus* (20.86%) and *Staphylococcus* (8.67%) were prominent in NEC patients and *Klebsiella* (42.15%) in LOS group. Both two groups addressed lower proportion of *Lactococcus* (7.98% and 13.76% in NEC and LOS group, respectively) than the control group (3.66%).]

Conclusions

Postpartum colonization pattern of gut microbiome might predispose preterm newborns to NEC or LOS, in which reduced diversity of the whole microbiota community and potentially pathogenic genus could have played an essential role in disease progression. Still, more studies are needed to identify etiological strains, underlying mechanisms and correspondent microbial patterns.

INTRODUCTION

Gut microbiota is a key contributor to human health and the dysbiosis of which are proven to be associated with diseases, such as atherosclerosis(Tang et al., 2017), obesity(Bouter et al., 2017), neuropathy(Sarkar et al., 2016), liver diseases(Tilg et al., 2016), etc. Temporal colonization pattern of the intestinal microbiota during early stages of life also provided evidence of its association with early life events, including Type 1 diabetes(Giongo et al., 2011; Vatanen et al., 2018), asthma(Stokholm et al., 2018) and allergy(Madan et al., 2012; Savage et al., 2018). In light of interindividual variability in gut maturity, innate immunity, birth mode and environmental exposures, preterm infants are predisposed to complications post-partum intestinal microbiota development

In preterm infants, necrotizing enterocolitis and late onset sepsis

Necrotizing enterocolitis, characterized by rapid progression, high morbidity and mortality, is one of the most devastating gastrointestinal neonatal emergencies, especially in preterm newborns. Previous studies have suggested well how intestinal microbiota pattern is implicated prior to, during and after the course of this condition. Mai et al. reported an increase in the Proteobacteria and a decrease in the Firmicutes phyla during three to seven days prior to NEC onset (Mai et al., 2011). Zhou et al. found a relatively higher abundance of *Clostridium* and Gamma-Proteobacteria in the proximity of NEC during early and late onset, respectively(Zhou et al., 2015).

late onset sepsis is

Among non-Western population, however, evidence (few studies on?) of microbiota chronological dysbiosis preceding necrotizing enterocolitis or late onset sepsis has been scant so far. Hence, conducting a prospective study among Chinese patients using high-throughput DNA sequencing, we aimed to profile and compare post-postpartum pattern of intestinal microbiota in preterm infants who subsequently developed necrotizing enterocolitis and late onset sepsis, which may be critical in the etiopathogenesis of disease.

METHODS

Ethics

This study was approved by the joint committee of ethics of Shanghai Children's Medical Center, School of Medicine Shanghai Jiao Tong University (SCMCIRB-K2013022). Detailed written informed consent was obtained from the parents prior to fecal sample collection.

Patients

Newly born preterm infants with gestational age less than 33 weeks, birth weight over 950g were enrolled from Neonatal Intensive Care Unit at Shanghai Children's Medical Center from July 2013 to December 2014. The exclusion criteria were 1) diagnosed with early-onset sepsis, 2) hepatic diseases, 3) renal impairment ($\text{Cr} > 88 \mu\text{M}$), 4) diagnosed with intestinal obstruction, 5) in foreseeable need of cardiovascular or abdominal surgeries (except for male circumcision or PDA ligation), 6) estimated parenteral support to supply over 50% of daily caloric intake for more than four days, 7) given intravenous antibiotics administration (except prophylactic regimen of cefotaxime, piperacillin-tazobactam and/or metronidazole), 8) history of oral antibiotics administration, 9) grossly bloody stools at admission, and 10) over five days old.

NEC cases were defined as infants who met the criteria for Stage II and Stage III NEC diagnosis (Bell et al., 1978), including radiographic intestinal dilation, ileus, pneumatosis intestinalis, and/or absent bowel sounds with or without abdominal tenderness, and/or mild metabolic acidosis and thrombocytopenia. LOS cases was diagnosed if 1) an infant had a positive hemoculture or other suspicious loci of infection after 72 hours of life, with septic signs/symptoms reviewed independently by at least two neonatologists, and had been treated with advanced antibiotics (e.g., Meropenem) after diagnosis. Infants with no infectious complications or sepsis were regarded as controls.

Sample collection and handling

Fecal samples collection began from neonatal meconium till discharge. Although we intended to collect fecal samples on a daily basis, due to working shifts and flexible clinical scheduling, we set seven days as the maximum interval between two collections from every infant. Every sample was collected within 30 minutes of defecation from patients' diaper with a sterile spatula. The samples were immediately placed in a cryogenic vial on dry ice and stored at 80°C within 30 minutes without additives. All samples were collected and stored before knowing the diagnosis of respective patients.

DNA extraction and quality control amplification and 16s rRNA gene sequencing

Microbial genomic DNA was isolated from each fecal specimen using the E.Z.N.A.® Soil DNA Kit (Omega Bio-Tek, Norcross, GA, U.S.) according to manufacturer's protocols. The concentration and purity of the DNA were determined by NanoDrop 2000 UV-vis spectrophotometer (Thermo Scientific, Wilmington, USA), and the DNA quality was checked by 1% agarose gel electrophoresis.

Broad-range PCR and High-throughput Sequencing of 16s rRNA gene amplicons

The V3-V4 hypervariable regions of the bacterial 16S rRNA gene were amplified from each sample using bacterial/archaeal primers 338F (5'-ACTCCTACGGGAGGCAGCAG-3') and 806R (5'-GGACTACHVGG GTWTCTAAT-3') using thermocycler PCR system (GeneAmp 9700, ABI, USA). The PCR reactions were as follows: 3 min of denaturation at 95°C , 27 cycles of 30 s at 95°C , 30 s annealing at 55°C and 45 s elongation at 72°C , and a final extension at 72°C for 10 min. The PCR reactions were performed in triplicate, with each 20 μL mixture containing 4 μL 5X FastPfu Buffer, 2 μL 2.5 mM dNTPs, 0.8 μL of each primer (5 μM), 0.4 μL FastPfu Polymerase and 10 ng template DNA. The PCR products were extracted from a 2% agarose gel and further purified using the AxyPrep DNA Gel Extraction Kit (Axygen Biosciences, Union City, CA, USA), and quantified using QuantiFluor™-ST (Promega, USA) according to the manufacturer's protocols.

Equimolar amounts of purified amplicons were pooled and paired-end sequenced (2 x 300) on an Illumina MiSeq platform (Illumina, San Diego, USA) according to the standard protocols of Majorbio Bio-Pharm Technology Co. Ltd. (Shanghai, China). The reads were de-multiplexed using the Illumina software and separate FASTQ files were generated for each specimen and deposited to the Sequence Read Archive NCBI under the BioProject accession PRJNA470548. Another public archive repository is available at figshare doi: 10.6084/m9.figshare.7205102

Raw Data Processing

After pyrosequencing, de-multiplexed sequence reads were subjected to quality filtering utilizing Trimmomatic software (Bolger et al., 2014) (version????), and were truncated at any site with an Phred score < 20 over a 50bp-sized window; barcode matching with the primer mismatch from 0 to 2 nucleotides was adopted and reads containing ambiguous characters were removed. After trimming, FLASH (Fast Length Adjustment of Short Read) (Magoč and Salzberg, 2011), a read pre-processing software, assembled and

merged the paired-end reads from fragments and generated >10 bp overlapped, with the dead match ratio 0.2. Unassembled reads were discarded. As a result, 7,472,400 V3-V4 tags of 16S rRNA gene were produced, with an average length of 448 bp.(Table S1)

To fairly compare all the samples at the same sequencing depth, the "sub.sample" command of mothur program(version1.30.1)(Schloss et al., 2009) was used for normalization to the smallest sample size. UCHIME Algorithm detected chimeric sequences, removed chimera to obtain effective reads, which were then sorted by cluster size and processed using Operational Taxonomic Units(OTUs) were clustered with 97% similarity cutoff using USEARCH v7(UPARSE version 7.1). The taxonomy of each 16S rRNA gene sequence was analyzed by RDP Classifier algorithm(Wang et al., 2007) against the Silva (SSU128)(Quast et al., 2012) 16S rRNA database using confidence threshold of 70%. Each sequence was assigned the taxonomy by QIIME(Caporaso et al., 2010). The representative sequences were allocated phylogenetically down to the domain, phylum, class, order, family, and genus levels. The relative abundance of a given taxonomic group was calculated as a percentage of the sequences number belonging to that group divided by the total number of obtained sequences.

Alpha diversity analysis, including Shannon index and Observed species richness (sobs), were obtained using the "summary.single" command of mothur program(version1.30.1).

Statistical and Bioinformatics Analysis

Demographics and Clinical Sample comparisons

Non-parametric tests, including Kruskal-Wallis test and Wilcoxon rank-sum test were used to compare gestational age, birth weight, age when the patients were diagnosed as NEC or LOS and length of stay at α level of 0.05 among three groups. All statistical test not involving microbiome 16S rRNA sequencing data was performed using "stats" package using R(v.3.5.1).

Microbiota and Bioinformatics Analyses

Figures

Figures were generated with the "ggpubr"(Kassambara, 2017) and "ggplot2"(Wickham, 2016) packages using R(v.3.5.1)

RESULTS

Patients and samples characteristics

Totally 130 infants met the criteria of our study, and 1698 samples were collected from them. We studied 192 samples from 24 well-sampled preterm infants, including four subsequently diagnosed with NEC (2 in stage IIA and 2 in stage IIB), three with LOS, and 17 matched controls (Table S2). Comparisons showed no significant difference in terms of gestational age, birth weight and gender proportions, diagnosed age among three groups (Table 1). Length of stay among three groups was significantly different however rational since NEC and LOS patients usually require longer period of healthcare because of their worse conditions compared with the control group. All infants were delivered by cesarean section and fed on infant formula. No one was prescribed probiotics during hospitalization.

DISCUSSION

Previous studies have

CONCLUSIONS

Necrotizing enterocolitis, a worldwide concern that threatens

We acknowledge that the sample size is limited since this study is single-center-based and the incidence of both diseases are relatively low: among the 1148 preterm infants admitted within July 2013 to December 2014, only five developed NEC.

ACKNOWLEDGMENTS

We appreciate the support from enrolled patients, their families, and all staffs at Shanghai Children's Medical Center.

178 **SOME L^AT_EX EXAMPLES**

179 Use section and subsection commands to organize your document. L^AT_EX handles all the formatting and
180 numbering automatically. Use ref and label commands for cross-references.

181 **Figures and Tables**

182 Use the table and tabular commands for basic tables — see Table 1, for example. You can upload a figure
183 (JPEG, PNG or PDF) using the project menu. To include it in your document, use the includegraphics
184 command as in the code for Figure 1 below.



Figure 1. An example image.

Item	Quantity
Widgets	42
Gadgets	13

Table 1. An example table.

185 **Citations**

186 LaTeX formats citations and references automatically using the bibliography records in your .bib file,
187 which you can edit via the project menu. Use the cite command for an inline citation, like ?, and the citep
188 command for a citation in parentheses (?).

189 **Mathematics**

L^AT_EX is great at typesetting mathematics. Let X_1, X_2, \dots, X_n be a sequence of independent and identically distributed random variables with $E[X_i] = \mu$ and $\text{Var}[X_i] = \sigma^2 < \infty$, and let

$$S_n = \frac{X_1 + X_2 + \dots + X_n}{n} = \frac{1}{n} \sum_i^n X_i$$

190 denote their mean. Then as n approaches infinity, the random variables $\sqrt{n}(S_n - \mu)$ converge in distribution
191 to a normal $\mathcal{N}(0, \sigma^2)$.

192 Lists

193 You can make lists with automatic numbering ...

- 194 1. Like this,
- 195 2. and like this.

196 ... or bullet points ...

- 197 • Like this,
- 198 • and like this.

199 ... or with words and descriptions ...

200 **Word** Definition

201 **Concept** Explanation

202 **Idea** Text

203 We hope you find write \LaTeX useful for your PeerJ submission, and please let us know if you have any
204 feedback. Further examples with dummy text are included in the following pages.

205 METHODS

206 Quisque ullamcorper placerat ipsum. Cras nibh. Morbi vel justo vitae lacus tincidunt ultrices. Lorem
207 ipsum dolor sit amet, consectetur adipiscing elit. In hac habitasse platea dictumst. Integer tempus
208 convallis augue. Etiam facilisis. Nunc elementum fermentum wisi. Aenean placerat. Ut imperdiet, enim
209 sed gravida sollicitudin, felis odio placerat quam, ac pulvinar elit purus eget enim. Nunc vitae tortor.
210 Proin tempus nibh sit amet nisl. Vivamus quis tortor vitae risus porta vehicula.

$$\cos^3 \theta = \frac{1}{4} \cos \theta + \frac{3}{4} \cos 3\theta \quad (1)$$

211 Fusce mauris. Vestibulum luctus nibh at lectus. Sed bibendum, nulla a faucibus semper, leo velit
212 ultricies tellus, ac venenatis arcu wisi vel nisl. Vestibulum diam. Aliquam pellentesque, augue quis sagittis
213 posuere, turpis lacus congue quam, in hendrerit risus eros eget felis. Maecenas eget erat in sapien mattis
214 porttitor. Vestibulum porttitor. Nulla facilisi. Sed a turpis eu lacus commodo facilisis. Morbi fringilla,
215 wisi in dignissim interdum, justo lectus sagittis dui, et vehicula libero dui cursus dui. Mauris tempor
216 ligula sed lacus. Duis cursus enim ut augue. Cras ac magna. Cras nulla. Nulla egestas. Curabitur a leo.
217 Quisque egestas wisi eget nunc. Nam feugiat lacus vel est. Curabitur consectetur.

218 Subsection

219 Suspendisse vel felis. Ut lorem lorem, interdum eu, tincidunt sit amet, laoreet vitae, arcu. Aenean faucibus
220 pede eu ante. Praesent enim elit, rutrum at, molestie non, nonummy vel, nisl. Ut lectus eros, malesuada
221 sit amet, fermentum eu, sodales cursus, magna. Donec eu purus. Quisque vehicula, urna sed ultricies
222 auctor, pede lorem egestas dui, et convallis elit erat sed nulla. Donec luctus. Curabitur et nunc. Aliquam
223 dolor odio, commodo pretium, ultricies non, pharetra in, velit. Integer arcu est, nonummy in, fermentum
224 faucibus, egestas vel, odio.

225 **Paragraph** Sed commodo posuere pede. Mauris ut est. Ut quis purus. Sed ac odio. Sed vehicula
226 hendrerit sem. Duis non odio. Morbi ut dui. Sed accumsan risus eget odio. In hac habitasse platea
227 dictumst. Pellentesque non elit. Fusce sed justo eu urna porta tincidunt. Mauris felis odio, sollicitudin sed,
228 volutpat a, ornare ac, erat. Morbi quis dolor. Donec pellentesque, erat ac sagittis semper, nunc dui lobortis
229 purus, quis congue purus metus ultricies tellus. Proin et quam. Class aptent taciti sociosqu ad litora
230 torquent per conubia nostra, per inceptos hymenaeos. Praesent sapien turpis, fermentum vel, eleifend
231 faucibus, vehicula eu, lacus.

232 **Paragraph** Pellentesque habitant morbi tristique senectus et netus et malesuada fames ac turpis egestas.
233 Donec odio elit, dictum in, hendrerit sit amet, egestas sed, leo. Praesent feugiat sapien aliquet odio.
234 Integer vitae justo. Aliquam vestibulum fringilla lorem. Sed neque lectus, consectetur at, consectetur
235 sed, eleifend ac, lectus. Nulla facilisi. Pellentesque eget lectus. Proin eu metus. Sed porttitor. In hac
236 habitasse platea dictumst. Suspendisse eu lectus. Ut mi mi, lacinia sit amet, placerat et, mollis vitae, dui.
237 Sed ante tellus, tristique ut, iaculis eu, malesuada ac, dui. Mauris nibh leo, facilisis non, adipiscing quis,
238 ultrices a, dui.

239 **Subsection**

240 Morbi luctus, wisi viverra faucibus pretium, nibh est placerat odio, nec commodo wisi enim eget quam.
241 Quisque libero justo, consectetur a, feugiat vitae, porttitor eu, libero. Suspendisse sed mauris vitae
242 elit sollicitudin malesuada. Maecenas ultricies eros sit amet ante. Ut venenatis velit. Maecenas sed mi
243 eget dui varius euismod. Phasellus aliquet volutpat odio. Vestibulum ante ipsum primis in faucibus orci
244 luctus et ultrices posuere cubilia Curae; Pellentesque sit amet pede ac sem eleifend consectetur. Nullam
245 elementum, urna vel imperdiet sodales, elit ipsum pharetra ligula, ac pretium ante justo a nulla. Curabitur
246 tristique arcu eu metus. Vestibulum lectus. Proin mauris. Proin eu nunc eu urna hendrerit faucibus.
247 Aliquam auctor, pede consequat laoreet varius, eros tellus scelerisque quam, pellentesque hendrerit ipsum
248 dolor sed augue. Nulla nec lacus.




Figure 2. In-text Picture

249 Reference to Figure 2.

250 **RESULTS AND DISCUSSION**

251 Suspendisse vitae elit. Aliquam arcu neque, ornare in, ullamcorper quis, commodo eu, libero. Fusce
252 sagittis erat at erat tristique mollis. Maecenas sapien libero, molestie et, lobortis in, sodales eget, dui.
253 Morbi ultrices rutrum lorem. Nam elementum ullamcorper leo. Morbi dui. Aliquam sagittis. Nunc
254 placerat. Pellentesque tristique sodales est. Maecenas imperdiet lacinia velit. Cras non urna. Morbi eros

255 pede, suscipit ac, varius vel, egestas non, eros. Praesent malesuada, diam id pretium elementum, eros sem
256 dictum tortor, vel consectetur odio sem sed wisi.

257 Subsection

258 Sed feugiat. Cum sociis natoque penatibus et magnis dis parturient montes, nascetur ridiculus mus. Ut
259 pellentesque augue sed urna. Vestibulum diam eros, fringilla et, consectetur eu, nonummy id, sapien.
260 Nullam at lectus. In sagittis ultrices mauris. Curabitur malesuada erat sit amet massa. Fusce blandit.
261 Aliquam erat volutpat. Aliquam euismod. Aenean vel lectus. Nunc imperdiet justo nec dolor.

262 Subsubsection

263 Etiam euismod. Fusce facilisis lacinia dui. Suspendisse potenti. In mi erat, cursus id, nonummy sed,
264 ullamcorper eget, sapien. Praesent pretium, magna in eleifend egestas, pede pede pretium lorem, quis
265 consectetur tortor sapien facilisis magna. Mauris quis magna varius nulla scelerisque imperdiet. Aliquam
266 non quam. Aliquam porttitor quam a lacus. Praesent vel arcu ut tortor cursus volutpat. In vitae pede quis
267 diam bibendum placerat. Fusce elementum convallis neque. Sed dolor orci, scelerisque ac, dapibus nec,
268 ultricies ut, mi. Duis nec dui quis leo sagittis commodo.

269 Subsubsection

270 Etiam ac leo a risus tristique nonummy. Donec dignissim tincidunt nulla. Vestibulum rhoncus molestie
271 odio. Sed lobortis, justo et pretium lobortis, mauris turpis condimentum augue, nec ultricies nibh arcu
272 pretium enim. Nunc purus neque, placerat id, imperdiet sed, pellentesque nec, nisl. Vestibulum imperdiet
273 neque non sem accumsan laoreet. In hac habitasse platea dictumst. Etiam condimentum facilisis libero.
274 Suspendisse in elit quis nisl aliquam dapibus. Pellentesque auctor sapien. Sed egestas sapien nec lectus.
275 Pellentesque vel dui vel neque bibendum viverra. Aliquam porttitor nisl nec pede. Proin mattis libero vel
276 turpis. Donec rutrum mauris et libero. Proin euismod porta felis. Nam lobortis, metus quis elementum
277 commodo, nunc lectus elementum mauris, eget vulputate ligula tellus eu neque. Vivamus eu dolor.

278 Subsection

279 Nulla in ipsum. Praesent eros nulla, congue vitae, euismod ut, commodo a, wisi. Pellentesque habitant
280 morbi tristique senectus et netus et malesuada fames ac turpis egestas. Aenean nonummy magna non leo.
281 Sed felis erat, ullamcorper in, dictum non, ultricies ut, lectus. Proin vel arcu a odio lobortis euismod.
282 Vestibulum ante ipsum primis in faucibus orci luctus et ultrices posuere cubilia Curae; Proin ut est.
283 Aliquam odio. Pellentesque massa turpis, cursus eu, euismod nec, tempor congue, nulla. Duis viverra
284 gravida mauris. Cras tincidunt. Curabitur eros ligula, varius ut, pulvinar in, cursus faucibus, augue.

285 Nulla mattis luctus nulla. Duis commodo velit at leo. Aliquam vulputate magna et leo. Nam
286 vestibulum ullamcorper leo. Vestibulum condimentum rutrum mauris. Donec id mauris. Morbi molestie
287 justo et pede. Vivamus eget turpis sed nisl cursus tempor. Curabitur mollis sapien condimentum nunc. In
288 wisi nisl, malesuada at, dignissim sit amet, lobortis in, odio. Aenean consequat arcu a ante. Pellentesque
289 porta elit sit amet orci. Etiam at turpis nec elit ultricies imperdiet. Nulla facilisi. In hac habitasse platea
290 dictumst. Suspendisse viverra aliquam risus. Nullam pede justo, molestie nonummy, scelerisque eu,
291 facilisis vel, arcu.

292 Curabitur tellus magna, porttitor a, commodo a, commodo in, tortor. Donec interdum. Praesent
293 scelerisque. Maecenas posuere sodales odio. Vivamus metus lacus, varius quis, imperdiet quis, rhoncus a,
294 turpis. Etiam ligula arcu, elementum a, venenatis quis, sollicitudin sed, metus. Donec nunc pede, tincidunt
295 in, venenatis vitae, faucibus vel, nibh. Pellentesque wisi. Nullam malesuada. Morbi ut tellus ut pede
296 tincidunt porta. Lorem ipsum dolor sit amet, consectetur adipiscing elit. Etiam congue neque id dolor.

297 Donec et nisl at wisi luctus bibendum. Nam interdum tellus ac libero. Sed sem justo, laoreet vitae,
298 fringilla at, adipiscing ut, nibh. Maecenas non sem quis tortor eleifend fermentum. Etiam id tortor ac
299 mauris porta vulputate. Integer porta neque vitae massa. Maecenas tempus libero a libero posuere dictum.
300 Vestibulum ante ipsum primis in faucibus orci luctus et ultrices posuere cubilia Curae; Aenean quis mauris
301 sed elit commodo placerat. Class aptent taciti sociosqu ad litora torquent per conubia nostra, per inceptos
302 hymenaeos. Vivamus rhoncus tincidunt libero. Etiam elementum pretium justo. Vivamus est. Morbi a
303 tellus eget pede tristique commodo. Nulla nisl. Vestibulum sed nisl eu sapien cursus rutrum.

304 Nulla non mauris vitae wisi posuere convallis. Sed eu nulla nec eros scelerisque pharetra. Nullam
305 varius. Etiam dignissim elementum metus. Vestibulum faucibus, metus sit amet mattis rhoncus, sapien
306 dui laoreet odio, nec ultricies nibh augue a enim. Fusce in ligula. Quisque at magna et nulla commodo

consequat. Proin accumsan imperdiet sem. Nunc porta. Donec feugiat mi at justo. Phasellus facilisis ipsum quis ante. In ac elit eget ipsum pharetra faucibus. Maecenas viverra nulla in massa.

Nulla ac nisl. Nullam urna nulla, ullamcorper in, interdum sit amet, gravida ut, risus. Aenean ac enim. In luctus. Phasellus eu quam vitae turpis viverra pellentesque. Duis feugiat felis ut enim. Phasellus pharetra, sem id porttitor sodales, magna nunc aliquet nibh, nec blandit nisl mauris at pede. Suspendisse risus risus, lobortis eget, semper at, imperdiet sit amet, quam. Quisque scelerisque dapibus nibh. Nam enim. Lorem ipsum dolor sit amet, consectetur adipiscing elit. Nunc ut metus. Ut metus justo, auctor at, ultrices eu, sagittis ut, purus. Aliquam aliquam.

ACKNOWLEDGMENTS

So long and thanks for all the fish.

REFERENCES

- Bell, M. J., Ternberg, J. L., Feigin, R. D., Keating, J. P., Marshall, R., Barton, L., and Brotherton, T. (1978). Neonatal necrotizing enterocolitis. therapeutic decisions based upon clinical staging. *Annals of surgery*, 187(1):1.
- Bolger, A. M., Lohse, M., and Usadel, B. (2014). Trimmomatic: a flexible trimmer for illumina sequence data. *Bioinformatics*, 30(15):2114–2120.
- Bouter, K. E., van Raalte, D. H., Groen, A. K., and Nieuwdorp, M. (2017). Role of the gut microbiome in the pathogenesis of obesity and obesity-related metabolic dysfunction. *Gastroenterology*, 152(7):1671–1678.
- Caporaso, J. G., Kuczynski, J., Stombaugh, J., Bittinger, K., Bushman, F. D., Costello, E. K., Fierer, N., Pena, A. G., Goodrich, J. K., Gordon, J. I., et al. (2010). Qiime allows analysis of high-throughput community sequencing data. *Nature methods*, 7(5):335.
- Giongo, A., Gano, K. A., Crabb, D. B., Mukherjee, N., Novelo, L. L., Casella, G., Drew, J. C., Ilonen, J., Knip, M., Hyöty, H., et al. (2011). Toward defining the autoimmune microbiome for type 1 diabetes. *The ISME journal*, 5(1):82.
- Kassambara, A. (2017). ggpubr:“ggplot2” based publication ready plots. r package version 0.1. 6.
- Madan, J. C., Farzan, S. F., Hibberd, P. L., and Karagas, M. R. (2012). Normal neonatal microbiome variation in relation to environmental factors, infection and allergy. *Current opinion in pediatrics*, 24(6):753.
- Magoč, T. and Salzberg, S. L. (2011). Flash: fast length adjustment of short reads to improve genome assemblies. *Bioinformatics*, 27(21):2957–2963.
- Quast, C., Pruesse, E., Yilmaz, P., Gerken, J., Schweer, T., Yarza, P., Peplies, J., and Glöckner, F. O. (2012). The silva ribosomal rna gene database project: improved data processing and web-based tools. *Nucleic acids research*, 41(D1):D590–D596.
- Sarkar, A., Lehto, S. M., Harty, S., Dinan, T. G., Cryan, J. F., and Burnet, P. W. (2016). Psychobiotics and the manipulation of bacteria–gut–brain signals. *Trends in neurosciences*, 39(11):763–781.
- Savage, J. H., Lee-Sarwar, K. A., Sordillo, J., Bunyavanich, S., Zhou, Y., O’connor, G., Sandel, M., Bacharier, L. B., Zeiger, R., Sodergren, E., et al. (2018). A prospective microbiome-wide association study of food sensitization and food allergy in early childhood. *Allergy*, 73(1):145–152.
- Schloss, P. D., Westcott, S. L., Ryabin, T., Hall, J. R., Hartmann, M., Hollister, E. B., Lesniewski, R. A., Oakley, B. B., Parks, D. H., Robinson, C. J., et al. (2009). Introducing mothur: open-source, platform-independent, community-supported software for describing and comparing microbial communities. *Applied and environmental microbiology*, 75(23):7537–7541.
- Stokholm, J., Blaser, M. J., Thorsen, J., Rasmussen, M. A., Waage, J., Vinding, R. K., Schoos, A.-M. M., Kunøe, A., Fink, N. R., Chawes, B. L., et al. (2018). Maturation of the gut microbiome and risk of asthma in childhood. *Nature communications*, 9(1):141.
- Tang, W. W., Kitai, T., and Hazen, S. L. (2017). Gut microbiota in cardiovascular health and disease. *Circulation research*, 120(7):1183–1196.
- Tilg, H., Cani, P. D., and Mayer, E. A. (2016). Gut microbiome and liver diseases. *Gut*, 65(12):2035–2044.
- Vatanen, T., Franzosa, E. A., Schwager, R., Tripathi, S., Arthur, T. D., Vehik, K., Lernmark, Å., Hagopian, W. A., Rewers, M. J., She, J.-X., et al. (2018). The human gut microbiome in early-onset type 1 diabetes from the teddy study. *Nature*, 562(7728):589.

359 Wang, Q., Garrity, G. M., Tiedje, J. M., and Cole, J. R. (2007). Naive bayesian classifier for rapid
360 assignment of rna sequences into the new bacterial taxonomy. *Applied and environmental microbiology*,
361 73(16):5261–5267.
362 Wickham, H. (2016). *ggplot2: Elegant Graphics for Data Analysis*. Springer-Verlag New York.
363 Zhou, Y., Shan, G., Sodergren, E., Weinstock, G., Walker, W. A., and Gregory, K. E. (2015). Longitudinal
364 analysis of the premature infant intestinal microbiome prior to necrotizing enterocolitis: a case-control
365 study. *PloS one*, 10(3):e0118632.