

1 **Patterned progression of gut microbiota in**
2 **association with preterm infants to**
3 **necrotizing enterocolitis and late onset**
4 **sepsis: prospective pilot data from a**
5 **non-Western population**

6 **Jiayi Liu¹, Jianhua Sun², Yuqing Li³, Yi Feng⁴, Liya Pan⁵, Zhoulonglong**
7 **Xie⁶, Zhilong Yan⁷, Jianhua Zhao⁸, and Li Hong⁹**

8 ¹**Department of Clinical Nutrition, Shanghai Children's Medical Center, School of**
9 **Medicine Shanghai Jiao Tong University, Shanghai, China**

10 ²**Department of Clinical Nutrition, Shanghai Children's Medical Center, School of**
11 **Medicine Shanghai Jiao Tong University, Shanghai, China**

12 ³**Department of Clinical Nutrition, Shanghai Children's Medical Center, School of**
13 **Medicine Shanghai Jiao Tong University, Shanghai, China**

14 ⁴**Department of Clinical Nutrition, Shanghai Children's Medical Center, School of**
15 **Medicine Shanghai Jiao Tong University, Shanghai, China**

16 ⁵**Department of Clinical Nutrition, Shanghai Children's Medical Center, School of**
17 **Medicine Shanghai Jiao Tong University, Shanghai, China**

18 ⁶**Department of Clinical Nutrition, Shanghai Children's Medical Center, School of**
19 **Medicine Shanghai Jiao Tong University, Shanghai, China**

20 ⁷**Department of Clinical Nutrition, Shanghai Children's Medical Center, School of**
21 **Medicine Shanghai Jiao Tong University, Shanghai, China**

22 ⁸**Shanghai Majorbio Bio-Pharm Technology Co., Ltd, Shanghai, China**

23 ⁹**Department of Clinical Nutrition, Shanghai Children's Medical Center, School of**
24 **Medicine Shanghai Jiao Tong University, Shanghai, China**

25 Corresponding author:

26 Li Hong⁹

27 Email address: hongli@scmc.com.cn

28 **ABSTRACT**

Background and Objectives

Intestinal microbiota dysbiosis might predispose preterm infants to necrotizing enterocolitis(NEC) and late onset sepsis(LOS). In this observational prospective study, we aimed to profile and compare post-partum microbiota progression patterns in non-Western preterm patients with either condition.

Methods

We enrolled preterm infants with gestational age less than 33 weeks and birth weight more than 950g, from July 2013 to December 2014. We began fecal sample collection from the the first stool after birth and prospectively collected until discharge. Bacterial V3 V4 region of 16s rRNA genes from each stool sample were amplified and sequenced. With the use of RM two-way ANOVA and Zero-Inflated Beta Random Effect models to account for repeated measures, we found out the development of NEC or LOS associated with gut bacterial communities.

Results

A total of 192 fecal samples from 24 patiens were studied, of whom four developed NEC, three LOS; the remaining 17 were used as controls. [The post-partum gut microbiota colonization started to diverge among NEC, LOS and their matched control groups, from the second week after birth. Microbiota of the LOS infants was the least diversified (Shannon index=1.66), while that of the control group was the most diversified(Shannon index=0.88, p=0.01). Potentially pathogenic genus *Enterococcus* (20.86%) and *Staphylococcus* (8.67%) were prominent in NEC patients and *Klebsiella* (42.15%) in LOS group. Both two groups addressed lower proportion of *Lactococcus* (7.98% and 13.76% in NEC and LOS group, respectively) than the control group (3.66%).]

Conclusions

post-partum colonization pattern of gut microbiome might predispose preterm newborns to NEC or LOS, in which reduced diversity of the whole microbiota community and potentially pathogenic genus could have played an essential role in disease progression. Still, more studies are needed to identify etiological strains, underlying mechanisms and correspondent microbial patterns.

INTRODUCTION

Gut microbiota is a key contributor to human health and the dysbiosis of which are proven to be associated with various diseases, such as atherosclerosis(Tang et al., 2017), obesity(Bouter et al., 2017), neuropathy(Sarkar et al., 2016), liver diseases(Tilg et al., 2016), etc. Temporal colonization pattern of the intestinal microbiota during early stages of life also provided evidence of its association with early life events, including Type 1 diabetes(Giongo et al., 2011; Vatanen et al., 2018), asthma(Stokholm et al., 2018) and allergy(Madan et al., 2012; Savage et al., 2018). In light of less gut maturity, innate immunity and more C-sections birth modes, microbiome assembly in pretem infants often differs from that of term infants, especially presenting with lower *Bifidobacterium* spp. abundance and higher *Escherichia coli*, *Enterococcus* sp., and *Klebsiella pneumoniae*(Schwartz et al., 2003; Bezirtzoglou et al., 2011). As as result, perturbation of post-partum microbiota haboring contributes to the vulnerablilty in preterm-associated health consequencese, such as necrotizing enterocolitis and late-onset sepsis.

Necrotizing enterocolitis, characterized by rapid progression, high morbidity and mortality, is one of the most devastating gastrointestinal neonatal emergencies, especially in preterm newborns; the etiologies of which remains elusive. Previous studies have suggested how intestinal microbiota pattern is implicated in the condition. Mai et al. reported an increase in the Proteobacteria and a decrease in the Firmicutes phyla during three to seven days prior to NEC onset (Mai et al., 2011). Zhou et al. reported a relatively higher abundance of Clostridium and Gamma-Proteobacteria in the proximity of NEC during early and late onset, respectively(Zhou et al., 2015). Dominance of (?) and/or *Klebsiella pneumoniae* was found to correlate with NEC risk in preterm infants

Among non-Western population, however, microbiota chronological dysbiosis preceding necrotizing enterocolitis or late onset sepsis remain scant so far. Hence, we conducted this prospective study with the aims to profile and compare postpost-partum pattern of intestinal microbiota in Chinese preterm infants who subsequently developed necrotizing enterocolitis and late onset sepsis, which may be critical in the etiopathogenesis of disease.

METHODS

Ethics

This study was approved by the joint committee of ethics of Shanghai Children's Medical Center, School of Medicine Shanghai Jiao Tong University (SCMCIRB-K2013022). Detailed written informed consent was obtained from the parents prior to fecal sample collection.

Patients

Newly born preterm infants with gestational age less than 33 weeks, birth weight over 950g were enrolled from Neonatal Intensive Care Unit at Shanghai Children's Medical Center from July 2013 to December 2014. The exclusion criteria were 1) diagnosed with early-onset sepsis, 2) hepatic diseases, 3) renal impairment ($\text{Cr} > 88 \mu\text{M}$), 4) diagnosed with intestinal obstruction, 5) in foreseeable need of cardiovascular or abdominal surgeries (except for male circumcision or PDA ligation), 6) estimated parenteral support to supply over 50% of daily caloric intake for more than four days, 7) given intravenous antibiotics administration (except prophylactic regimen of cefotaxime, piperacillin-tazobactam and/or metronidazole), 8) history of oral antibiotics administration, 9) grossly bloody stools at admission, and 10) over five days old.

NEC cases were defined as infants who met the criteria for Stage II and Stage III NEC diagnosis (Bell et al., 1978), including radiographic intestinal dilation, ileus, pneumatosis intestinalis, and/or absent bowel sounds with or without abdominal tenderness, and/or mild metabolic acidosis and thrombocytopenia. LOS cases was diagnosed if 1) an infant had a positive hemoculture or other suspicious loci of infection after 72 hours of life, with septic signs/symptoms reviewed independently by at least two neonatologists, and had been treated with advanced antibiotics (e.g., Meropenem) after diagnosis. Infants with no infectious complications or sepsis were regarded as controls.

Sample collection and handling

Fecal samples collection began from neonatal meconium till discharge. Although we intended to collect fecal samples on a daily basis, due to working shifts and flexible clinical scheduling, we set seven days as the maximum interval between two collections from every infant. Every sample was collected within 30 minutes of defecation from infants' diaper with a sterile spatula. The samples were immediately placed in a cryogenic vial on dry ice and stored at -80°C within 30 minutes without additives. All samples were collected and stored before knowing the diagnosis of respective patients.

DNA extraction and quality control amplification and 16s rRNA gene sequencing

Microbial genomic DNA was isolated from each fecal specimen using the E.Z.N.A.® Soil DNA Kit (Omega Bio-Tek, Norcross, GA, U.S.) according to manufacturer's protocols. The concentration and purity of the DNA were determined by NanoDrop 2000 UV-vis spectrophotometer (Thermo Scientific, Wilmington, USA), and the DNA quality was checked by 1% agarose gel electrophoresis.

Broad-range PCR and High-throughput Sequencing of 16s rRNA gene amplicons

The V3-V4 hypervariable regions of the bacterial 16S rRNA gene were amplified from each sample using bacterial/archaeal primers 338F (5'-ACTCCTACGGGAGGCAGCAG-3') and 806R (5'-GGACTACHVGG GTWTCTAAT-3') using thermocycler PCR system (GeneAmp 9700, ABI, USA). The PCR reactions were as follows: 3 min of denaturation at 95°C , 27 cycles of 30 s at 95°C , 30 s annealing at 55°C and 45 s elongation at 72°C , and a final extension at 72°C for 10 min. The PCR reactions were performed in triplicate, with each 20 μL mixture containing 4 μL 5X FastPfu Buffer, 2 μL 2.5 mM dNTPs, 0.8 μL of each primer (5 μM), 0.4 μL FastPfu Polymerase and 10 ng template DNA. The PCR products were extracted from a 2% agarose gel and further purified using the AxyPrep DNA Gel Extraction Kit (Axygen Biosciences, Union City, CA, USA), and quantified using QuantiFluor™-ST (Promega, USA) according to the manufacturer's protocols.

Equimolar amounts of purified amplicons were pooled and paired-end sequenced (2 x 300) on an Illumina MiSeq platform (Illumina, San Diego, USA) according to the standard protocols of Majorbio Bio-Pharm Technology Co. Ltd. (Shanghai, China). The reads were de-multiplexed using the Illumina software and separate FASTQ files were generated for each specimen and deposited to the Sequence Read Archive NCBI under the BioProject accession PRJNA470548. Another public archive repository is available at figshare doi: 10.6084/m9.figshare.7205102

Raw Data Processing

After pyrosequencing, de-multiplexed sequence reads were subjected to quality filtering utilizing Trimmomatic software (version 0.36) (Bolger et al., 2014), and were truncated at any site with an Phred score <20 over a 50bp-sized window; barcode matching with the primer mismatch from 0 to 2 nucleotides was adopted and reads containing ambiguous characters were removed. After trimming, FLASH (Fast Length Adjustment of Short Read) (Magoč and Salzberg, 2011), a read pre-processing software, assembled and merged the paired-end reads from fragments and generated >10 bp overlapped, with the dead match ratio 0.2. Unassembled reads were discarded.

To fairly compare all the samples at the same sequencing depth, the "sub.sample" command of mothur program (version 1.30.1) (Schloss et al., 2009) was used for normalization to the smallest sample size. UCHIME Algorithm detected chimeric sequences, removed chimera to obtain effective reads, which were then sorted by cluster size and processed using Operational Taxonomic Units (OTUs) with 97% similarity cutoff UPARSE-OTU algorithm (implementing "cluster_otus" command) (Edgar, 2013) in USEARCH (v10) (UPARSE version 7.1). The taxonomy of each 16S rRNA gene sequence was analyzed by RDP Classifier algorithm (Wang et al., 2007) against the Silva (SSU128) (Quast et al., 2012) 16S rRNA database using confidence threshold of 70%. Each sequence was assigned the taxonomy by QIIME (Caporaso et al., 2010). The representative sequences were allocated phylogenetically down to the domain, phylum, class, order, family, and genus levels. The relative abundance of a given taxonomic group was calculated as a percentage of the sequences number belonging to that group divided by the total number of obtained sequences.

Within-sample diversity (alpha diversity) analysis, including Shannon index and Observed species richness (sobs), were obtained using the "summary.single" command of mothur program (version 1.30.1) (Schloss et al., 2009). Between-sample diversity (beta diversity) analysis was obtained estimating weighted UniFrac distances between samples.

Statistical and Bioinformatics Analysis

Demographics and Clinical Sample comparisons

Kruskal-Wallis test and Wilcoxon rank-sum test were used to identify statistically significant differences in continuous variables, including gestational age, birth weight, age when the patients were diagnosed and length of hospitalisation. The χ^2 , or Fisher's exact test were used to identify differences in gender composition. α level was considered 0.05 for all statistical tests. All statistical test not involving microbiome 16S rRNA sequencing data was performed using "stats" package using R (v.3.5.1).

Microbiota and Bioinformatics Analyses

Disease-related Time Interval Definition Under the circumstance that the sampling and disease onset timepoints for each patient were not perfectly universal, to illustrate the continuous longitudinal and repeated nature of the sampling and its relationship with onset and progression of diseases, we splitted the whole sampling span into 7 time intervals:

1. early post-partum (EPP): within 3 days afterbirth
2. early pre-onset (EPO): from the end of EPP to at least four days before disease onset
3. late pre-onset (LPO): from the end of EPO to the start of onset; for control group patients, the onset time is set at 16 days of life, as is the average diagnosis age of NEC and LOS groups.
4. early disease (ED): first third interval of whole disease span
5. middle disease (MD): second third interval of whole disease span
6. late disease (LD): last third interval of whole disease span
7. post disease (PD): from the end of disease to discharge timepoint

Diversity Analyses The average of α diversity, if more than two were available within one analysis interval, of each patient was calculated. Repeated measures two-way ANOVA, with time intervals (EPP, EPO, LPO, ED, MD, LD, PD) as a within-subject factor and groups (NEC, LOS, control) as a between-subject factor, and Kruskal Wallis tests were used to test shannon diversity differences either among groups within a time interval or among time intervals within a certain group.

Modeling Strategies for Taxonomy Comparisons To compare the dynamics of microbiota diversity and relative taxonomic abundance preceding the disease, we took into account the EPP, EPO, LPO and ED interval among all patients and fit (Supplementary matrix 1). To compare the microbiome profile right

after birth until disease alleviation, we selected EPP, EPO, LPO, ED, MD and LD interval of NEC and LOS patients(Supplementary matrix2). The average taxonomy relative abundances, if more than two were available within one analysis interval, of each patient was calculated. Zero-Inflated Beta Regression Model with Random Effects (ZIBR) and Linear Mixed-effects Model(LME) were used to test the association between OTU relative abundance and clinical covariates (diseases-related time intervals) for longitudinal microbiome data (Chen and Li, 2016). *ZIBR* R package were utilized for both models. Figures were generated with the "*ggpubr*"(Kassambara, 2017) and "*ggplot2*"(Wickham, 2016) packages using R(v.3.5.1). Scripts for modeling and figures plotting, input and output files, figures are available at our github repository.

Scripts and Figures Archiving

Figures were generated with the "*ggpubr*"(Kassambara, 2017) and "*ggplot2*"(Wickham, 2016) packages using R(v.3.5.1)

RESULTS

Overall 16s Sequencing Report and Coverage

A total of 7,472,400 optimized V3-V4 tags of 16s rRNA gene sequences were produced from 192 fecal samples, with an average read length of 448 bp (Table S1).

Patients characteristics

Totally 130 infants met the criteria of our study, and 1698 samples were collected from them in the neonatal intensive care unit (NICU) at Shanghai Children's Medical Center from July 2013 to December 2014. Among whom, we studied 192 fecal samples from 24 well-sampled preterm infants, including four subsequently diagnosed with NEC (2 in stage IIA and 2 in stage IIB), three with LOS, and 17 matched controls (Figure1 ,Supplemantary Table S2). Fecal samples were collected between days 1 and 69 of life. Sampling timepoints and numbers of samples varied among each infant.




Figure 1. Flow of Study Design

All infants were delivered by cesarean section and fed on infant formula. No one was prescribed probiotics during hospitalization. Comparisons showed no significant difference in terms of gestational age, birth weight and gender proportions, diagnosed age among three groups (Table 1). Lenth of stay mong three groups was significantly different however rational since NEC and LOS patients usually require longer period of healthcare because their worse health status often required longer health care time than the control group. All infants were delivered by cesarean section and fed on infant formula. No one was prescribed probiotics during hospitalization.

Dynimics of Microbiome Diversity in diseases onset and progression

Mircobiome Richness Plays Minor Roles

Overallly, all three groups shared similar microbiota richness trend after birth, that the observed richness (Sobs) decreased significantly from post-partum period until the late disease stage (Fig2a. NEC group, p =

0.044; b. LOS group, $p = 0.013$; c. control group, $p < 0.01$). Nevertheless, both the inter-group test between adjacent time-intervals (Fig2 and the inter-time-interval comparison among groups (supplementary sobs-time-group!!!!) didn't show any significant alterations and differences in sobs, indicating the minor effect of microbiota richness on the diseases acquisition.

Mircobiome Eveness

Microbiome Composition

Taxonomic Determinants

DISCUSSION

are consistent with the hypothesis that dysbiosis precedes this severe event. The Our study has its limitations. We acknowledge that the sample size is limited since this study is single-center-based and the incidence of both diseases are relatively low: among the 1148 preterm infants admitted within July 2013 to December 2014, only five developed NEC. Our results, however, showed the needs of larger study population and longer follow ups. Furthermore, the resultant overfitting possibility inevitably rose up, which became the pitfall in understanding the true microbiota patterns preceding NEC and LOS.

CONCLUSIONS

ACKNOWLEDGMENTS

We appreciate the support from enrolled patients, their families, and all staffs at Shanghai Children's Medical Center.

SOME L^AT_EX EXAMPLES

Use section and subsection commands to organize your document. L^AT_EX handles all the formatting and numbering automatically. Use ref and label commands for cross-references.

Figures and Tables

Use the table and tabular commands for basic tables — see Table 2, for example. You can upload a figure (JPEG, PNG or PDF) using the project menu. To include it in your document, use the includegraphics command as in the code for Figure 3 below.

Table 1. Demographic characteristics of Preterm NEC, LOS and control groups.

	NEC (N=3)	LOS (N=4)	Control (N=17)	Statistical Test	<i>p value</i>
Gestational Age (weeks)	29(29-30)	30(29-31)	31(28-33)	Kruskal-Wallis test	0.074
Birth Weight(g)	1416.3 (773.4-2149.1)	1141.7 (633.4-1649.9)	1527.4 (1391.6-1663.1)	Kruskal-Wallis test	0.111
Gender				Fisher's exact test	0.82
Female	3(75%)	2(67%)	9(53%)		
male	1(25%)	1(33%)	8(47%)		
Diagnosis Age(days)	16(11-19)	16(10-22)	—	Wilcoxon rank-sum test	0.629
Length of Stay(d)	54.3 (13.5-95.0)	60.0 (24.8-95.2)	32.9 (26.3-39.5)	Kruskal-Wallis test	0.046
Number of Samples	46	42	103	—	—




Figure 2. Post-partum microbiome richness trend (Sobs)

Citations

LaTeX formats citations and references automatically using the bibliography records in your .bib file, which you can edit via the project menu. Use the cite command for an inline citation, like ?, and the citep command for a citation in parentheses (?).

Mathematics

LaTeX is great at typesetting mathematics. Let X_1, X_2, \dots, X_n be a sequence of independent and identically distributed random variables with $E[X_i] = \mu$ and $\text{Var}[X_i] = \sigma^2 < \infty$, and let

$$S_n = \frac{X_1 + X_2 + \dots + X_n}{n} = \frac{1}{n} \sum_{i=1}^n X_i$$

denote their mean. Then as n approaches infinity, the random variables $\sqrt{n}(S_n - \mu)$ converge in distribution to a normal $\mathcal{N}(0, \sigma^2)$.

Lists

You can make lists with automatic numbering ...

1. Like this,




Figure 3. An example image.

Item	Quantity
Widgets	42
Gadgets	13

Table 2. An example table.

254 2. and like this.

255 ... or bullet points ...

- 256 • Like this,
- 257 • and like this.

258 ... or with words and descriptions ...

259 **Word** Definition

260 **Concept** Explanation

261 **Idea** Text

262 We hope you find write \LaTeX useful for your PeerJ submission, and please let us know if you have any
 263 feedback. Further examples with dummy text are included in the following pages.

264 METHODS

265 Quisque ullamcorper placerat ipsum. Cras nibh. Morbi vel justo vitae lacus tincidunt ultrices. Lorem
 266 ipsum dolor sit amet, consectetur adipiscing elit. In hac habitasse platea dictumst. Integer tempus
 267 convallis augue. Etiam facilisis. Nunc elementum fermentum wisi. Aenean placerat. Ut imperdiet, enim
 268 sed gravida sollicitudin, felis odio placerat quam, ac pulvinar elit purus eget enim. Nunc vitae tortor.
 269 Proin tempus nibh sit amet nisl. Vivamus quis tortor vitae risus porta vehicula.

$$\cos^3 \theta = \frac{1}{4} \cos \theta + \frac{3}{4} \cos 3\theta \quad (1)$$

270 Fusce mauris. Vestibulum luctus nibh at lectus. Sed bibendum, nulla a faucibus semper, leo velit
 271 ultricies tellus, ac venenatis arcu wisi vel nisl. Vestibulum diam. Aliquam pellentesque, augue quis sagittis

272 posuere, turpis lacus congue quam, in hendrerit risus eros eget felis. Maecenas eget erat in sapien mattis
273 porttitor. Vestibulum porttitor. Nulla facilisi. Sed a turpis eu lacus commodo facilisis. Morbi fringilla,
274 wisi in dignissim interdum, justo lectus sagittis dui, et vehicula libero dui cursus dui. Mauris tempor
275 ligula sed lacus. Duis cursus enim ut augue. Cras ac magna. Cras nulla. Nulla egestas. Curabitur a leo.
276 Quisque egestas wisi eget nunc. Nam feugiat lacus vel est. Curabitur consectetur.

277 Subsection

278 Suspendisse vel felis. Ut lorem lorem, interdum eu, tincidunt sit amet, laoreet vitae, arcu. Aenean faucibus
279 pede eu ante. Praesent enim elit, rutrum at, molestie non, nonummy vel, nisl. Ut lectus eros, malesuada
280 sit amet, fermentum eu, sodales cursus, magna. Donec eu purus. Quisque vehicula, urna sed ultricies
281 auctor, pede lorem egestas dui, et convallis elit erat sed nulla. Donec luctus. Curabitur et nunc. Aliquam
282 dolor odio, commodo pretium, ultricies non, pharetra in, velit. Integer arcu est, nonummy in, fermentum
283 faucibus, egestas vel, odio.

284 **Paragraph** Sed commodo posuere pede. Mauris ut est. Ut quis purus. Sed ac odio. Sed vehicula
285 hendrerit sem. Duis non odio. Morbi ut dui. Sed accumsan risus eget odio. In hac habitasse platea
286 dictumst. Pellentesque non elit. Fusce sed justo eu urna porta tincidunt. Mauris felis odio, sollicitudin sed,
287 volutpat a, ornare ac, erat. Morbi quis dolor. Donec pellentesque, erat ac sagittis semper, nunc dui lobortis
288 purus, quis congue purus metus ultricies tellus. Proin et quam. Class aptent taciti sociosqu ad litora
289 torquent per conubia nostra, per inceptos hymenaeos. Praesent sapien turpis, fermentum vel, eleifend
290 faucibus, vehicula eu, lacus.

291 **Paragraph** Pellentesque habitant morbi tristique senectus et netus et malesuada fames ac turpis egestas.
292 Donec odio elit, dictum in, hendrerit sit amet, egestas sed, leo. Praesent feugiat sapien aliquet odio.
293 Integer vitae justo. Aliquam vestibulum fringilla lorem. Sed neque lectus, consectetur at, consectetur
294 sed, eleifend ac, lectus. Nulla facilisi. Pellentesque eget lectus. Proin eu metus. Sed porttitor. In hac
295 habitasse platea dictumst. Suspendisse eu lectus. Ut mi mi, lacinia sit amet, placerat et, mollis vitae, dui.
296 Sed ante tellus, tristique ut, iaculis eu, malesuada ac, dui. Mauris nibh leo, facilisis non, adipiscing quis,
297 ultrices a, dui.

298 Subsection

299 Morbi luctus, wisi viverra faucibus pretium, nibh est placerat odio, nec commodo wisi enim eget quam.
300 Quisque libero justo, consectetur a, feugiat vitae, porttitor eu, libero. Suspendisse sed mauris vitae
301 elit sollicitudin malesuada. Maecenas ultricies eros sit amet ante. Ut venenatis velit. Maecenas sed mi
302 eget dui varius euismod. Phasellus aliquet volutpat odio. Vestibulum ante ipsum primis in faucibus orci
303 luctus et ultrices posuere cubilia Curae; Pellentesque sit amet pede ac sem eleifend consectetur. Nullam
304 elementum, urna vel imperdiet sodales, elit ipsum pharetra ligula, ac pretium ante justo a nulla. Curabitur
305 tristique arcu eu metus. Vestibulum lectus. Proin mauris. Proin eu nunc eu urna hendrerit faucibus.
306 Aliquam auctor, pede consequat laoreet varius, eros tellus scelerisque quam, pellentesque hendrerit ipsum
307 dolor sed augue. Nulla nec lacus.

308 Reference to Figure 4.

309 RESULTS AND DISCUSSION

310 Suspendisse vitae elit. Aliquam arcu neque, ornare in, ullamcorper quis, commodo eu, libero. Fusce
311 sagittis erat at erat tristique mollis. Maecenas sapien libero, molestie et, lobortis in, sodales eget, dui.
312 Morbi ultrices rutrum lorem. Nam elementum ullamcorper leo. Morbi dui. Aliquam sagittis. Nunc
313 placerat. Pellentesque tristique sodales est. Maecenas imperdiet lacinia velit. Cras non urna. Morbi eros
314 pede, suscipit ac, varius vel, egestas non, eros. Praesent malesuada, diam id pretium elementum, eros sem
315 dictum tortor, vel consectetur odio sem sed wisi.

316 Subsection

317 Sed feugiat. Cum sociis natoque penatibus et magnis dis parturient montes, nascetur ridiculus mus. Ut
318 pellentesque augue sed urna. Vestibulum diam eros, fringilla et, consectetur eu, nonummy id, sapien.
319 Nullam at lectus. In sagittis ultrices mauris. Curabitur malesuada erat sit amet massa. Fusce blandit.
320 Aliquam erat volutpat. Aliquam euismod. Aenean vel lectus. Nunc imperdiet justo nec dolor.




Figure 4. In-text Picture

321 **Subsubsection**

322 Etiam euismod. Fusce facilisis lacinia dui. Suspendisse potenti. In mi erat, cursus id, nonummy sed,
 323 ullamcorper eget, sapien. Praesent pretium, magna in eleifend egestas, pede pede pretium lorem, quis
 324 consectetur tortor sapien facilisis magna. Mauris quis magna varius nulla scelerisque imperdiet. Aliquam
 325 non quam. Aliquam porttitor quam a lacus. Praesent vel arcu ut tortor cursus volutpat. In vitae pede quis
 326 diam bibendum placerat. Fusce elementum convallis neque. Sed dolor orci, scelerisque ac, dapibus nec,
 327 ultricies ut, mi. Duis nec dui quis leo sagittis commodo.

328 **Subsubsection**

329 Etiam ac leo a risus tristique nonummy. Donec dignissim tincidunt nulla. Vestibulum rhoncus molestie
 330 odio. Sed lobortis, justo et pretium lobortis, mauris turpis condimentum augue, nec ultricies nibh arcu
 331 pretium enim. Nunc purus neque, placerat id, imperdiet sed, pellentesque nec, nisl. Vestibulum imperdiet
 332 neque non sem accumsan laoreet. In hac habitasse platea dictumst. Etiam condimentum facilisis libero.
 333 Suspendisse in elit quis nisl aliquam dapibus. Pellentesque auctor sapien. Sed egestas sapien nec lectus.
 334 Pellentesque vel dui vel neque bibendum viverra. Aliquam porttitor nisl nec pede. Proin mattis libero vel
 335 turpis. Donec rutrum mauris et libero. Proin euismod porta felis. Nam lobortis, metus quis elementum
 336 commodo, nunc lectus elementum mauris, eget vulputate ligula tellus eu neque. Vivamus eu dolor.

337 **Subsection**

338 Nulla in ipsum. Praesent eros nulla, congue vitae, euismod ut, commodo a, wisi. Pellentesque habitant
 339 morbi tristique senectus et netus et malesuada fames ac turpis egestas. Aenean nonummy magna non leo.
 340 Sed felis erat, ullamcorper in, dictum non, ultricies ut, lectus. Proin vel arcu a odio lobortis euismod.
 341 Vestibulum ante ipsum primis in faucibus orci luctus et ultrices posuere cubilia Curae; Proin ut est.
 342 Aliquam odio. Pellentesque massa turpis, cursus eu, euismod nec, tempor congue, nulla. Duis viverra
 343 gravida mauris. Cras tincidunt. Curabitur eros ligula, varius ut, pulvinar in, cursus faucibus, augue.

344 Nulla mattis luctus nulla. Duis commodo velit at leo. Aliquam vulputate magna et leo. Nam

345 vestibulum ullamcorper leo. Vestibulum condimentum rutrum mauris. Donec id mauris. Morbi molestie
346 justo et pede. Vivamus eget turpis sed nisl cursus tempor. Curabitur mollis sapien condimentum nunc. In
347 wisi nisl, malesuada at, dignissim sit amet, lobortis in, odio. Aenean consequat arcu a ante. Pellentesque
348 porta elit sit amet orci. Etiam at turpis nec elit ultricies imperdiet. Nulla facilisi. In hac habitasse platea
349 dictumst. Suspendisse viverra aliquam risus. Nullam pede justo, molestie nonummy, scelerisque eu,
350 facilisis vel, arcu.

351 Curabitur tellus magna, porttitor a, commodo a, commodo in, tortor. Donec interdum. Praesent
352 scelerisque. Maecenas posuere sodales odio. Vivamus metus lacus, varius quis, imperdiet quis, rhoncus a,
353 turpis. Etiam ligula arcu, elementum a, venenatis quis, sollicitudin sed, metus. Donec nunc pede, tincidunt
354 in, venenatis vitae, faucibus vel, nibh. Pellentesque wisi. Nullam malesuada. Morbi ut tellus ut pede
355 tincidunt porta. Lorem ipsum dolor sit amet, consectetur adipiscing elit. Etiam congue neque id dolor.

356 Donec et nisl at wisi luctus bibendum. Nam interdum tellus ac libero. Sed sem justo, laoreet vitae,
357 fringilla at, adipiscing ut, nibh. Maecenas non sem quis tortor eleifend fermentum. Etiam id tortor ac
358 mauris porta vulputate. Integer porta neque vitae massa. Maecenas tempus libero a libero posuere dictum.
359 Vestibulum ante ipsum primis in faucibus orci luctus et ultrices posuere cubilia Curae; Aenean quis mauris
360 sed elit commodo placerat. Class aptent taciti sociosqu ad litora torquent per conubia nostra, per inceptos
361 hymenaeos. Vivamus rhoncus tincidunt libero. Etiam elementum pretium justo. Vivamus est. Morbi a
362 tellus eget pede tristique commodo. Nulla nisl. Vestibulum sed nisl eu sapien cursus rutrum.

363 Nulla non mauris vitae wisi posuere convallis. Sed eu nulla nec eros scelerisque pharetra. Nullam
364 varius. Etiam dignissim elementum metus. Vestibulum faucibus, metus sit amet mattis rhoncus, sapien
365 dui laoreet odio, nec ultricies nibh augue a enim. Fusce in ligula. Quisque at magna et nulla commodo
366 consequat. Proin accumsan imperdiet sem. Nunc porta. Donec feugiat mi at justo. Phasellus facilisis
367 ipsum quis ante. In ac elit eget ipsum pharetra faucibus. Maecenas viverra nulla in massa.

368 Nulla ac nisl. Nullam urna nulla, ullamcorper in, interdum sit amet, gravida ut, risus. Aenean ac
369 enim. In luctus. Phasellus eu quam vitae turpis viverra pellentesque. Duis feugiat felis ut enim. Phasellus
370 pharetra, sem id porttitor sodales, magna nunc aliquet nibh, nec blandit nisl mauris at pede. Suspendisse
371 risus risus, lobortis eget, semper at, imperdiet sit amet, quam. Quisque scelerisque dapibus nibh. Nam
372 enim. Lorem ipsum dolor sit amet, consectetur adipiscing elit. Nunc ut metus. Ut metus justo, auctor at,
373 ultrices eu, sagittis ut, purus. Aliquam aliquam.

374 ACKNOWLEDGMENTS

375 So long and thanks for all the fish.

376 REFERENCES

- 377 Bell, M. J., Ternberg, J. L., Feigin, R. D., Keating, J. P., Marshall, R., Barton, L., and Brotherton, T.
378 (1978). Neonatal necrotizing enterocolitis. therapeutic decisions based upon clinical staging. *Annals of*
379 *surgery*, 187(1):1.
- 380 Bezirtzoglou, E., Tsiotsias, A., and Welling, G. W. (2011). Microbiota profile in feces of breast-and
381 formula-fed newborns by using fluorescence in situ hybridization (fish). *Anaerobe*, 17(6):478–482.
- 382 Bolger, A. M., Lohse, M., and Usadel, B. (2014). Trimmomatic: a flexible trimmer for illumina sequence
383 data. *Bioinformatics*, 30(15):2114–2120.
- 384 Bouter, K. E., van Raalte, D. H., Groen, A. K., and Nieuwdorp, M. (2017). Role of the gut microbiome in
385 the pathogenesis of obesity and obesity-related metabolic dysfunction. *Gastroenterology*, 152(7):1671–
386 1678.
- 387 Caporaso, J. G., Kuczynski, J., Stombaugh, J., Bittinger, K., Bushman, F. D., Costello, E. K., Fierer, N.,
388 Pena, A. G., Goodrich, J. K., Gordon, J. I., et al. (2010). Qiime allows analysis of high-throughput
389 community sequencing data. *Nature methods*, 7(5):335.
- 390 Chen, E. Z. and Li, H. (2016). A two-part mixed-effects model for analyzing longitudinal microbiome
391 compositional data. *Bioinformatics*, 32(17):2611–2617.
- 392 Edgar, R. C. (2013). Uparse: highly accurate otu sequences from microbial amplicon reads. *Nature*
393 *methods*, 10(10):996.
- 394 Giongo, A., Gano, K. A., Crabb, D. B., Mukherjee, N., Novelo, L. L., Casella, G., Drew, J. C., Ilonen, J.,
395 Knip, M., Hyöty, H., et al. (2011). Toward defining the autoimmune microbiome for type 1 diabetes.
396 *The ISME journal*, 5(1):82.

397 Kassambara, A. (2017). ggpubr:“ggplot2” based publication ready plots. r package version 0.1. 6.

398 Madan, J. C., Farzan, S. F., Hibberd, P. L., and Karagas, M. R. (2012). Normal neonatal microbiome
399 variation in relation to environmental factors, infection and allergy. *Current opinion in pediatrics*,
400 24(6):753.

401 Magoč, T. and Salzberg, S. L. (2011). Flash: fast length adjustment of short reads to improve genome
402 assemblies. *Bioinformatics*, 27(21):2957–2963.

403 Quast, C., Pruesse, E., Yilmaz, P., Gerken, J., Schweer, T., Yarza, P., Peplies, J., and Glöckner, F. O.
404 (2012). The silva ribosomal rna gene database project: improved data processing and web-based tools.
405 *Nucleic acids research*, 41(D1):D590–D596.

406 Sarkar, A., Lehto, S. M., Harty, S., Dinan, T. G., Cryan, J. F., and Burnet, P. W. (2016). Psychobiotics and
407 the manipulation of bacteria–gut–brain signals. *Trends in neurosciences*, 39(11):763–781.

408 Savage, J. H., Lee-Sarwar, K. A., Sordillo, J., Bunyavanich, S., Zhou, Y., O’connor, G., Sandel, M.,
409 Bacharier, L. B., Zeiger, R., Sodergren, E., et al. (2018). A prospective microbiome-wide association
410 study of food sensitization and food allergy in early childhood. *Allergy*, 73(1):145–152.

411 Schloss, P. D., Westcott, S. L., Ryabin, T., Hall, J. R., Hartmann, M., Hollister, E. B., Lesniewski, R. A.,
412 Oakley, B. B., Parks, D. H., Robinson, C. J., et al. (2009). Introducing mothur: open-source, platform-
413 independent, community-supported software for describing and comparing microbial communities.
414 *Applied and environmental microbiology*, 75(23):7537–7541.

415 Schwirtz, A., Gruhl, B., Löbnitz, M., Michel, P., Radke, M., and Blaut, M. (2003). Development of the
416 intestinal bacterial composition in hospitalized preterm infants in comparison with breast-fed, full-term
417 infants. *Pediatric research*, 54(3):393.

418 Stokholm, J., Blaser, M. J., Thorsen, J., Rasmussen, M. A., Waage, J., Vinding, R. K., Schoos, A.-M. M.,
419 Kunø, A., Fink, N. R., Chawes, B. L., et al. (2018). Maturation of the gut microbiome and risk of
420 asthma in childhood. *Nature communications*, 9(1):141.

421 Tang, W. W., Kitai, T., and Hazen, S. L. (2017). Gut microbiota in cardiovascular health and disease.
422 *Circulation research*, 120(7):1183–1196.

423 Tilg, H., Cani, P. D., and Mayer, E. A. (2016). Gut microbiome and liver diseases. *Gut*, 65(12):2035–2044.

424 Vatanen, T., Franzosa, E. A., Schwager, R., Tripathi, S., Arthur, T. D., Vehik, K., Lernmark, Å., Hagopian,
425 W. A., Rewers, M. J., She, J.-X., et al. (2018). The human gut microbiome in early-onset type 1 diabetes
426 from the teddy study. *Nature*, 562(7728):589.

427 Wang, Q., Garrity, G. M., Tiedje, J. M., and Cole, J. R. (2007). Naive bayesian classifier for rapid
428 assignment of rna sequences into the new bacterial taxonomy. *Applied and environmental microbiology*,
429 73(16):5261–5267.

430 Wickham, H. (2016). *ggplot2: Elegant Graphics for Data Analysis*. Springer-Verlag New York.

431 Zhou, Y., Shan, G., Sodergren, E., Weinstock, G., Walker, W. A., and Gregory, K. E. (2015). Longitudinal
432 analysis of the premature infant intestinal microbiome prior to necrotizing enterocolitis: a case-control
433 study. *PloS one*, 10(3):e0118632.