
MA685 HW1#9

Jiayuan Shi 01/25/2016

library(ISLR)

9a.

pairs(Auto)



9b.

cor(subset(Auto, select=-name))

```
##
                     mpg cylinders displacement horsepower
                                                            weight
## mpg
               1.0000000 -0.7776175
                                    -0.8051269 -0.7784268 -0.8322442
## cylinders
              -0.7776175 1.0000000
                                     ## displacement -0.8051269 0.9508233
                                     1.0000000 0.8972570
                                                         0.9329944
## horsepower
              -0.7784268 0.8429834
                                     0.8972570 1.0000000 0.8645377
## weight
              -0.8322442 0.8975273
                                     0.9329944 0.8645377 1.0000000
## acceleration 0.4233285 -0.5046834
                                    -0.5438005 -0.6891955 -0.4168392
```

```
## year
                 0.5805410 -0.3456474
                                        -0.3698552 -0.4163615 -0.3091199
## origin
                 0.5652088 -0.5689316
                                        -0.6145351 -0.4551715 -0.5850054
##
                acceleration
                                   year
                                            origin
                   0.4233285 0.5805410 0.5652088
## mpg
## cylinders
                  -0.5046834 -0.3456474 -0.5689316
## displacement
                 -0.5438005 -0.3698552 -0.6145351
## horsepower
                  -0.6891955 -0.4163615 -0.4551715
## weight
                  -0.4168392 -0.3091199 -0.5850054
## acceleration
                  1.0000000 0.2903161 0.2127458
## year
                  0.2903161
                             1.0000000
                                        0.1815277
## origin
                  0.2127458 0.1815277 1.0000000
```


9c.


```
lm1 = lm(mpg~.-name, data=Auto)
summary(lm1)
```

```
##
## Call:
## lm(formula = mpg ~ . - name, data = Auto)
##
## Residuals:
                10 Median
                                3Q
                                       Max
  -9.5903 -2.1565 -0.1169
                           1.8690 13.0604
##
##
## Coefficients:
##
                  Estimate Std. Error t value Pr(>|t|)
## (Intercept) -17.218435
                            4.644294
                                       -3.707 0.00024 ***
## cylinders
                 -0.493376
                            0.323282
                                      -1.526 0.12780
## displacement
                 0.019896
                            0.007515
                                        2.647
                                               0.00844 **
                 -0.016951
                            0.013787
                                       -1.230
                                              0.21963
## horsepower
## weight
                 -0.006474
                            0.000652
                                       -9.929
                                              < 2e-16
                                       0.815 0.41548
## acceleration
                 0.080576
                            0.098845
## year
                  0.750773
                            0.050973
                                       14.729
                                              < 2e-16 ***
                                       5.127 4.67e-07 ***
## origin
                  1.426141
                            0.278136
## ---
## Signif. codes: 0 '***' 0.001 '**' 0.05 '.' 0.1 ' ' 1
## Residual standard error: 3.328 on 384 degrees of freedom
## Multiple R-squared: 0.8215, Adjusted R-squared: 0.8182
## F-statistic: 252.4 on 7 and 384 DF, p-value: < 2.2e-16
```

- i. Yes, there is a relatioship between the predictors and the response by testing the null hypothesis of whether all the regression coefficients are zero. The F -statistic is has a small p-value, less than 0.01, indicating that we can reject the null hypothesis. Therefore, there is a significant relatioship between the predictors and the response.
- ii. From the p-values of each predictor, displacement, weight, year, and origin have a statistically significant relationship to the response (since they have small p-values < 0.01), while cylinders, horsepower, and acceleration do not.
- iii. The coefficient for year, 0.7508, suggests that for every one year increase, mpg also increases by 0.7508.


9d.





There are three problems with the fit. One is that the curve pattern in the residuals plot indicates non-linearity in the data. The other is from the leverage plot, point 14 appears to have high leverage, although not a high magnitude residual.

plot(predict(lm1), rstudent(lm1))



Also there may be some outliers in the plot of studentized residuals because there exist some values greater than 3.

9e.

```
lm2 = lm(mpg~cylinders*displacement+displacement*weight, data=Auto)
summary(lm2)
```

```
##
## Call:
  lm(formula = mpg ~ cylinders * displacement + displacement *
##
##
       weight, data = Auto)
##
## Residuals:
##
        Min
                  1Q
                       Median
                                     3Q
                                             Max
  -13.2934
            -2.5184
                      -0.3476
                                1.8399
                                        17.7723
##
##
  Coefficients:
##
                            Estimate Std. Error t value Pr(>|t|)
## (Intercept)
                           5.262e+01
                                      2.237e+00
                                                  23.519
                                                          < 2e-16 ***
## cylinders
                           7.606e-01
                                                   0.992
                                                            0.322
                                      7.669e-01
## displacement
                          -7.351e-02
                                      1.669e-02
                                                  -4.403 1.38e-05 ***
## weight
                          -9.888e-03
                                      1.329e-03
                                                  -7.438 6.69e-13 ***
## cylinders:displacement -2.986e-03
                                      3.426e-03
                                                  -0.872
                                                            0.384
## displacement:weight
                                                   4.254 2.64e-05 ***
                           2.128e-05
                                      5.002e-06
##
## Signif. codes: 0 '***' 0.001 '**' 0.05 '.' 0.1 ' ' 1
## Residual standard error: 4.103 on 386 degrees of freedom
```

```
## Multiple R-squared: 0.7272, Adjusted R-squared: 0.7237
## F-statistic: 205.8 on 5 and 386 DF, p-value: < 2.2e-16</pre>
```


From the correlation matrix, I got the two highest correlated pairs (>0.9) and used them in picking my interaction effects. From the p-values, we can see that the interaction between displacement and weight is statistically significant with p-value <0.01, while the interaction between cylinders and displacement is not.


9f.

par(mfrow=c(2,2))

plot(lm3)

```
lm3 = lm(log(mpg)~.-name, data=Auto)
summary(lm3)
##
## Call:
## lm(formula = log(mpg) ~ . - name, data = Auto)
##
## Residuals:
       Min
##
                 1Q
                      Median
                                   3Q
                                           Max
## -0.40955 -0.06533 0.00079 0.06785 0.33925
##
## Coefficients:
##
                 Estimate Std. Error t value Pr(>|t|)
## (Intercept)
                1.751e+00 1.662e-01 10.533 < 2e-16 ***
## cylinders
               -2.795e-02 1.157e-02 -2.415 0.01619 *
## displacement 6.362e-04 2.690e-04
                                      2.365 0.01852 *
## horsepower
               -1.475e-03 4.935e-04 -2.989 0.00298 **
## weight
               -2.551e-04 2.334e-05 -10.931 < 2e-16 ***
## acceleration -1.348e-03 3.538e-03 -0.381 0.70339
## year
                2.958e-02 1.824e-03 16.211 < 2e-16 ***
                4.071e-02 9.955e-03
                                     4.089 5.28e-05 ***
## origin
## ---
## Signif. codes: 0 '***' 0.001 '**' 0.05 '.' 0.1 ' ' 1
##
## Residual standard error: 0.1191 on 384 degrees of freedom
## Multiple R-squared: 0.8795, Adjusted R-squared: 0.8773
## F-statistic: 400.4 on 7 and 384 DF, p-value: < 2.2e-16
```





predict(lm3)

From the correlation matrix in 9a., displacement, horsepower and weight show a similar nonlinear pattern against the response mpg. This nonlinear pattern is very close to a log form. So we use log(mpg) as our response variable.

The outputs show that log transform of mpg yield better model fitting. For example, R^2 increases from 0.8215 to 0.8795. Residuals tend to be more normal. The curve pattern in the residuals plot indicates more linearity in the data. Also, there are less outliers in the plot of studentized residuals because all values are less than 3.