

ACCESS-Fire report

Jesse Greenslade

Mika Peace

Harvey Ye

Bureau of Meteorology

Jeff Kepert

Kevin Tory

May 4, 2020

Contents

1	Introduction	2
2	Model	3
2.1	ACCESS	3
2.2	Coupled Fire Model	3
2.3	Model runs	3
2.4	Modelled Weather	7
2.5	Data flow	7
3	Pyrocumulonimbus	9
4	Ember storm	10

Abstract

This is the abstract ...

1 Introduction

Introduction to project and model. Bushfires are extremely dangerous for many reasons. Firefighters depend on accurate weather forecasting and information in order to mitigate risks both to themselves and the community in general. One issue is that fires can generate their own local weather such as vortexes [1] and storms (pyrocumulonimbus or PCB) [2] that generate lightning and wind gusts that have not been forecast by models.

Model output is examined to improve understanding of weather phenomena in the context of bushfires.

Outline The remainder of this article is organized as follows...

2 Model

2.1 ACCESS

TODO: Blurb about ACCESS

Model output are 4-dimensional: latitude, longitude, vertical level (θ or ρ), and time. Most fields are defined at their grid-box midpoints. Wind speed and direction is defined at the direction-dependent gridbox edge.

2.2 Coupled Fire Model

TODO: Details about fire model

Model output has the same latitude and longitude as ACCESS output, with more frequent time steps and no vertical component. A fire front field is produced that is negative where the fire has burnt, and positive elsewhere.

2.3 Model runs

Over the course of the project the ACCESS-Fire model has been run and updated multiple times, over two regions where large scale forest fires occurred. The model runs in a nested fashion, with three to four nests so that the highest resolution possible is produced in a reasonable manner at the site of the fires (e.g. Figure 1).

The first fire under examination occurred a few kilometers East from Waroona, a suburb South of Perth. The second occurred a few hundred kilometres North of Sydney. Both locations are modelled at high resolution using ACCESS-Fire in a nested setup (Figure 1).

This report analyses output from the latest iteration of ACCESS-Fire, earlier iterations are listed here as they may help elucidate model parameterisation decisions.

Waroona_{oldold}

- Run in the depths of the past (Oct 2016?)
- Output at 30 minute resolution
- Meteorology not affected by fire model



Figure 1: Three nests run by ACCESS-Fire model are shown as black/red rectangles. The outermost nest has resolution of approximately 3.5 km by 3.5 km, The intermediary Nest is at 1.0 km by 1.0 km, and the smallest nest (red) is at 0.3 km by 0.3 km. A fourth nest is sometimes added with .1 km by .1 km resolution. Fire coupling is enabled only within the higher resolution (red) nests.

- slightly different grid to other model runs

Waroona_old

- Run in Aug, 2018
- Output at 30 minute resolution
- Run crashed after 21 hours due to runaway model physics (vertical wind speeds $> 1\text{km/s}$)
- Fast fire spread parameters
- Clear PCB creation around 1100

Waroona_run1

- Run in Aug, 2019
- Output at 10 minute resolution
- Increased boundary layer stability to prevent crash
- Slower updated fire spread parameters
- No real PCB creation seen

Waroona_run2

- Run in Dec, 2019 (Raijin's last encore)
- Output at 10 minute resolution
- Increased boundary layer stability
- Fire spread parameters: Faster (older, matching Waroona_old)
- can be compared to Waroona_run2uc, which has identical settings but no fire coupling

Waroona_run3

- Run in Feb, 2020 (GADI)
- Output at 10 minute resolution
- Increased boundary layer stability
- Fire spread parameters: ?




Figure 2: Model run fire fronts at start of burn and end of simulation

2.4 Modelled Weather

Weather is examined at various scales by sub-selecting extents and vertical levels, then viewing horizontal and vertical winds, cloud content, and other available metrics. This section outlines in chronological order the overall weather in the latest model run for each site.

The Waroona simulation starts 1 hour before midnight, and winds from ESE near the surface form horizontal rolls for several hours (Figure TODO). Fire ignition occurs after about 5 hours (4AM), and progresses westwards in the easterly winds. The fire is seen to influence vertical motion a long way down-wind, before clear East-West oriented horizontal rolls occur in the vertical motion around midday (Figure TODO). At about 2 in the afternoon, some westward winds occur at the base of the escarpment, causing updraughts and disrupting the horizontal rolls. Coincidentally the vertical motion near the fire front can be seen to have pushed into the upper troposphere leading to heightened cloud activity (Figure 3), This feature is discussed more in Section 3. Easterly winds re-establish dominance by 7PM, getting stronger especially down the escarpment until the end of the simulation. The fire spreads down the escarpment on this first evening exactly when these winds pick up, this feature is discussed in more detail in Section 4.

2.5 Data flow

TODO: Data flow diagram for project

waroona_run3 weather 2016 Jan 06 06:00 (UTC)




Figure 3: Left panels show horizontal wind speed (coloured contour map) and wind direction (arrows). Right panels show vertical wind speed (coloured contour map) and cloud content (diagonal stippling). The wind speeds are averaged into vertical bins based on height above the ground, showing lower to higher altitudes from the first row to the last row respectively. Cloud content is summed within each vertical bin, and the areas marked have at least 0.1 gkg^{-1} of water and ice.

3 Pyrocumulonimbus

Formation General hand wavey physics etc.

Model formation Details regarding how PCB is found/examined in model output. Story about how the parameters affect model PCB formation. Some stuff about vorticity and other metrics?

3.1 Waroona

Figure ?? shows horizontal model slices (at varying altitudes) of vertical wind motion (filled contour map), and cloud content (black outlines).

PCB Formation Threshold (PFT) Kevin's PFT - point to his publication. Calculation within model output Summary showing that when fire spreads fast the output is higher. High output coinciding with low PFT is exactly when PCB occurs.

```
.../.../figures/waroona_run3/vert_motion_slices/fig_201601060630.png
```

Figure 4: Top-down views of vertical motion on model levels of increasing altitude from left to right, top to bottom. A red contour shows the fire front, and cloud content above the 0.1 gkg threshold is marked by stippling.

4 Ember storm




Figure 5: Top panel: Top down view of topography, surface horizontal winds (quivers), and 300m above ground vertical wind contour. Vertical winds are shown with dashed lines (at 3 m/s) upwards and downwards (pink, blue respectively). Waroona is marked by a grey dot. Middle panel: Potential temperature transect along dashed black line in panel 1. An arrow points to the X-axis depicting the west-most point of the fire front, and an arrow along the Y-axis showing the altitude of the vertical wind contours shown in the top panel. Bottom panel: Vertical motion transect (coloured) with quivers showing wind speed and direction on the east-west-vertical plane.

References

- [1] FORTHOFER, J. M., AND GOODRICK, S. L. Review of vortices in wildland fire. *Journal of Combustion* 2011, Figure 1 (2011).
- [2] PEACE, M., MCCAW, L., SANTOS, B., KEPERT, J. D., AND FAWCETT, R. J. B. Meteorological drivers of extreme fire behaviour during the Waroona bushfire , Western Australia , January 2016. 79–101.