

UNIVERSITY OF WOLLONGONG

DOCTORAL THESIS

Thesis Title

Author:
Jesse GREENSLADE

Supervisor:
Dr. Jenny FISHER

*A thesis submitted in fulfillment of the requirements
for the degree of Doctor of Philosophy
in the*

Centre of Atmospheric Chemistry
Chemistry Department

March 23, 2018

Declaration of Authorship

I, Jesse GREENSLADE, declare that this thesis titled, "Thesis Title" and the work presented in it are my own. I confirm that:

- This work was done wholly or mainly while in candidature for a research degree at this University.
- Where any part of this thesis has previously been submitted for a degree or any other qualification at this University or any other institution, this has been clearly stated.
- Where I have consulted the published work of others, this is always clearly attributed.
- Where I have quoted from the work of others, the source is always given. With the exception of such quotations, this thesis is entirely my own work.
- I have acknowledged all main sources of help.
- Where the thesis is based on work done by myself jointly with others, I have made clear exactly what was done by others and what I have contributed myself.

Signed:

Date:

“Thanks to my solid academic training, today I can write hundreds of words on virtually any topic without possessing a shred of information, which is how I got a good job in journalism.”

Dave Barry

Contents

Declaration of Authorship	iii
1 Introduction and Literature Review	1
1.1 The atmosphere	1
1.1.1 Structure	2
1.1.2 Composition	3
Hydroxyl radicals	3
1.2 Ozone	3
NOx	4
1.2.1 Stratosphere to troposphere transport	6
1.2.2 Chemical production	7
1.3 VOCs	8
1.3.1 Emissions	10
1.3.2 Isoprene	10
1.3.3 The isoprene cascade	11
Oxidation	11
Low NOx scenario	12
Ozonolysis	13
Oxidation by NO ₃	14
1.3.4 Radiative Forcing	14
1.4 HCHO	17
1.4.1 Sources and sinks	17
1.4.2 Measurement techniques	18
Satellite measurements	20
1.5 Atmospheric Chemistry Modelling	21
1.5.1 Box models	22
1.5.2 Emissions	23
1.5.3 Uncertainties?	23
Emissions Inventories	24
Resolution	24
Chemistry mechanisms	24
Clouds	25
Soil Moisture	25
1.6 Australia and the southern hemisphere	26
1.6.1 Ozone	28
1.6.2 VOCs	28
1.6.3 Measurements	29
1.7 Aims	30

2 Data and Modelling	31
2.1 Introduction	31
2.2 List of runs and outputs used in my work TODO: good place for this?	31
2.2.1 GEOS-Chem	31
2.2.2 CAABA/MECCA	32
2.2.3 Reading Data	32
CAABA/MECCA outputs	32
GEOS-Chem Satellite output	32
HEMCO diagnostics	32
2.3 GEOS-Chem	32
2.3.1 GEOS-Chem isoprene modelling	33
Outline	33
Emissions from MEGAN	35
2.3.2 Chemical Mechanisms	37
2.3.3 Running GEOS-Chem (before isop?)	37
Installation and requirements	37
Outputs	37
Tropospheric chemistry run	38
UCX run	38
2.3.4 Run comparisons	38
2.4 Measurement Techniques	42
2.4.1 DOAS	42
2.4.2 Satellites	44
OMI	44
AMF	45
Uncertainties	46
2.4.3 OMI Satellite measurements	48
OMI HCHO	48
OMI NO ₂	48
2.5 Campaigns	48
2.5.1 Marine and Urban MBA ? (MUMBA)	48
2.5.2 Sydney Particle Studies (SPS1, SPS2)	48
2.6 Analysing output	49
2.6.1 Circadian emissions cycle	49
2.6.2 HCHO: Simulated vs Measured	49
2.6.3 Accounting for Fires	50
2.6.4 Accounting for NO _x	50
2.6.5 HCHO Comparisons	63
2.7 Data Access	63
Bibliography	65

List of Figures

1.1	Pressure (red) logarithmically decreasing, shown with percentage of atmosphere below at several points. Temperature (green) changes throughout the atmosphere. Figure edited from https://climate.ncsu.edu/edu/Structure	2
1.2	Tropospheric ozone processes, Figure 1 in Young et al., 2017. DOI: https://doi.org/10.1525/elementa.265.f1	5
1.3	Ozone production rate dependent on NO _x and VOC concentrations (Mazzuca et al., 2016).	8
1.4	The overall radiative forcings and uncertainties of several atmospheric constituents This is an image taken from Forster et al., 2007, found at https://www.ipcc.ch/publications_and_data/ar4/wg1/en/faq-2-1.html	15
1.5	The overall radiative forcings and uncertainties of several atmospheric constituents This is an image taken from IPCC, 2013: <i>Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change</i> , chapter 8.	16
1.6	HCHO spectrum, with a typical band of wavelengths used for DOAS path measurements. This is a portion of an image from Davenport et al., 2015.	19
1.7	An example spectrum showing interferences used for species concentration measurements by GOME-2. Image by EUMETSAT and ESA (EUMETSAT, 2015).	20
1.8	Standard box model parameters, image taken from Jacob, 1999.	22
1.9	Forest types in Australia http://www.agriculture.gov.au/abares/forestsaustralia/australias-forests	27
1.10	Part of a figure from Guenther et al., 2006 showing global isoprene emission factors.	27
2.1	MEGAN schematic, copied from Guenther (2016)	35
2.2	Surface HCHO simulated by GEOS-Chem with UCX (top left), and without UCX (top right), along with their absolute and relative differences(bottom left, right respectively). Amounts simulated by GEOS-Chem for the 1st of January, 2005.	39
2.3	As figure 2.2, except looking at isoprene.	40
2.4	As figure 2.2, except looking at ozone.	41
2.5	Image from Lee et al., 2015.	43

2.6	Figure 1 and Table 1 from Schenkeveld et al., 2017, with the following caption “An impression of OMI flying over the Earth. The spectrum of a ground pixel is projected on the wavelength dimension of the charge-coupled device (CCD; the columns). The cross-track ground pixels are projected on the swath dimension of the CCD (the rows). The forward speed of 7 kms^{-1} and an exposure time of 2 s lead to a ground pixel size of 13 km in the flight direction. The viewing angle of 114° leads to a swath width on the ground of 2600 km.” The table shows the optical properties for OMIs three channels.	45
2.7	Top panel: surface temperature averaged over January 2005. Bottom panel: surface temperature correlated against temperature over January 2005, with different colours for each gridbox, and the combined correlation. A reduced major axis regression is used within each gridbox (shown in top panel) using daily overpass time surface temperature and HCHO amounts (ppbv). The distribution of slopes and regression corellation coefficients (one datapoint per gridbox) for the exponential regression is shown in the embedded plot.	51
2.8	As figure 2.7 but for northern Australia.	52
2.9	As figure 2.7 but for south-western Australia.	53
2.10	GEOS-Chem mid-day tropospheric column NO_2 vs OMNO2d columns (averaged to match GEOS-Chems lower resolution). Absolute and relative differences, along with corellation shown on bottom row.	54
2.11	Top row (left to right): GEOS-Chem NO_2 mid-day tropospheric columns, OMNO2d NO_2 columns, modelled anthropogenic NO emissions. Second row: absolute and relative difference between GEOS-Chem and OMI NO_2 data, and the corellation. Third row: corellation between GEOS-Chem tropospheric column NO_2 and emitted NO, then between the model-satellite bias and the emissions. All corellation plots are coloured by emission rates.	55
2.12	As figure 2.11, for Autumn 2005.	56
2.13	As figure 2.11, for Winter 2005.	57
2.14	As figure 2.11, for Spring 2005.	58
2.15	As figure 2.11, except anthropogenic NO emissions are replaced by soil NO emissions.	59
2.16	As figure 2.11, for Autumn 2005, with soil NO emissions replacing anthropogenic NO emissions.	60
2.17	As figure 2.11, for Winter 2005, with soil NO emissions replacing anthropogenic NO emissions.	61
2.18	As figure 2.11, for Spring 2005, with soil NO emissions replacing anthropogenic NO emissions.	62

List of Tables

Chapter 1

Introduction and Literature Review

1.1 The atmosphere

The atmosphere is made up of gases held to the earth's surface by gravity. These gases undergo transport on all scales, from barbecue smoke being blown about the garden, to smoke plumes from forest fires travelling across the world and depositing in the Antarctic snow. They take part in innumerable chemical reactions along the way, largely driven by solar input and interactions with each other. Many gases are lofted into the atmosphere by soil, trees, factories, cars, seas and oceans. They are also deposited back to the surface both directly and in rainfall.

The atmosphere is made up of nitrogen (N_2 : ~ 78%), oxygen (O_2 : ~ 21%), and argon (Ar : ~ 1%), along with water (H_2O) and *trace gases* (those that make up less than 1% of the atmosphere). Water (H_2O) ranges from 0.001 to 1% depending on evaporation and precipitation. Beyond these major constituents the atmosphere has a vast number of *trace gases*, including carbon dioxide (CO_2 : ~ 0.4%), Ozone (O_3 : .000001 to 0.001%), and methane (CH_4 : ~ 0.4%) (Brasseur and Jacob, 2017, Ch. 2). Trace gases in the atmosphere can have a large impact on living conditions. They react in complex ways with other elements (anthropogenic and natural), affecting all surface ecosystems upon which life depends.

Ozone in the lower atmosphere is a serious hazard that causes health problems (Hsieh and Liao, 2013), damages agricultural crops worth billions of dollars (Avnery et al., 2013; Yue et al., 2017), and increases the rate of climate warming (Myhre and Shindell, 2013). Around 5 to 20 percent of all air pollution related deaths are due to ozone (Monks et al., 2015), roughly .8 million deaths per year (Lelieveld et al., 2013). In the short term, ozone concentrations of ~50-60 ppbv over eight hours or ~80 ppbv over one hour are agreed to constitute a human health hazard (Ayers and Simpson, 2006; Lelieveld et al., 2009). Long term exposure causes problems with crop loss and ecosystem damage (Ashmore, Emberson, and Murray Frank, 2003), and concentrations may get worse in the future (Lelieveld et al., 2009; Stevenson et al., 2013). Further tropospheric ozone enhancements are projected to drive reductions in global crop yields equivalent to losses of up to \$USD₂₀₀₀ 35 billion per year by 2030 (Avnery et al., 2013), along with detrimental health outcomes equivalent to ~\$USD₂₀₀₀ 11.8 billion per year by 2050 (Selin et al., 2009). Recently Yue et al., 2017 showed that the net effect of near-surface ozone on is a ~ 14% decrease in net primary productivity (NPP) in China. They state that reducing this decrease by ~ 70% before 2030 would require drastic measures.

This is a much better start. I'd suggest an additional framing paragraph about trop. O₃ impacts on climate

Don't
capitalise
Ozone

acronym not
used again,
delete (NPP)

Transition needed, e.g. to link
from generic trace gases to
specific focus on Ozone.

May be a good place to state
that this thesis focuses
specifically on ozone in the
troposphere.




FIGURE 1.1: Pressure (red) logarithmically decreasing, shown with percentage of atmosphere below at several points. Temperature (green) changes throughout the atmosphere. Figure edited from <https://climate.ncsu.edu/Structure>.

1.1.1 Structure

Most of the atmosphere ($\sim 85\%$) is within 10 km of the earth's surface. This is due to air pressure, which decreases logarithmically with altitude. Any entity is subjected to the weight of all the air above it, and the density of the atmosphere is driven by this pressure.

The atmosphere extends above us to the edges of space. This is split into various layers, defined by the *lapse rate*: the decrease in temperature with increasing altitude, or $\frac{-dT}{dz}$. Figure 1.1 shows the pressure and temperature profiles against altitude through the atmosphere. The first layer is the troposphere, which extends to roughly 10 km and is characterised by positive lapse rate (or decreasing temperature with altitude). At the top of the troposphere (the tropopause) the temperature stops decreasing, and then the stratosphere is defined by a negative lapse rate. This is due to UV radiation being absorbed by ozone, and leads to a very vertically stable environment.

In addition to these atmospheric layers, the troposphere can be subdivided into a boundary layer and the free troposphere. The *boundary layer* is the lowest layer and involves increased atmospheric mixing due to ground heating and friction effects. It generally extends anywhere from 200 - 1000 m, above which the ground effects have fewer direct impacts. The *free troposphere* is the remainder of the troposphere and is more affected by transport, both horizontally and from the stratosphere.

too informal - Earth's surface?
lower case z, all vars should be italicised, need to explicitly state what T & z are.

1.1.2 Composition & Chemistry?

Good idea →

To address 'Composition' in title, perhaps a sentence or two about diversity of trace gases.


TODO overview here Oxidation and photolysis are the two main processes whereby compounds are broken down in the atmosphere.

Hydroxyl radicals - I don't think you need a section heading here

OH and HO₂ concentrations largely determine the oxidative capacity of the atmosphere. The OH radical drives many processes in the atmosphere, especially during the day when photolysis of ozone drives OH concentrations (Atkinson, 2000). OH is a key species which reacts with nearly all the organic compounds in the troposphere, with only a few exceptions (Atkinson, 2000). Over land, isoprene (C₅H₈) and monoterpenes (C₁₀H₁₆) account for 50% and 30% of the OH reactivity (defined in section TODO) respectively (Fuentes et al., 2000).

Since radicals are involved in all oxidative chemistry in the atmosphere it is important for models to accurately represent them (eg. Travis2014). This is difficult as they are coupled with so many other species and measurements of OH are not readily available on a global scale. In the late 90's it was thought that OH radicals were formed exclusively from photolysis of O₃, HONO, HCHO, and other carbonyls (R₂C=O) Atkinson, 2000. It has been shown since that TODO. Isoprene (C₅H₈) was thought to be a sink of OH until it was shown by Paulot et al. (2009b) that the radicals are recycled. This recycling process is discussed in more detail in section 1.3.3.

Ozone is an important precursor to OH, as excited oxygen atoms (O(¹D)) are created through its photolysis, which then go on to react with water to form OH, as shown in this reaction sequence (Atkinson, 2000; Atkinson and Arey, 2003):



Where $h\nu$ represents radiation and M is an inert molecule. This shows that some of the O(¹D) recycles back to ozone, while some forms OH. Atkinson and Arey, (2003) discuss the relative rates of these reactions.

here i elsewhere:
Authors (year) show...


1.2 Ozone

1.2.1 Stratospheric Ozone

Ozone (O₃) is mostly located in the stratosphere, where it prevents much of the shorter wavelength (UV) solar radiation from reaching the earth's surface. In the stratosphere ozone production is driven by the Chapman mechanism, as high energy radiation (with wavelengths $\lambda < 242$ nm) photolyses the molecular oxygen (O₂) in the atmosphere (Brasseur and Jacob, 2017, Chapter 3, section 2).

Not a meaningful sentence.
Either summarise what →
they say or delete.

The Chapman mechanism involves several reactions which lead to rough equilibrium of O, O₂, O₃ and pressure, as follows:



The high energy photons ($\lambda < 242$ nm) are present from the top of the atmosphere but are mostly removed before reaching the troposphere as their energy is used to split the O₂ molecules. The lifetime of O against loss by O₂ is less than a second in the troposphere, and produced O₃ quickly returns to O and O₂, as low energy ($\lambda < 1180$ nm) photons and M are abundant. The reduced light penetration towards the surface, in addition to the logarithmic increase in atmospheric pressure (which affects M abundance) drives the vertical profile of ozone into what is called the *ozone layer*. This is a layer of relative ozone abundance within the stratosphere. The Chapman mechanism requires radiation so only takes place during the daytime, during the night this process slows to a halt, and the ozone concentrations remain stable unless NO_x pollution intrudes (Jacob, 1999, Chapter 10).

Since the Montreal Protocol on Substances that Deplete the Ozone Layer was established in August 1987, and ratified in August 1989, several satellites and many measurement stations were set up to monitor ozone in the stratosphere. However, in the southern hemisphere there are relatively few records of ozone (Huang et al., 2017). This affects our ability to accurately determine sources of ozone in the troposphere.

Generally there are two main drivers of tropospheric ozone concentrations; transport from the stratosphere and chemical production due to emissions of precursors. Tropospheric ozone is regulated by NO and NO₂ concentrations, which form an equilibrium (Cape, 2008; Young et al., 2017). However NO_x or VOC emissions affect this equilibrium and can lead to enhanced ozone formation, this can be seen in figure 1.2. At small to medium scales, pyrogenic (fire) and anthropogenic (man-made) emissions can be important. Smoke plumes from biomass burning can carry ozone precursors, creating higher ozone concentrations downwind of the plume's source. Emissions of precursors from large cities (such as NO_x emissions from traffic and power production) can impact ozone concentrations. These impacts are not always straightforwards due to the nonlinear relationship between ozone and its precursors.

A summary of processes affecting tropospheric ozone, copied from Young et al., 2017, is shown in Figure 1.2. This shows the major processes and emissions which affect tropospheric ozone concentrations. My work involves improving the highly uncertain natural emissions of NMVOCs from Australia. ↗ move to be first text in 'Tropospheric ozone' Section 1.2.2. ↗ This thesis

NO_x *not yet defined*

NO_x ($\equiv \text{NO}_2 + \text{NO}$) is another important chemical family in the atmosphere which interacts with ozone and regulates the atmospheric oxidative capacity. NO_x compounds are short lived, with emissions (Power generation and combustion transport) being the main driver of concentrations (Delmas, Serca, and Jambert, 1997). NO_x and O₃ relative concentrations during the day are regulated by the following reactions

Instead of a separate sub-section, integrate into the above where you state that NO & NO₂ regulate trop. O₃.





FIGURE 1.2: Tropospheric ozone processes, Figure 1 in Young et al., 2017. DOI: <https://doi.org/10.1525/elementa.265.f1>

(Sillman, 1999; Atkinson, 2000):



This doesn't look right.
Should be NO_2 photolysis not
 NO_3 . Check refs & textbooks (e.g.
Jacob 1999).

NO_x is removed primarily by conversion to nitric acid (HNO_3) followed by wet or dry deposition (Ayers and Simpson, 2006).

1.2.1 Stratosphere to troposphere transport

citcs papers from 2000s... is this even needed?

Historically (in the late 1990's), ozone transported down from the stratosphere was thought to contribute 10-40 ppb to tropospheric ozone levels, matching tropospheric production (Atkinson, 2000; Stohl et al., 2003). This number was revised down over the years as measurement and modelling campaigns improved our understanding of global scale transport, mixing, and chemistry (Monks et al., 2015). Recently Kuang et al., 2017 analysed tropospheric ozone measurements in the southeast USA and observed STT influence which can be seen to affect surface ozone levels. In their work they used several different instruments and analysed the structure and temporal evolution of ozone and the local weather systems. - again only include if you are going to summarise what they found

Ozone transported to the troposphere from the stratosphere can occur through diffusion (relatively slowly) or direct mixing. Intrusions of stratospheric air into the troposphere are often called Stratosphere to Troposphere Transport events (STT). STT often occur as tongues of stratospheric air descend and get disconnected from the stratosphere, potentially due to low pressure systems and jet streams Sprenger, Croci Maspoch, and Wernli, 2003. Recently global chemical transport models (CTMs) have been used to trace how much ozone is being transported to the troposphere in this manner. There are a few methods of doing this, such as Ojha et al., 2016, who use a model that keeps track of ozone formed in (and transported from) the stratosphere. Model based estimates require validation against actual measurements, such as those from ozonesondes or satellites. Hegglin and Shepherd, 2009 estimate that climate change will lead to increased STT due to an acceleration in the Brewer Dobson circulation. They estimate ~ 30 , and $\sim 121 \text{ Tg yr}^{-1}$ increases (relative to 1965) in the southern and northern hemispheres respectively. No context for these numbers. Increase relative to what??

Liu et al., 2017 examine southern hemispheric ozone and the processes which control its inter-annual variability (IAV). IAV is the standard deviation of ozone anomalies from the monthly mean. They show that ozone transported from the stratosphere plays a major role in the upper troposphere, especially over the southern Indian ocean during austral winter. While stratospheric transport mostly impacts the upper troposphere, some areas are impacted right down to the surface. Liu et al., 2017 look at modelled tropospheric ozone sensitivity to changes in stratospheric ozone, ozone precursor emissions, and lightning over the southern hemisphere from 1992–2011. Their work suggests ozone at 430 hPa (roughly 6 km altitude) is mostly stratospheric in

tell us what they found, not what they did.

need to explain what this is if you refer to it.

Description of Liu could be shortened.

Note: For published papers use past tense when you describe what they did; e.g. Authors used, authors found.

If finding is generally applicable, that can be present tense e.g. Authors found STT explains 10% of trop. O₃ or whatever. Fix throughout.

September over 20°S to 60°S at all longitudes. They also see tropospheric ozone sensitivity to emissions from South America (0–20°S, 72.5–37.5°W), southern Africa (5–10°S, 12–38°E), and South to Southeast Asia (70–125°E, 10°S–40°N). In the USA recent work by Lin et al., 2015 suggests that intrusions during spring are increasing surface ozone levels. Their work also recommends that understanding of the frequency and cause of STT needs to be improved to effectively implement air quality standards.

1.2.2 Chemical production

Ozone produced in the troposphere from precursors and radiation drive ozone levels, especially in the lower (near-surface) troposphere. The main processes involved are shown in figure 1.2, with ozone regulated by reactions 1.3. An analysis of the Atmospheric Chemistry and Climate Model Inter-comparison Project (ACCMIP) simulations by Young et al., 2013 found STT is responsible for $540 \pm 140 \text{ Tg yr}^{-1}$, equivalent to ~11% of the tropospheric ozone column, with the remainder produced photochemically (Monks et al., 2015). A recent summary by Young et al., 2017 estimates ozone production and loss in the troposphere to be $\sim 4900 \text{ Tg yr}^{-1}$, and $\sim 4500 \text{ Tg yr}^{-1}$ respectively, while STT sources are $\sim 500 \text{ Tg yr}^{-1}$. These numbers are at the global scale, and in order to understand the processes driving ozone concentrations at any specific site measurements must be analysed.

doesn't mean much → just reference large spatial variability

Tropospheric ozone concentrations rely on climate and ozone precursor emissions; including NO, NO₂, CO, VOCs, and HCHO (Atkinson, 2000; Young et al., 2013; Marvin et al., 2017). Ozone predictions are uncertain due to the vagaries of changing climate which affects both transport, deposition, destruction, and plant based precursor emissions. All of these processes are tightly coupled and difficult to accurately model, as they depend on uncertain assumptions such as CO₂ dependency (Young et al., 2013). Even with all the work done over the prior decades there remains large uncertainties about ozone precursors in the troposphere (Mazzuca et al., 2016).

Ozone is formed in the troposphere through oxidation of VOCs (described in Section 1.3) in the presence of NO_x. Net formation or loss of O₃ is determined by interactions between VOCs, NO_x, and HO_x, and is a complicated system of positive and negative feedbacks (Atkinson, 2000). Figure 1.3 shows the non-linear effect of NO_x and VOC concentrations on ozone production over Houston, as modelled in Mazzuca et al., 2016. Recently the relationship has been examined on the intradiel timescale showing that ozone production can be more or less sensitive to VOCs at different hours depending on location various other factors (Mazzuca et al., 2016). This shows how important it is to correctly determine the precursors concentrations in order to estimate ozone levels and production.

Tropospheric ozone is lost via chemical destruction and dry deposition, estimated to be $4700 \pm 700 \text{ Tg yr}^{-1}$ and $1000 \pm 200 \text{ Tg yr}^{-1}$, respectively (Stevenson et al., 2006; Young et al., 2017). The main loss channel is through equation 1.1, where photolysis and pressure create OH from the O₃.

precision! "Pressure" doesn't "create OH!"

*What about VOC chemistry?
Need to show that
somewhere*

*put this in
previous section*

*"rely on" not quite right...
require?*

HCHO is a VOC..

*not sure
this is
needed*

*include example
reaction sequence.*

*frame this discussion
more generally, with more
detail about NO_x vs. VOC
regimes. Mazzuca just a
recent example of long known
phenomenon, doesn't need
special attention.*




FIGURE 1.3: Ozone production rate dependent on NO_X and VOC concentrations (Mazzuca et al., 2016).

1.3 VOCs Spell out

The least well understood precursors to tropospheric ozone production belong to a class of organic compounds. Organic compounds are members of a large class of chemicals whose molecules contain carbon, with the exception of a few compounds such as carbides, carbonates (CO₃), and simple oxides of carbon and cyanides. Organic compounds can be categorised based on their vapour pressure, which is the tendency of a liquid or solid to vaporise. Compounds with high vapour pressures at standard temperature are classed as volatile, and have a **felicity** to evaporate at low temperatures. Plants contain tens of thousands of organic compounds, it's likely that fewer than 40 have high enough volatility to be emitted (Guenther et al., 2000).

Atmospheric organic compounds are legion and differ by orders of magnitude with respect to their fundamental properties, such as volatility, reactivity, and cloud droplet formation propensity, etc. Volatile organic compounds (VOCs) have vapour pressure greater than 10⁻⁵ atm, and are mostly generated naturally by plants, which emit around 1000 Tg yr⁻¹ (Guenther et al., 1995; Glasius and Goldstein, 2016). Due to their high volatility these compounds are generally seen in the gas phase. Organic compounds with a lower volatility are classed as semi-volatile (SVOCs: vapour pressure between 10⁻⁵ and 10⁻¹¹ atm) are seen in both gas and particle phase depending on temperature and pressure. Organic compounds with even lower vapour pressure are generally found in the particle phase in aerosol particulate matter (Glasius and Goldstein, 2016). Understanding the drivers of trends in biogenic VOC emissions (BVOCs) is required in order to estimate future carbon fluxes, changes in the water cycle, ozone production, air quality, and other climate responses (Yue, Unger, and

not correct as written (CO₂⁻)
but not sure why it's here when
no other chem. formulas given.

?
avoid 'it's'. Also should be
a new sentence.

by who? (word choice...)

Zheng, 2015). In the last 20 years anthropogenic emissions of VOCs have been increasing while biogenic VOC emissions have decreased, due to rapid economic growth and lower annual temperatures (Stavrakou et al., 2014; Kwon et al., 2017).

in what sense? I'd delete


"include"
isoprene is an alkene...
"abundant?"

Methane (CH_4) is one of the more abundant and potent VOCs, however it is often classified separately and compared against non-methane VOCs (NMVOCs). NMVOCs are alkanes, alkenes, aromatic hydrocarbons and isoprene, with isoprene being the most prominent. Methane is relatively long lived (years) and is well mixed in the atmosphere while other VOC levels are spatially diverse due to their shorter lifetimes. In this thesis I work towards a better understanding of the NMVOC emissions coming from Australia.

just isoprene right?

VOCs are an important driver of atmospheric processes, especially near forests. VOCs are broken down into HCHO , O_3 , CO_2 and many other species, mainly through oxidation by OH. VOC emissions result in radical cycling, acid deposition, production of tropospheric ozone, and secondary organic aerosols (SOAs) (Atkinson, 2000; Kanakidou et al., 2005). VOC emissions affect surface pollution levels, potentially enhancing particulate matter (PM) and ozone levels. A regional-model study in Europe (Aksoyoglu et al., 2017) has also shown VOCs impact secondary inorganic aerosol concentrations. These have impacts on climate (through radiative forcing) and air quality (from ozone and SOA enhancements), affecting both human health and crop yields (Forster et al., 2007; Avnery et al., 2013; Lelieveld et al., 2015).

do you have a ref. for this? Ozone in rural areas is often higher than in populous cities, due to the more abundant VOCs concentrations. This occurs through the following reaction sequence (Sillman, 1999):



The reactions of VOCs or CO with OH convert NO to NO_2 , which leads to ozone formation as NO_2 production in reaction 1 of 1.3 is bypassed. R and R' are organic species in these reactions.

One problem associated with VOC emissions is the production of aerosols. Aerosols are suspended particulates and liquid compounds in the atmosphere, of which particulate matter (PM) is an important subset. PM in the atmosphere is a major problem, causing an estimated 2-3 million deaths annually (Hoek et al., 2013; Krewski et al., 2009; Silva et al., 2013; Lelieveld et al., 2015). Fine particulate matter ($\text{PM}_{2.5}$) penetrates deep into the lungs and is detrimental to human health. Some PM comes from small organic aerosols (OA) emitted in the particulate phase and referred to as primary OA (POA). A substantial amount of PM is due to gaseous organic compounds transforming in the troposphere leading to what's known as secondary OA (SOA) (Kroll and Seinfeld, 2008).

Formation of SOA is generally due to VOC oxidation and subsequent reactions, while removal from the atmosphere is largely due to wet or dry deposition, and cloud scavenging (Kanakidou et al., 2005). It can be difficult to attribute PM formation, in part due to the complex relationship between NO_x , OH, O_3 , and the uncertainty surrounding precursor emissions. Improved concentration estimates of these organic

"problem" implies value judgement. Stay neutral.
isn't PM just another name
for aerosols? Unless you
mean $\text{PM}_{2.5}$?

avoid "what's"

is this about PM or SOA?
be consistent

compounds requires a better understanding of their emissions, which is one of the foci in this thesis.

1.3.1 Emissions

VOC emissions are often classified as either anthropogenic, biogenic, or pyrogenic. There are ten times as many emissions of NMVOCs from natural sources compared to anthropogenic sources (Guenther et al., 2006; Kanakidou et al., 2005; Millet et al., 2006). Methane and isoprene each comprise around a third of the global total emissions of VOCs (Guenther et al., 2006). Major emitters are tropical broadleafs (notably eucalypts), and shrubs (Guenther et al., 2006; Arneth et al., 2008; Niinemets et al., 2010; Monks et al., 2015). TODO: why do plants emit? increased canopy light penetration? Emissions are affected by various factors such as temperature, sunlight, soil moisture, etc.

It used to be thought that emissions of anthropogenic and biogenic VOCs (AVOCs, BVOCs respectively) were roughly similar (Müller, 1992, TODO: more cites). In the 1990's it became clear that biogenic emissions are in fact dominant. Global VOC levels are estimated at 85 %, 13 %, and 3 % from biogenic, anthropogenic, and pyrogenic sources respectively (Kanakidou et al., 2005; Kefauver, Filella, and Peñuelas, 2014). Although methane makes up a third of atmospheric VOCs, its emissions are low. The World Meteorological Organisation (WMO) estimated that we are emitting 360 Mt yr^{-1} of methane, compared to biogenic emissions of around 200 Mt yr^{-1} (Atkinson, 2000). In 1995 emissions of other VOCs were estimated at 1150 Tg C yr^{-1} from biogenic sources, and 100 Tg C yr^{-1} from anthropogenic sources (Guenther et al., 1995; Atkinson, 2000). The main non-methane BVOC emissions are isoprene (44%) and monoterpenes (11%) (Guenther et al., 2000; Kefauver, Filella, and Peñuelas, 2014). Land use changes can drastically affect isoprene sources, for instance in the tropics where large scale deforestation has converted forest into crop lands (Kanakidou et al., 2005).

VOCs are removed by wet and dry deposition, OH oxidation, reaction with NO_3^- , ozonolysis (at night time or in polluted areas) or photolysis (Atkinson and Arey, 2003; Brown et al., 2009). The process of deposition only accounts for a small fraction of the VOC loss, with the possible exception of the long lived methane compound (Atkinson and Arey, 2003).

1.3.2 Isoprene

Isoprene, or 2-methylbuta-1,3-diene, is a VOC with the chemical formula C_5H_8 . It is of major importance to the atmosphere, as it is involved in various processes which alter the oxidative capacity of the atmosphere. Guenther et al., 1995, and subsequent updates (Guenther et al., 2000; Guenther et al., 2006; Guenther et al., 2012), have been used ubiquitously by the atmospheric community as a global estimate of isoprene emissions, at roughly 500-600 Tg yr^{-1} , emitted mostly during the day. Recently an estimate of global isoprene emissions has been made using a completely different model, [of around 465 Tg C yr^{-1}] by (Messina et al., 2016) using ORCHIDEE. The global emission factors ~~model~~ used to derive both these estimates are based on modelling emissions from different plant species (phenotypes), and relatively few Australian species are used when forming in these estimates.

careful w/ language. Could mean more species emitted or more mass (I think the latter)

really? I thought eucalypts largely temperate?

probably irrelevant

?? Your next sentence confusing too. Just state earlier that you'll only discuss NMVOC and remove methane stuff.

1995! Use more recent estimate

} doesn't belong in section called "emissions"

Since so much of your thesis is about reducing uncertainty in isoprene emissions, you probably want to expand the discussion (and refs) around emission uncertainties and how they differ from place to place.

1.3. VOCs

11

HO_x?

Isoprene affects NO_x and HO_x cycling, see for example formulae 1.1, 1.3. In the presence of NO_x, isoprene forms tropospheric ozone and SOAs (Wagner, 2002; Mille et al., 2006). It has a short lifetime during the day, roughly an hour due to OH oxidation (Atkinson and Arey, 2003)).

As before, only include if you are going to summarise findings (and they are relevant). No one cares what others did.

Measurements of isoprene are often uncertain or difficult to make accurately. Kanakidou et al., 2005 summarised how chamber experiments used to measure isoprene reactions may be unsuitable for comparison to the natural atmosphere. In Nguyen et al., 2014 many scientists and groups worked together on chamber measurements, to improve understanding of ambient atmospheric oxidation mechanisms of biogenic hydrocarbons (such as isoprene).

Isoprene emissions estimates are still fairly uncertain, as global measurements are difficult and regional emissions can be very different. The global uncertainty of isoprene emission was estimated to be a factor of 2 to 5 (250-750 Tg yr⁻¹) (Kanakidou et al., 2005). Improvements over the years have been incremental, and generally localised to regions of particular interest for air quality such as China and the USA TODO: find recent uncertainty estimate improvements examples. The lack of accuracy in BVOC emissions estimates prevents accurate determinations of the sources and distribution of pollutants including ozone and organic aerosols. Most of the tropospheric SOA comes from biogenic precursors, the evidence for this has grown over the last two decades (Guenther et al., 1995; Kanakidou et al., 2005; Guenther et al., 2012). Accuracy in VOC measurements is important: it has been shown that even the diurnal pattern of isoprene emissions has an effect on modelling ground level ozone (Hewitt et al., 2011; Fan and Zhang, 2004).

1.3.3 The isoprene cascade Isoprene chemistry?

relevant

Isoprene forms many products with various lifetimes, here I will present an overview of some important mechanisms and products which are useful to my work. Isoprene is emitted and enters the atmosphere in the gas phase, where it reacts quickly with OH and other radicals. One common compound which is produced by these reactions is HCHO, which is easier to measure and often used to estimate how much isoprene is being emitted. The reactions which occur are important to understand due to their impacts on air quality, ozone, and physical properties in the lower troposphere. The many children processes and products which begin with isoprene oxidation are often called the isoprene (photochemical) cascade (Paulot 2012; Crounse et al., 2012; Wolfe et al., 2016.)

Photolysis and oxidation of many VOCs initially form alkyl radicals (\dot{R}). Alkenes (VOCs with double bonded carbon, such as isoprene) react with ozone leading to organic peroxy radicals (RO \dot{O}). These go on to form many products and lead to (amongst other things) aerosol, formaldehyde, and ozone formation, depending on sunlight and NO_x concentrations (Atkinson, 2000).

Oxidation

The primary first step for atmospheric isoprene is photooxidation, reacting with OH to form isoprene hydroxyperoxy radicals (RO \dot{O}) (Patchen 2017; Wolfe et al., 2016; Marvin et al., 2017). There is still uncertainty about which pathways are most important

Agree - therefore you need to show at least part of this reaction sequence!

Your dots are in the wrong place. RO \dot{O} , R \cdot

instead of generic R, use isoprene products here!

There is a bit of redundancy and also some confusion in this section. Take the time to write out the first steps of the reaction sequence, specifically for isoprene (not generic R) and make sure you understand the steps and associated terminology.

following RO \cdot production: HO $_2$ reactions predominantly produce hydroxyhydroperoxides (ISOPOOH), NO reactions largely produce methyl vinyl ketone (MVK) and methacrolein (MCR), and RO $_2$ reactions are also possible (Liu et al., 2016b)

First isoprene has its double bond replaced by OH, as summarised by the equation from PATCHEN et al., 2007: R-CH=CH-R' + OH \longrightarrow R-CH(OH)CH-R' where R and R' represent hydrocarbons. This OH adduct then reacts with O $_2$ to produce a hydroxyperoxy radical (RO \cdot), which can be any of six different isomers (PATCHEN et al., 2007). These RO \cdot (also called organic-peroxy/alkyl-peroxyl/ISOPOO radicals, or RO $_2$) react with HO $_2$ or NO and produce stable products (often called oxidised VOCs or OVOCs) (Nguyen et al., 2014). Most of these reaction pathways produce HCHO (Wolfe et al., 2016).

RO \cdot reaction pathways depend on the NO $_x$ concentrations. Reactions with NO can lead to ozone production in environments rich in isoprene or other non-methane organic compounds (NMOCs) and NMVOCs (PATCHEN et al., 2007; Atkinson and Arey, 2003). These reactions are complex and coupled, for example NO $_2$ concentrations can be increased by NMOC and NO reactions (Atkinson and Arey, 2003).

In the presence of NO $_x$, the RO \cdot may form organic nitrates after reacting with NO. Any organic nitrates which are formed affect levels of both HO x (H, OH, peroxy radicals) and NO x , acting as a sink (Mao et al., 2013 and references therein). Reaction with NO $_2$ forms isoprene nitrates, or hydroxynitrate (RONO $_2$). The first generation of organic nitrates produced by isoprene oxidation range from 7% to 12%, shown in laboratory experiments (Paulot et al., 2009a; Mao et al., 2013). A portion of isoprene nitrates are recycled back to NO x , so may serve as a reservoir of nitrogen and allow its transport to the boundary layer of remote regions (PATCHEN et al., 2007; Paulot et al., 2009a). The nitrates can also build up in the winter, when removal processes are not as dominant (Lelieveld et al., 2009).

Oxidation reactions are important and quickly stabilise the ratio of NO to NO $_2$. There is still large uncertainty around the fate of various RO \cdot , which limits understanding of the relative importance of some chemical processes (Crounse et al., 2013). Some portion of emitted isoprene leads to SOA, potentially through the formation of methacrylic acid epoxide (MAE) formed by decomposition of methacryloylperoxynitrile (MPAN, a second generation product of isoprene oxidation) as shown in smog chambers and field studies in Lin et al., 2013.

Again, very patchy referencing. Not the dominant isop \rightarrow SOA pathway. Unless you are more comprehensive, delete.

Low NO x scenario

Isoprene oxidation by OH is less well understood when lower concentrations of NO are present in the atmosphere. Initially isoprene was thought to be a sink for atmospheric oxidants (e.g. Guenther et al., 2000). It was thought that in low NO environments, like those far from anthropogenic pollution and fires, oxidation of isoprene would create ISOPOOH and lead to low concentrations of OH and HO $_2$ (Paulot et al., 2009b). In Paulot et al., 2009b, the HO x levels are shown to be largely unaffected by isoprene concentrations. They show that ISOPOOH is formed in yields > 70%, and MACR and MVK is formed with yields < 30%. The formation of MACR and MVK produces some HO x , although not enough to close the gap. Paulot et al., 2009b goes on to suggest (and provide experimental evidence) that dihydroxyperoxides (IEPOX) are formed from oxidation of the ISOPOOH, which form precursors for SOAs as well

just another name for NMVOC

?

hot always! See my 2016 paper for discussion of possible fates.

You're missing a lot of the more recent lit here and this really isn't that relevant for your work, so I would delete

Such as...?

Condense. Your intro does not need to be a history lesson. Tell us the current state of the science. Again, focus on what found, not what done. Should see more "Thing we know (author, year" than "author (year) did thing."

as closing the HO_x concentration gap. They then use GEOS-Chem, modified to include IEPOX formation, to estimate that one third of isoprene peroxy radicals react with HO₂, and two thirds react with NO. They estimated $95 \pm 45 \text{ Tg yr}^{-1}$ IEPOX being created in the atmosphere, which (at the time) was not modelled by CTMs. Their work showed another pathway for isoprene based SOA creation, through these IEPOX creation and HO_x recycling mechanisms. Peeters and Muller, 2010 suggested that the work of Paulot et al., 2009b only partially bridges the gap between clean air OH concentration measurements and models. They suggested four new mechanisms for OH recycling in these pristine conditions. These can be summarised as OH regenerating reactions which occur during photolysis of hydroperoxy-methyl-buteneals (HPALDs), and resulting photolabile peroxy-acid-aldehydes (PACALDs). These reactions are highly non-linear and subject to large uncertainty, however when compared against several campaigns they were shown to improve modeled HO_x concentrations. In Crounse et al., 2012, MACR products are examined and hydroxy recycling is observed in low NO conditions, backing up results from Peeters and Muller, 2010. Peeters and Muller, 2010 showed that HO₂ is produced at near unity yields following isoprene oxidation initiated by OH. TODO: read more Peeters2010

Nguyen et al., 2014 examine various measurement techniques to determine isoprene reactions in non-laboratory conditions. Their work discussed how large uncertainties persist in isoprene oxidation, which carries through to predictions by atmospheric models. Nguyen et al., 2014 show preliminary estimates of low-NO yields of MVK and MCR to be $6 \pm 3\%$ and $4 \pm 2\%$ respectively, consistent with TODO:Liu2013 but only when cold-trapping methods are employed. These yields each increase (due to interference by OVOCs) to greater than 40% when directly sampled by GC-FID.

Even with the recent boom in isoprene analysis, uncertainties remain in the isoprene oxidation mechanisms. Examples (taken from Nguyen et al., 2014) include isoprene nitrate yields, which range from 4-15% (Paulot et al., 2009a), 90% disagreements in MAC and MVK yields TODO:(Liu2013), various possible sources for SOA TODO:(Chan2010; "Reactive intermediates revealed in secondary organic aerosol formation from isoprene"; Lin et al., 2013), unknown HPALD fates, incomplete O₂ incorporation TODO:(Peeters2009; Crounse et al., 2013), and under-characterized RO₂ lifetime impacts TODO:(Wolfe2012). TODO: get those citations and read abstracts.

Ozononlysis only relevant if you focus on isoprene ozonolysis...

Ozononlysis is the splitting of carbon chains by ozone molecules, and is among the primary oxidation pathways for volatile alkenes (Nguyen et al., 2016). Criegee intermediates (carbonyl oxides with two charge centres) are formed when isoprene reacts with O₃. Nguyen et al., 2016 examine in detail a few of these, with proposed mechanisms for C₁ and C₄ Criegee intermediate reactions. The C₁ stabilised Criegee (CH₂OO, ~61%) is therein proposed to react with water yielding 73% hydroxymethyl hydroperoxide (HMHP), 6% HCHO + H₂O₂, and formic acid + H₂O, and the same products with yields of 40, 6, and 54% respectively when this Criegee reacts with (H₂O)₂.

Oxidation by NO_x

At night when OH concentrations have dropped, isoprene can remain in the atmosphere ~~to be transported~~. Typically less than half of this night time isoprene is removed through ozonolysis (Atkinson and Arey, 2003), however, in polluted areas where high levels of NO_x exist, isoprene is consumed by ~~a different radical~~. During the night ~~isoprene is oxidised~~ no need for suspense! by nitrate radicals (NO₃), which joins to one of the double bonds and produces organic nitrates in high yield (65% to 85%) (Mao et al., 2013). (todo: read mao2013 para 3 cites) NO₃ are largely formed through ozone reactions, as in equation 1.3. A build up of NO₃ radicals can be seen at night, when photolysis is not removing them (Atkinson, 2000; Brown et al., 2009).

In areas with high NO_x levels, greater than 20% of the isoprene emitted late in the day ends up being oxidised by the NO₃ radical over ~~night~~ (Brown et al., 2009). At night isoprene has ~~affects~~ on both NO_x concentrations and ozone levels, and can form harmful organic nitrates and SOAs (Brown et al., 2009; Mao et al., 2013). These nitrates go on to produce further SOAs, largely due to NO₃ reacting with first generation isoprene oxidation products (Rollins et al., 2009). The night-time concentrations of OH and ozone also have a complex effect on NO_x removal in high latitude winters, when photolysis and NO reactions are reduced (Ayers and Simpson, 2006).

1.3.4 Radiative Forcing

This section doesn't really belong here. You could condense and put early in VOC section as motivation (1 paragraph max). It would move IPCC AR5 fig. to start of Chapter and use to highlight O3 RF.

One of the larger uncertainties in atmospheric modelling is how particles in the atmosphere affect radiative forcing. For 12 years it has been understood that most OA cool the atmosphere, with smaller particles having a larger effect as they matching the wavelengths of visible light (Kanakidou et al., 2005). Gas phase emissions with higher vapour pressures can be oxidised into lower vapour pressure products which will partition between gas and particle phase, often called semi or non-volatile. The aerosol products from these gas phase emissions (or the children thereof) are SOA (Kanakidou et al., 2005). SOA plays an indirect and complex role in cloud properties, with a net cooling effect (IPCC, 2013: *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*, Chapter 7,8)

Transport and indirect effects complicate matters further, with cloud creation and modification of cloud properties being quite difficult to accurately predict. In the third IPCC report (*Intergovernmental Panel on Climate Change (IPCC): Climate Change: The Scientific Basis* 2001), the uncertainty involved if OA forcing was a factor of 3 times the estimated effect. This has since been improved however OA and cloud formation still remains a large uncertainty in more recent IPCC reports (Forster et al., 2007). Figure 1.4 shows the radiative forcing (RF) of various atmospheric constituents, it's clear that OA uncertainty dominates. Figure 1.5 shows the same summary updated in chapter 8 of the fifth report, where the SOA uncertainty remains quite large.

In order to improve understanding of processes involved in radiative forcing, Kanakidou et al., 2005 highlight the need for improving VOC emissions and flux measurements. They also advocate utilising satellite data in models as a means of improving the emissions inventories. VOCs can lead to changes in cloud formation, as nucleation

- Two other notes:
 (1) can't just equate OA and total aerosol.
 (2) Hard to use a 13 year old paper as basis to frame current uncertainties.



FIGURE 1.4: The overall radiative forcings and uncertainties of several atmospheric constituents This is an image taken from Forster et al., 2007, found at https://www.ipcc.ch/publications_and_data/ar4/wg1/en/faq-2-1.html.




Figure 8.15 | Bar chart for RF (hatched) and ERF (solid) for the period 1750–2011, where the total ERF is derived from Figure 8.16. Uncertainties (5 to 95% confidence range) are given for RF (dotted lines) and ERF (solid lines).

Remove original legend.

FIGURE 1.5: The overall radiative forcings and uncertainties of several atmospheric constituents. This is an image taken from *IPCC, 2013: Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*, chapter 8.

can arise from the subsequent SOA. Kanakidou et al., 2005 concluded that it is very likely that organics contribute to particle growth and formation rates.


1.4 HCHO *spell full name*

One of the major products of isoprene chemistry is HCHO. HCHO is important both for its own atmospheric impacts, and as a proxy for determination of isoprene emissions. Given a modelled yield of HCHO from isoprene, it is possible to work backwards from measured HCHO concentrations to determine the isoprene emissions.

HCHO, aka methanal, methyl aldehyde, or methylene oxide, is of the aldehyde family. HCHO is an OVOC which is toxic, allergenic, and a potential carcinogen. It is dangerous at low levels, with WHO guidelines for prolonged exposure at 80 ppb. HCHO is used as an adhesive in plywood, carpeting, and in the creation of paints and wallpapers. Emissions in enclosed spaces can build up to dangerous levels, especially if new furnishings are installed (Davenport et al., 2015). At global scales HCHO in furniture is less important, as concentrations are driven by photochemical reactions with methane and other VOCs.

1.4.1 Sources and sinks

Background levels of HCHO in the atmosphere are driven by the oxidation of methane (CH_4) by the hydroxyl radical (OH^{\cdot}). Atkinson, 2000 summarised the background formation of HCHO with the following reaction:



which shows that photolysis and oxidation of methane forms HCHO and ozone in a process that regenerates the OH radicals. CH_4 concentrations are relatively well constrained in models, with the ACCMIP comparison showing only $\sim 3\%$ inter-quartile range (Young et al., 2013). There is a complex relationship between VOCs, HO_x , and NO_x : with higher levels of NO_x increase the speed that VOCs are converted into HCHO (Wolfe et al., 2016).

Within the continental boundary layer, the major source of HCHO enhancement is VOC emissions (which reacting with OH radicals in the presence of NO_x) (Wagner, 2002; Millet et al., 2006; Kefauver, Filella, and Peñuelas, 2014). Enhancements to regional and continental HCHO are largely driven by isoprene emissions (Guenther et al., 1995; Palmer, 2003; Shim et al., 2005; Kefauver, Filella, and Peñuelas, 2014). This is true except near fires or anthropogenic sources of HCHO and precursors (Guenther et al., 1995; Kefauver, Filella, and Peñuelas, 2014; Wolfe et al., 2016). Biomass burning (BB) can be a source of HCHO, and various other pollutants, precursors, and aerosols (Guenther et al., 1995; Andreae, 2001). Additionally HCHO is emitted into the atmosphere directly through fossil fuel combustion, natural gas flaring, ethanol refining, and agricultural activity (Wolfe et al., 2016).

Other terpenoids (monoterpenes, sesquiterpenes, etc.) can also produce HCHO, although generally to a lesser extent than isoprene, methane and biomass burning (Guenther et al., 2012). Many of the HCHO yields from terpenoids are estimated

"low" is relative; typical outdoor values at ppt levels, right?

Better to state outright why relevant to this thesis and just focus on outdoor chem.

OH^{\cdot}

Check grammar in this sentence

through chamber studies which examine the products molecular mass and charge after mixing the compound of choice into a known volume of air (Nguyen et al., 2014, eg.). These conditions generally don't match those of the real world, where ambient air will have a cocktail of these compounds and other reactants. Nguyen et al., 2014 state that one of their goals is to recreate ambient atmosphere in their chamber studies with more accuracy, in order to improve interpretations and allow more accurate model parameters.

Millet et al., 2008 show that anthropogenic emissions of HCHO in America are mostly negligible, although improved sensitivity from oversampling allowed satellite detection of enhanced HCHO concentrations over Houston and Texas (Zhu et al., 2014). This is not the case in China, since massive population centres and industrial districts are emitting huge amounts of VOCs into the atmosphere (Fu et al., 2007). Fu et al., 2007 use GOME measurements over Asia and derive biogenic, anthropogenic, and pyrogenic VOC emissions, and Zhu et al., 2014 use oversampling of the OMI HCHO measurements to determine anthropogenic highly-reactive VOC emissions. Then with their updated emissions they show how surface ozone is affected, with a seasonal increase of 5-20 ppb simulated by GEOS-Chem.

In the past, HCHO levels were underestimated by models, often with large discrepancies, due to the poor understanding of methyl peroxy radical (CH_3OO) chemistry (Wagner, 2002). Nowadays HCHO concentrations are better understood, however precursor emissions are one of the main unknowns (Emmerson et al., 2016; Marvin et al., 2017, eg.). Marvin et al., 2017 found that discrepancies in modelled HCHO concentrations are primarily due to second and later generation isoprene oxidation chemistry.

HCHO has two major sinks, one being reactions with OH (oxidation), the other being photolysis: the process of being broken apart by photons (CRUTZEN, LAWRENCE, and PÖSCHL, 1999; Wagner, 2002; Levy, 1972; Kefauver, Filella, and Peñuelas, 2014). These reactions lead to a daytime lifetime of a few hours (Atkinson, 2000; Millet et al., 2006). Both these loss processes (photolysis, oxidation) form CO and hydroperoxyl radicals (HO_2), and have global significance to radiative forcing and oxidative capacity (Franco et al., 2015). The other notable sinks are wet and dry deposition, although these are not as significant (Atkinson, 2000) (TODO: add more cites here).

only relevant if they did it and found something new!

this paragraph wanders, relevance not clear

you have used/defined previously, don't re-define

"notable" and "not significant"? give numbers to back-up.

1.4.2 Measurement techniques

There are a few ways to measure HCHO, including Fourier Transform Infra-Red (FTIR) Spectrometry and Differential Optical Absorption Spectroscopy (DOAS). FTIR examines the Fourier transform of a measured spectrum in order to determine what trace gases are interfering within the IR range of light. DOAS methods are based on light interference and absorption through air masses.

The DOAS technique takes advantage of the optically thin nature of HCHO in order to linearise the radiance differential through air masses with and without HCHO, using the Beer-Lambert intensity law. This method is used both in the home, and from space, globally for HCHO detection (Guenther et al., 1995; Gonzalez Abad et al., 2015; Davenport et al., 2015). As a trace gas HCHO interferes with light over a few wavelength bands, which allows instruments to detect concentrations between a known light source and a detector. Figure 1.6 shows the interference spectrum of HCHO

need to clean up language in here.

? You mean from ground?




FIGURE 1.6: HCHO spectrum, with a typical band of wavelengths used for DOAS path measurements. This is a portion of an image from Davenport et al., 2015.

On prior page you said
optically thin was an
advantage. Now a difficulty:

along with a typical band used to examine interference in the DOAS technique. One difficulty is that this interference is relatively small (HCHO is optically thin) and other compounds absorb light at similar wavelengths (Davenport et al., 2015).

FTIR and DOAS measurements have a range of uncertainties, including systematic and random measurement errors and uncertain apriori shape factors and water profiles (eg: Franco et al., 2015). Other types of measurement involve directly measuring the air, and determining chemical compounds through their physical properties. ??

how is this relevant
to HCHO measurement?

Vague

[/] A proton transfer reaction mass spectrometer (PTR-MS) can be used to determine gas phase evolution of terpene oxidation products (Lee et al., 2006; Nguyen et al., 2014; Wolfe et al., 2016, eg.). This is done through analysis of mass to charge ratios (m/z) of ionised air masses which are then identified as chemical compounds. Nguyen et al., 2014 use and compare several instruments (including one which is PTR-MS based) in the analysis of isoprene and monoterpene products. A Gas Chromatography mass spectrometer (GC-MS) is also able to identify isoprene, monoterpenes, and their products Lerner2017; Nguyen et al., 2014, eg.

Other measurement techniques include chromatographic and fluorimetric methods, both of which differ widely from each other and the spectroscopic methods (Hak et al., 2005). Hak et al., 2005 examine a single air mass with 8 instruments using the four techniques (MAX-DOAS, FTIR, chromatographic, and fluorimetric), and show that reasonable agreements can be achieved. Generally the measurements were somewhat close, the five Hantzsch instruments agreeing to within 11% (after removing two potentially faulty measurements), although different calibration standards were used. Titration for the different calibration solutions could not be resolved, which may account for absolute offsets up to 30%. These differences and non-uniformities between measurements (even among identical instruments) are part of the reason HCHO does not have a consistent network for global measurements like those for GHGs or Ozone (Chevallier et al., 2012). spell don't capitalise




FIGURE 1.7: An example spectrum showing interferences used for species concentration measurements by GOME-2. Image by EUMETSAT and ESA (EUMETSAT, 2015).

Satellite measurements

generic – you're talking just about HCHO, right?

Several satellites provide long term **trace gas** observations with near complete global coverage, including the ERS-2 launched in April 1995 which houses the GOME ultraviolet and visible (UV-Vis) spectrometer, the AURA launched in July 2004 which houses the OMI UV-Vis spectrometer, the MetOp-A and B launched in October 2006 and September 2012 respectively both housing a GOME-2 UV-Vis spectrometer. These satellites are on Low Earth Orbit (LEO) trajectories and overpass any area up to once per day. Satellites ~~can~~ use DOAS techniques with radiative transfer calculations on solar radiation absorption spectra to measure column HCHO. An example of a spectrum retrieved from the GOME-2 instrument is given in figure 1.7.

In conjunction with atmospheric chemistry and radiative models, satellite measurements can be used to quantify the abundance of **several chemical species in the atmosphere**. Isoprene is hard to measure directly due to its short lifetime and weak spectral absorption, instead HCHO is often used as a proxy (Millet et al., 2006; Fu et al., 2007; Dufour et al., 2008; Marais et al., 2012; Bauwens et al., 2013; Kefauver, Filella, and Peñuelas, 2014; Bauwens et al., 2016). This leads to a method of isoprene emissions estimation termed top-down (as opposed to bottom-up estimates). The existence of satellite data covering remote areas provides an opportunity to improve VOC emissions estimates leading to more robust models of global climate and chemistry. Satellite data allows us to **verify** large scale models of natural emissions, and their subsequent chemistry.

State in here that this Method is described in detail in Chapter xx.

again, stay focused. In your thesis, only used for HCHO/ isoprene.

Not necessarily verify, but provide alternative estimate

avoid using "chemical" - "chemical species" or "trace gas" generally ok.
(as a noun)

1.5 Atmospheric Chemistry Modelling

too informal

keep focused - much of
this is not how you
use models...

I think you are discussing
CTMs here, so state that.
Be specific.

Models can fill the gaps (both spatial and temporal) in measurement records, and can help us improve our understanding of the world around us. They are used ideally to steer us away from unsustainable pollution by showing us the future outcomes resulting from our emissions, from small to large scales. They can be used to increase measurement accuracy (for instance in satellite measurements) and determine where we lack information, while also checking the performance of new instruments. Precisely representing various chemicals and reactions in the atmosphere allows efficient mitigation of pollution, since we can compare scenarios against one another. Models can always be expanded to include new compounds or processes, however validation is always necessary. Currently they require improved isoprene emissions and subsequent chemistry understanding for effective air quality determination (Marvin et al., 2017).

Atmospheric chemical models provide a simulation of chemical densities and transport over time, through the atmosphere. They require many inputs (such as wind velocities) in order to accurately represent scenarios or regions on earth. Models of emissions are often used as drivers for atmospheric chemistry models, which require initial and boundary conditions in order to run. Chemistry in the atmosphere is a complex system of coupled reactions and dynamics, which can be solved using numerical partial differential equation solvers.

Chemical Transport Models (CTMs) simulate production, loss, and transport of chemical species. This is generally calculated using one or both of the Eulerian (box) or Lagrangian (puff) frames of reference. CTMs normally solve the continuity equations simultaneously with chemical production and loss for chemicals under inspection. The continuity equations describe transport of a conserved quantity such as mass, which, solved together with production and loss of a chemical can provide detailed simulations of natural processes.

? Solve for
many coupled
species, interesting
or not!

The general continuity equation links a quantity of a substance (q) to the field in which it flows and can be described by the formula:

$$\frac{\partial \rho}{\partial t} + \nabla \cdot j = \sigma$$

where ρ is density of q in the field, t is time, ∇ is divergence, j is the flux (q per unit area per unit time entering or leaving the field), and σ is the generation or loss of q per unit volume per unit time.

The type of model best suited to modelling the entire earth uses the Eulerian frame of reference, where the atmosphere is broken up into 3-D boxes with densities and transport calculated and stored for sequential steps in time at each location. The mass balance equation must be satisfied in any realistic long term box model and is as follows:

$$\begin{aligned}\frac{dm}{dt} &= \sum \text{sources} - \sum \text{sinks} \\ &= F_{in} + E + P - F_{out} - L - D\end{aligned}$$

where m is mass of a chemical, E and D are emission and deposition, P and L are




FIGURE 1.8: Standard box model parameters, image taken from Jacob, 1999.

production and loss, and F is chemical transport in and out, as shown in figure 1.8. Many chemical species interact with each other through production and loss. Any large chemical model will solve this mass balance equation over highly coupled arrays of partial differential equations, which becomes computation time expensive when looking at many species or resolved over many grid-boxes.

Contemporary models generally use mathematical differential solving tools of various complexity to solve chemical equations and reaction rates (often called chemical mechanisms) in order to predict an environment's evolution over time. Different solvers may be slower or faster and more suited to particular situations based on the stability of the equations and systems involved, and chemical mechanisms may vary in how many reactions and chemicals are listed and grouped together. For example: Since $[O] \ll [O_3]$ the chemical family Ox ($Ox \equiv O + O_3$) can be used to simplify chemistry simulations and approximate O_3 concentrations (Brasseur and Jacob, 2017, Chapter 3). Zhang et al., 2012 examine the outputs from a regional model (WRF/Chem) using three different chemical mechanisms, and they show some model output sensitivity to the choice of mechanism. relevance?

too informed
You aren't "solving reaction rates!"

families have more to do w/
lifetime against intrconversion
vs. terminal sinks.

1.5.1 Box models

Box models are much smaller scale than global CTMs, examining one uniform environment with many parametrisations such as transport and emissions. Box models can be used to check chemical mechanisms in specific scenarios, such as high or low NO_x environments. Marvin et al., 2017 use a box model matching conditions in southeast USA to evaluate isoprene mechanisms from several models. A box model involves modelling chemistry in a singular set of conditions without transport or any spatial gradients.

Should come before CTMs..

By allowing for interactions between boxes this concept can be extended to multiple-box models. These are simply multiple instances of single boxes with the addition of transport between them, which requires meteorological fields such as wind velocities and turbulence. The meteorology fields can be modelled, and/or input as parameters.

1.5.2 Emissions

There are two commonly used ways of estimating isoprene emissions, top-down or bottom-up. Bottom-up emission estimates generally model the flora and events which emit isoprene, like Eucalypts, factories, shrubs, etc. They use various properties of the emitters in order to estimate how much isoprene is being produced. Some of these properties include leaf areas, speciated responses to sunlight and temperature, moisture stress, etc (Guenther et al., 1995; Guenther et al., 2006). Understanding how much isoprene is emitted, when and by what, is complicated. One frequently used bottom up emissions model is the Model of Emissions of Gases and Aerosols from Nature (MEGAN, Guenther et al., 1995). Since little data exists with which to verify many of these bottom-up emission inventories, they can be uncertain on a large scale.

In many CTMs the isoprene emissions are calculated separately (for example by running MEGAN), and then used as boundary conditions (EG: Guenther et al., 2006). This can speed up calculations as the transport and concentrations can be simulated in various conditions without recalculating the emissions. Trace gases with short lifetimes and complex chemistry such as isoprene are often hard to measure which makes verifying model estimates difficult.

Bottom up models of VOC emissions are sensitive to parameters. For example Stavrakou et al., 2014 examined modelled Asian emissions and altered model parameters for temperature, plant type emission factors, incoming solar radiation (insolation) intensity, land use changes, and palm tree forest expansion. Changes were constrained by a network of radiation measurements and some experiments with south east Asian forest emissions - and led to reduction in isoprene emissions by a factor of two over the region. relative to what?

Marais et al., 2014 examine factors affecting isoprene emissions, showing their sensitivity to environmental factors. Their work used MEGAN (Guenther et al., 1995) and GEOS-Chem to look at how these factors affect surface ozone and particulate matter in Africa. One of the important uncertainties seen in MEGAN within this work is the isoprene emissions due to plant type. Canopy level isoprene measurements are made using relaxed eddy accumulation (REA) at several sites in Africa. One plant type near a measurement site emits more than other species and it's actual distribution on a larger scale is completely unknown - leading to possible overestimations in MEGAN. Current emissions estimates require more validation against observations, and recently a comparison of two major VOC models (MEGAN and ORCHIDEE) was undertaken by Messina et al., 2016 reiterating this requirement. In their work they examine model sensitivities and show that the important parameters are leaf area index (LAI), emission factors (EF), plant functional type (PFT), and light density fraction (LDF). There is high uncertainty in LAI and EF, which require more or improved measurements at the global scale. LDF parameterisation needs improvement and these models require more PFTs. Global emissions inventories like MEGAN often have large areas based on extrapolations which can introduce uncertainties (Miller et al., 2014).

1.5.3 Uncertainties?

Here I will attempt to list and partially explain the major uncertainties models have in relation to VOCs, SOAs, and ozone. TODO: Is this a good idea or should I put any

Expand bottom-up discussion and be more specific about what is actually included

is this relevant to your work?

Condense these paragraphs and link studies together more clearly. Make sure you have an answer to "what is the point of this paragraph? How is it relevant to my thesis?"

pertinent uncertainties with the associated work/descriptions? *The second one. Your intro should weave these into the narrative wherever possible.*

Emissions Inventories

Using different emissions inventories in an ACM can have large impacts on the simulation. Natural (biogenic or pyrogenic) and human driven (anthropogenic) emissions often drive a large fraction of atmospheric oxidation and radical chemistry, especially in the continental boundary layer. Zeng et al., 2015 examine the affects on CO and HCHO when running simulations with two different inventories. TODO: find where I took notes about Zeng2015 and put them here.

It is important to note that many estimates of isoprene emission are based on a few algorithms which can depend greatly on input parameters (Arneth et al., 2008; Niinemets et al., 2010). Arneth et al., 2008 argue that this monopoly of emissions estimates may be leading us to an incorrect understanding of isoprene chemistry. Yue, Unger, and Zheng, 2015 has shown that this is still a problem by looking at land carbon fluxes and modelling the sensitivity to VOC emissions estimates using two independent models of VOC emission. One model is photosynthesis based and estimates isoprene emissions using electron transfer energies and leaf physiology (Niinemets et al., 1999), while the other (MEGAN) uses the light and canopy temperature ((Guenther et al., 1995; Arneth et al., 2007) TODO: Read Arneth et al., 2007; Unger et al., 2013). Both are sensitive to light and temperature parameterisations.

Resolution - see also Yu 2016 paper

GEOS-Chem simulations are somewhat sensitive to the resolution *at which you run*. For example: Wild and Prather, 2006 show that reduced resolution increases OH concentrations and ozone production rates. Christian, Brune, and Mao, 2017 find small changes in OH (< 10%) in OH, HO₂ and ozone concentrations local to the north american arctic, when changing from 4 by 5 to 2 by 2.5 °resolution, *however they continue at lower resolution to save computational time.*

informal

irrelevant.

For many global scale analyses, errors from resolution are less important than those from chemistry, meteorology, and emissions (Christian, Brune, and Mao, 2017). Many models lack in-situ measurements with which to verify their chemical mechanisms, leading to large discrepancies, as seen in Marvin2017a TODO: briefly talk about Marvin2017a takeaways. Christian, Brune, and Mao, 2017 used GEOS-Chem v9-02 with 4° × 5° resolution, and while the low resolution adds errors in OH concentrations and O₃ production rates, the errors from chemistry, meteorology, and emissions are much larger.

Already stated important parts

Chemistry mechanisms

There is still much work to be done in models to correctly simulate the precursor emissions and processes which lead to HCHO. Often HCHO is used as a way of checking if these precursors are correctly modelled since HCHO measurements are more readily available (for instance from satellites). GEOS-Chem has recently been analysed for ozone and oxidant (OH and HO₂) sensitivity to the processes within the model along with inputs which drive it (Christian, Brune, and Mao, 2017). Christian, Brune, and

emissions, not chem mech.

and ozone??

Mao, 2017 found that GEOS-Chem ozone was most sensitive to NO₂ photolysis, the NO₂ + OH reaction rate, and various emissions.

Marvin et al., 2017 suggest that isoprene mechanisms in several contemporary models (including GEOS-Chem) are inadequate. They show that for a specific measurement campaign, the HCHO concentrations are underestimated in a way that can not be easily fixed through rate constant changes. Recently Marvin et al., 2017 compared five global ACMs isoprene mechanisms by evaluating simulated HCHO mixing ratios compared to in situ measurements from the Southeast Nexus (SENEX) aircraft campaign (in southeastern USA). They compared five models (GEOS-Chem, CB05, CB6r2, MCMv3.2, and MCMv3.3.1) and found all of them underestimated the HCHO concentrations (by 15 – 30%).

Another important factor in determining the yield of HCHO and other products from BVOCs is the local concentration of NO_x. Travis et al., 2016 show how modelled surface ozone is overestimated due to high estimates of NO_x emissions, which affect oxidative capacity and VOC reactions. Only for us...

Understanding of OH production/recycling in low NO conditions has been improved (see section 1.3.3), however many observations of OH were still quite underpredicted in models (Mao et al., 2012). Mao et al., 2012 showed how traditional OH measurements may be overestimated due to VOC oxidation. They looked at measurements in a remote forest in California and found that the instruments were generating OH internally. Nguyen et al., 2014 also see this VOC oxidation interference in measurements using a gas chromatographer (GC) with an equipped flame ionisation detector (FID). This lends more credence to the current understanding of VOC oxidation as it closed the gap between measurements and model predictions (Mao et al., 2012).

Clouds

One of the major uncertainties in chemical, climate, radiation, and weather models is cloud formation and dynamics. Clouds are remarkably complex at a much finer scale than can be accurately modelled by global chemistry models (with current processing power). Globally over half (50-60%) of the world is covered by clouds, with ~ 10% of them being rain-clouds (Kanakidou et al., 2005). Wet scavenging performed in clouds not only depends on large scale cloud processes, but also on the micro-physics of aerosols being scavenged, differing between aerosol sizes and hygroscopic properties.

Soil Moisture

Rowntree and Bolton, 1983 show how quickly soil moisture anomalies affect rainfall and other weather systems, while Chen and Dudhia, 2001 specifically show how important fine scale soil moisture information is when modelling land surface heat flux, and energy balances. Modelled emissions are sensitive to soil moisture, especially near the soil moisture threshold (or wilting point), below which trees stop emitting isoprene and other VOCs completely as they can no longer draw water (Bauwens et al., 2016). MEGAN accounts for soil moisture by applying it as an emission factor (EF) which scales the emission rate of various species.

Ozone or HCHO?

not really an uncertainty
in model chemical
mechanisms...

How is this relevant
to your thesis?

relevance to your thesis?

Droughts affects can be difficult to measure, as it is a multi-scale problem which affects various aspects of the land-air interface including plant emissions and dry deposition (Wang et al., 2017). The Standardised Precipitation Evapotranspiration Index (SPEI) is a measure of drought using TODO SPEI Drought Index. This product covers 1901 - 2011, and uses the average over that period as the background, in order to compare drought stressed regions against those with sufficient or excess water SPEI Drought Index.

not relevant!

1.6 Australia and the southern hemisphere

- much of this should go in VOC section. This should be an overview focused on stuff relevant to VOCs and O₃.

Australia has a unique climate, along with soil moisture, clay content and other important properties which affect VOC emissions. These properties are poorly understood in Australia due to the continents size and the relative sparsity of population centres, which make many areas very difficult or expensive to reach. In Australia most long term air quality or composition measurements are performed in or near large cities. Australia is dominated by areas with little anthropogenic influence and no ground based measurements of the natural emissions taking place (VanDerA et al., 2008). Due to the lack of in-situ ground based measurements, estimates of VOC emissions are uncertain, with large scale extrapolation required (Millet et al., 2006). Since many Australian cities are on the edge of regions with rich VOC emissions, it is very important to clarify the quantity, type, and cause of VOC emissions. Understanding of emissions from these areas is necessary to inform national policy on air pollution levels.

poorly understood or sparsely measured?

Can you definitively say that?

Vegetation The trees in Australia are diverse, a great summary is provided by ABARES using the national forest inventory at <http://www.agriculture.gov.au/abares/forestsaustralia/australias-forests>. Figure 1.9 shows the different forest types and their locations within Australia, highlighting that much of our forested lands are near population centres along the east coast. 16% of Australia is covered by forest, most (75%) of which is Eucalyptus.

informal

Fire emissions include a range of chemicals and each year the affects of fire or burning seasons blanket the northern and southern hemispheres independently. Biomass burning in southern Africa and South America has previously been shown to have a major influence on atmospheric composition in Australia (Oltmans et al., 2001; Gloudemanns et al., 2007; Edwards et al., 2006), particularly from July to December (Pak et al., 2003; Liu et al., 2016a). The ocean plays a role in VOC emissions as well. The Asian region is shown to have a strong correlation with the Oceanic Niño Index (ONI), with positive anomalies associated with El Niño (Stavrakou et al., 2014).

Why are you talking about Asia in section On Australia and SH?

Guenther et al., 2006 estimated that the Australian outback is among the worlds strongest isoprene emitters with forests in SE Australia having emission factors greater than $16 \text{ mg m}^{-2} \text{ h}^{-1}$ (see figure 1.10). Measurement campaigns in SE Australia have since cast doubt on the emission factors used by MEGAN, as the Eucalyptus trees and soil moisture were poorly studied Emmerson et al., 2016. These emissions factor estimates are not well verified and measurements of isoprene (or other BVOC) emissions barely cover Australia either spatially or temporally. However, comprehensive coverage of one high yield product (HCHO) in the atmosphere over Australia exists in the form of satellite measurements. Not the right place.

is that really what you mean?

are sparse and infrequent




FIGURE 1.9: Forest types in Australia <http://www.agriculture.gov.au/abares/forestsaustralia/australias-forests>




Fig. 2. Global distribution of landscape-average isoprene emission factors ($\text{mg isoprene m}^{-2} \text{ h}^{-1}$). Spatial variability at the base resolution ($\sim 1 \text{ km}$) is shown by regional images of the southeastern U.S. and southeastern Australia.

FIGURE 1.10: Part of a figure from Guenther et al., 2006 showing global isoprene emission factors.

1.6.1 Ozone

reference?

Ozone levels over Australia are relatively low, however it remains unclear how much we would expect this to change in the future as relatively little is known about precursors and influx for the continent. Australian air quality is monitored independently within each state, using several metrics. These metrics are measured by varying numbers of monitoring stations in each state. In New South Wales (NSW) the metrics used to determine air quality are: particulate matter (PM), O₃, CO, NO₂, SO₂, and visibility. An air quality index equal to the worst of these metrics is used for NSW as shown at <http://www.environment.nsw.gov.au/aqms/aqitable.htm>. Similar methods are used in other states to get an idea of air quality. Measurement stations are generally located in population centres, and don't regularly measure precursor emissions. This is an important omission as naturally emitted precursor gases often get transported into cities where they affect air quality through production of O₃ and other pollutants.

Generally STT over Australia affects the upper troposphere only, however ozone enhancements can reach quite low during heavy storms and cyclonic weather patterns (Alexander et al., 2013).

*Need to expand - remember ~1/3 - 1/2 your thesis is about O₃!
Air quality is not your focus, so don't force everything to centre on relevance to air quality.*

1.6.2 VOCs

Bottom up inventories of VOCs remain largely uncertain due to extensive extrapolation over plant functional types, changing land cover, and parameterised environmental stressors (Guenther et al., 2000; Kanakidou et al., 2005; Millet et al., 2006). VOC emission estimates are highly sensitive to many factors, several of which are not well characterised in Australia (Sindelarova et al., 2014; Bauwens et al., 2016). Changes in parameterisation of soil moisture in the MEGAN lead to massive changes in Australian isoprene emission estimates (Sindelarova et al., 2014). Over Australia MEGAN has problems involving unpublished plant functional types and their emissions, as well as poorly optimised soil moisture parameterisation (Emmerson et al., 2016).

Australia has the potential to be a major hotspot of isoprene emissions according to Guenther et al., 2006; Guenther et al., 2012, which shows heavy emissions factors in the region. Although recent work suggests that some Australian eucalypts may not be as egregious isoprene emitters as once thought Emmerson et al., 2016. Emissions in MEGAN are based on plant functional types, which can vary heavily even within species. TODO: more on Muller2008 Australia also lacks a clear estimate of emitted monoterpenes.

?

Emmerson et al., 2016 analyse EF sensitivity of a high resolution model of atmospheric chemistry over southeast Australia, comparing isoprene and monoterpene emissions against 4 separate campaigns. They show that the effect on total emissions is roughly linear and that no blanket EF changes are appropriate for all regions/seasons. They also mention that Australian eucalypt emissions are based on samples from young trees, which may emit more isoprene than older trees. Emmerson et al., 2016 suggest that monoterpenes may be emitted in similar quantities to isoprene, with more measurements required to determine if this is so. They compare emissions estimates from MEGAN against data from several field campaigns and see overestimated isoprene emissions, as well as underestimated monoterpene emissions.

Word choices...

*paragraph in a weird order.
distil to what is most relevant for your work
(i.e. not EF sensitivity)*

Their work suggests that MEGAN estimates of isoprene emissions may be 2-6 times too high, and monoterpene emissions \sim 3 times too low over southeast Australia.

This problem is even more pronounced in Australia due to poor characterisation, or because emission factors are based on northern hemispheric data. Many plant emissions rates have not been published, such as those for any Australian acacias. Additionally soil moisture is not well quantified which has a large effect on emissions. Müller et al., 2008 show how isoprene is poorly captured by the MEGAN model and analyse the affect of changing the soil moisture parameter, which can reduce the overall bias for Australia. Sindelarova et al., 2014 show reductions in modelled Australian isoprene emissions of 50% when incorporating soil moisture in MEGAN estimates. Uncertainties in isoprene emissions could explain why models of HCHO over Australia are poor at reproducing satellite measurements (Stavrakou et al., 2009).

Improvements to emissions models require improved understanding of regions and their behaviour. Inaccuracies can arise due to lack of data, such as the large and sparsely measured Australian outback. MEGAN has been shown to overpredict isoprene and underpredict monoterpene emissions in southeast Australia, with peaks and troughs captured but not at the right magnitude (Emmerson et al., 2016). MEGAN output in Australia is adversely affected by poor emission factor estimation. An example can be seen in Müller et al. (2008) where MEGAN overestimates isoprene in northern Australia. Underestimates of monoterpenes may be due simply to underestimated emission rates for many Eucalypt species (Winters et al., 2009).

1.6.3 Measurements

TODO: Brief overview of all the measurement campaigns, pointing to Modelling and Data chapter for more details. There are relatively few measurements of isoprene in the southern hemisphere, including MUMBA(TODO CITE), SPSS(cite), and that girl from Macquarie University with an instrument in the daintree rainforest(TODO CITE, DESCRIBE). For details on the MUMBA campaign see Section ???. An airflight campaign (HIPPO) measuring isoprene was also performed in 2009-2011? TODO: ask Jenny re this one.

A particulate and air quality measurement campaign took place in Sydney using PTR-MS and GC-FID, for details see Section ??.

One method of measuring ozone in the troposphere and stratosphere is by releasing weather balloons (with attached ozone detectors) which take readings as they rise up to around 30 km, giving a vertical profile of concentrations. Since 1986, Lauder, New Zealand (45° S, 170° E) has released ozonesondes allowing a multi-decadal analysis of ozone concentrations over the city (Brinksma et al., 2002). Kerguelan Island (49.2° S, 70.1° E), also has a record of ozonesonde profiles, which are directly in the path of biomass burning smoke plumes transported off shore from Africa (Baray et al., 2012). SHADOZ is the southern hemispheric additional ozone project, which have released sondes from 15 sites at different times <http://tropo.gsfc.nasa.gov/shadoz/>.

A smaller network of ozonesonde release sites is operated by TODO: get details for sondes I use in ozone chapter.

more pronounced in Australia than in SC Australia?

lots of redundancy in this paragraph.

+ Tumbarumba, see Emmerson paper

Start from overall aim to understand natural contributions to ozone over Australia/southern Ocean.
Then dive into specifics. 1 aim per chapter, same order as chapters.

1.7 Aims

TODO: outline of aims here (FIND THESE THEY ARE SOMEWHERE)

One of the aims in this thesis is to use the available satellite measurements to improve the estimates of isoprene emissions in Australia. Satellites which overpass daily record reflected solar (and emitted terrestrial) radiation, and give us measurements over all of Australia. Combining satellite data with model outcomes provides a platform for the understanding of natural processes which is especially useful over Australia. Due to the low availability of in-situ data over most of the Australian continent, a combination of the models with satellite can fill the gap of understanding of emissions from Australian landscapes. Improved emissions estimates will in turn improve the accuracy of CTMs, providing better predictions of atmospheric composition and its response to ongoing environmental change.

Calculation of isoprene to HCHO yields over Australia is required to create top-down estimates. This requires among other things an idea of which VOCs are present and their yields of HCHO. The technique of determining isoprene emissions from satellite detected HCHO is called satellite inversion. Another aim to this end is to ~~run and become familiar with~~ GEOS-Chem in order to determine Australian emissions and yields, and the importance of the relevant parameters. Soil moisture plays an important role in VOC emissions, as trees under stress may stop emitting various chemicals. This is especially true for Australia due to frequent droughts and wildfires. The argument for improved understanding of land surface properties, specifically soil moisture, is an old one(Mintz, 1982; Rountree and Bolton, 1983; Chen and Dudhia, 2001). But you don't include soil moisture!! Why draw attention to it here?

"use"? it is not an appropriate aim for a PhD to do something. Focus on science

To improve understanding of ozone over the southern hemisphere including Australia. Meteorology and precursor emissions are the largest drivers of tropospheric ozone concentrations, and an improved understanding of their effects in Australia would be facilitated by an analysis of STT as well as more confidence in the emitted precursors.

Bibliography

- Brasseur, Guy P and Daniel J Jacob (2017). *Modeling of Atmospheric Chemistry*. Cambridge University Press. DOI: [10.1017/9781316544754](https://doi.org/10.1017/9781316544754).
- Hsieh, Nan-Hung and Chung-Min Liao (2013). "Fluctuations in air pollution give risk warning signals of asthma hospitalization". In: *Atmospheric Environment* 75, pp. 206–216. DOI: [10.1016/j.atmosenv.2013.04.043](https://doi.org/10.1016/j.atmosenv.2013.04.043). URL: <http://dx.doi.org/10.1016/j.atmosenv.2013.04.043>.
- Avnery, Shiri et al. (2013). "Global crop yield reductions due to surface ozone exposure: 2. Year 2030 potential crop production losses and economic damage under two scenarios of O₃ pollution". In: *Atmospheric Environment* 71.13, pp. 408–409. ISSN: 13522310. DOI: [10.1016/j.atmosenv.2012.12.045](https://doi.org/10.1016/j.atmosenv.2012.12.045). URL: <http://dx.doi.org/10.1016/j.atmosenv.2011.01.002>.
- Yue, Xu et al. (2017). "Ozone and haze pollution weakens net primary productivity in China". In: *Atmospheric Chemistry and Physics* 17.9, pp. 6073–6089. ISSN: 1680-7324. DOI: [10.5194/acp-17-6073-2017](https://doi.org/10.5194/acp-17-6073-2017). URL: <https://www.atmos-chem-phys.net/17/6073/2017/>.
- Myhre, G and D Shindell (2013). *Chapter 8: Anthropogenic and Natural Radiative Forcing, in Climate Change 2013: The Physical Science Basis, Working Group 1 Contribution to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*, 2013. Fifth Assessment Report of the Intergovernmental Panel on Climate Change, 2013.
- Monks, P. S. et al. (2015). "Tropospheric ozone and its precursors from the urban to the global scale from air quality to short-lived climate forcer". In: *Atmospheric Chemistry and Physics* 15.15, pp. 8889–8973. ISSN: 1680-7324. DOI: [10.5194/acp-15-8889-2015](https://doi.org/10.5194/acp-15-8889-2015). URL: <https://www.atmos-chem-phys.net/15/8889/2015/>.
- Lelieveld, J. et al. (2013). "Model calculated global, regional and megacity premature mortality due to air pollution". In: *Atmospheric Chemistry and Physics* 13.14, pp. 7023–7037. ISSN: 16807324. DOI: [10.5194/acp-13-7023-2013](https://doi.org/10.5194/acp-13-7023-2013).
- Ayers, James D and William R Simpson (2006). "Measurements of N₂O₅ near Fairbanks, Alaska". In: *Journal of Geophysical Research: Atmospheres* 111.D14, n/a–n/a. ISSN: 2156-2202. DOI: [10.1029/2006JD007070](https://doi.org/10.1029/2006JD007070). URL: <http://dx.doi.org/10.1029/2006JD007070>.
- Lelieveld, J. et al. (2009). "Severe ozone air pollution in the Persian Gulf region". In: *Atmospheric Chemistry and Physics* 9, pp. 1393–1406. ISSN: 1680-7324. DOI: [10.5194/acp-9-1393-2009](https://doi.org/10.5194/acp-9-1393-2009).
- Ashmore, M R, Lisa. Emberson, and Murray Frank (2003). *Air pollution impacts on crops and forests : a global assessment*. Ed. by Lisa Emberson, Mike Ashmore, and Frank Murray. Imperial College Press London ; River Edge, NJ, xiii, 372 p. : ISBN: 186094292.

- Stevenson, D. S. et al. (2013). "Tropospheric ozone changes, radiative forcing and attribution to emissions in the Atmospheric Chemistry and Climate Model Intercomparison Project (ACCMIP)". In: *Atmospheric Chemistry and Physics* 13.6, pp. 3063–3085. ISSN: 16807316. DOI: [10.5194/acp-13-3063-2013](https://doi.org/10.5194/acp-13-3063-2013).
- Selin, N E et al. (2009). "Global health and economic impacts of future ozone pollution". In: *Environmental Research Letters* 4.4, p. 044014. ISSN: 1748-9326. DOI: [10.1088/1748-9326/4/4/044014](https://doi.org/10.1088/1748-9326/4/4/044014).
- Atkinson, Roger (2000). "Atmospheric chemistry of VOCs and NO(x)". In: *Atmospheric Environment* 34.12-14, pp. 2063–2101. ISSN: 13522310. DOI: [10.1016/S1352-2310\(99\)00460-4](https://doi.org/10.1016/S1352-2310(99)00460-4).
- Fuentes, J. D. et al. (2000). "Biogenic Hydrocarbons in the Atmospheric Boundary Layer: A Review". In: *Bulletin of the American Meteorological Society* 81.7, pp. 1537–1575. ISSN: 00030007. DOI: [10.1175/1520-0477\(2000\)081<1537:BHITAB>2.3.CO;2](https://doi.org/10.1175/1520-0477(2000)081<1537:BHITAB>2.3.CO;2). arXiv: [arXiv : 1011.1669v3](https://arxiv.org/abs/1011.1669v3). URL: [http://journals.ametsoc.org/doi/abs/10.1175/1520-0477\(2000\)081%3C1537%3ABHITAB%3E2.3.CO%3B2](http://journals.ametsoc.org/doi/abs/10.1175/1520-0477(2000)081%3C1537%3ABHITAB%3E2.3.CO%3B2).
- Paulot, Fabien et al. (2009b). "Unexpected Epoxide Formation in the". In: *Science* 325.2009, pp. 730–733. ISSN: 0036-8075. DOI: [10.1126/science.1172910](https://doi.org/10.1126/science.1172910).
- Atkinson, Roger and Janet Arey (2003). "Gas-phase tropospheric chemistry of biogenic volatile organic compounds: A review". In: *Atmospheric Environment* 37.SUPPL. 2. ISSN: 13522310. DOI: [10.1016/S1352-2310\(03\)00391-1](https://doi.org/10.1016/S1352-2310(03)00391-1).
- Jacob, Daniel J (1999). *Introduction to Atmospheric Chemistry*. Ed. by Daniel J Jacob. Princeton University Press. URL: <http://acmg.seas.harvard.edu/people/faculty/djj/book/index.html>.
- Huang, Guanyu et al. (2017). "Validation of 10-year SAO OMI Ozone Profile (PROFOZ) Product Using Aura MLS Measurements". In: *Atmospheric Measurement Techniques Discussions*, pp. 1–25. ISSN: 1867-8610. DOI: [10.5194/amt-2017-92](https://doi.org/10.5194/amt-2017-92). URL: <https://www.atmos-meas-tech-discuss.net/amt-2017-92/>.
- Cape, J. N. (2008). "Surface ozone concentrations and ecosystem health: Past trends and a guide to future projections". In: *Science of the Total Environment* 400.1-3, pp. 257–269. ISSN: 00489697. DOI: [10.1016/j.scitotenv.2008.06.025](https://doi.org/10.1016/j.scitotenv.2008.06.025). URL: [http://dx.doi.org/10.1016/j.scitotenv.2008.06.025](https://doi.org/10.1016/j.scitotenv.2008.06.025).
- Young, P J et al. (2017). "Tropospheric Ozone Assessment Report (TOAR): Assessment of global-scale model performance for global and regional ozone distributions, variability, and trends". In: *Elementa: Science of the Anthropocene*, pp. 0–84. ISSN: 2325-1026. DOI: [10.1525/elementa.265](https://doi.org/10.1525/elementa.265). URL: http://eprints.lancs.ac.uk/88836/1/TOAR_Model_Performance_07062017.pdfhttp://www.igacproject.org/sites/default/files/2017-05/TOAR-Model_Performance_draft_for_open_comment.pdf.
- Delmas, R, D Serca, and C Jambert (1997). "Global inventory of NOx sources". In: *Nutrient cycling in agroecosystems* 48.x, pp. 51–60. ISSN: 1385-1314. DOI: [10.1023/A:1009793806086](https://doi.org/10.1023/A:1009793806086). URL: [http://link.springer.com/article/10.1023/A:1009793806086](https://doi.org/10.1023/A:1009793806086).
- Sillman, Sanford (1999). "The relation between ozone , NO and hydrocarbons in urban and polluted rural environments". In: *Atmospheric Environment* 33. DOI: [https://doi.org/10.1016/S1352-2310\(98\)00345-8](https://doi.org/10.1016/S1352-2310(98)00345-8). URL: <http://www-personal>.

- umich . edu /\$ \sim \\$ sillman / web - publications / Sillmanreview99 . pdf https : //www.sciencedirect.com/science/article/pii/S1352231098003458.
- Stohl, Andreas et al. (2003). "A new perspective of stratosphere-troposphere exchange". In: *Bulletin of the American Meteorological Society* 84.11, pp. 1565–1573+1473. ISSN: 00030007. DOI: 10.1175/BAMS-84-11-1565.
- Kuang, Shi et al. (2017). "Summertime tropospheric ozone enhancement associated with a cold front passage due to stratosphere-to-troposphere transport and biomass burning: Simultaneous ground-based lidar and airborne measurements". In: *Journal of Geophysical Research: Atmospheres* 122.2, pp. 1293–1311. ISSN: 21698996. DOI: 10.1002/2016JD026078. URL: <http://doi.wiley.com/10.1002/2016JD026078>.
- Sprenger, Michael, Mischa Croci Maspoli, and Heini Wernli (2003). "Tropopause folds and cross-tropopause exchange: A global investigation based upon ECMWF analyses for the time period March 2000 to February 2001". In: *Journal of Geophysical Research* 108.D12. ISSN: 2156-2202. DOI: 10.1029/2002JD002587. URL: <http://dx.doi.org/10.1029/2002JD002587>.
- Ojha, Narendra et al. (2016). "Secondary ozone peaks in the troposphere over the Himalayas". In: *Atmospheric Chemistry and Physics Discussions* 17.November, pp. 1–25. ISSN: 1680-7375. DOI: 10.5194/acp-2016-908. URL: <http://www.atmos-chem-phys-discuss.net/acp-2016-908/>.
- Hegglin, Michaela I and Theodore G Shepherd (2009). "Large climate-induced changes in ultraviolet index and stratosphere-to-troposphere ozone flux". In: *Nature Geoscience* 2.10, pp. 687–691. DOI: 10.1038/ngeo604. URL: <http://dx.doi.org/10.1038/ngeo604>.
- Liu, Junhua et al. (2017). "Causes of interannual variability over the southern hemispheric tropospheric ozone maximum". In: *Atmos. Chem. Phys* 17.5, pp. 3279–3299. ISSN: 1680-7324. DOI: 10.5194/acp-17-3279-2017. URL: www.atmos-chem-phys.net/17/3279/2017/.
- Lin, Meiyun et al. (2015). "Climate variability modulates western US ozone air quality in spring via deep stratospheric intrusions." In: *Nature communications* 6.May, p. 7105. ISSN: 2041-1723. DOI: 10.1038/ncomms8105. URL: <http://www.nature.com/ncomms/2015/150512/ncomms8105/full/ncomms8105.html>.
- Young, P. J. et al. (2013). "Preindustrial to present-day changes in tropospheric hydroxyl radical and methane lifetime from the Atmospheric Chemistry and Climate Model Intercomparison Project (ACCMIP)". In: *Atmospheric Chemistry and Physics* 13.10, pp. 5277–5298. ISSN: 16807316. DOI: 10.5194/acp-13-5277-2013.
- Marvin, Margaret R. et al. (2017). "Impact of evolving isoprene mechanisms on simulated formaldehyde: An inter-comparison supported by in situ observations from SENEX". In: *Atmospheric Environment* 164, pp. 325–336. ISSN: 13522310. DOI: 10.1016/j.atmosenv.2017.05.049. URL: https://ac.els-cdn.com/S1352231017303618/1-s2.0-S1352231017303618-main.pdf?_tid=3de7eaaa-06ff-11e8-99a9-0000aacb360&acdnat=1517455576_09d7334af609ed43470155c1c42fad5fhttp://www.sciencedirect.com/science/article/pii/S1352231017303618.
- Mazzuca, Gina M. et al. (2016). "Ozone production and its sensitivity to NO_x and VOCs: Results from the DISCOVER-AQ field experiment, Houston 2013". In: *Atmospheric Chemistry and Physics* 16.22, pp. 14463–14474. ISSN: 16807324. DOI: 10.5194/acp-16-14463-2016.

- Stevenson, D S et al. (2006). "Multimodel ensemble simulations of present-day and near-future tropospheric ozone". In: *Journal of Geophysical Research* 111.D8. DOI: [10.1029/2005jd006338](https://doi.org/10.1029/2005jd006338). URL: <http://dx.doi.org/10.1029/2005JD006338>.
- Guenther, Alex et al. (2000). "Natural emissions of non-methane volatile organic compounds, carbon monoxide, and oxides of nitrogen from North America". In: *Atmospheric Environment* 34.12-14, pp. 2205–2230. ISSN: 13522310. DOI: [10.1016/S1352-2310\(99\)00465-3](https://doi.org/10.1016/S1352-2310(99)00465-3).
- Guenther, Alex et al. (1995). "A global model of natural volatile organic compound emissions". In: *Journal of Geophysical Research* 100.D5, pp. 8873–8892. ISSN: 0148-0227. DOI: [10.1029/94JD02950](https://doi.org/10.1029/94JD02950). URL: <http://onlinelibrary.wiley.com/doi/10.1029/94JD02950/full>.
- Glasius, Marianne and Allen H. Goldstein (2016). "Recent Discoveries and Future Challenges in Atmospheric Organic Chemistry". In: *Environmental Science and Technology* 50.6, pp. 2754–2764. ISSN: 15205851. DOI: [10.1021/acs.est.5b05105](https://doi.org/10.1021/acs.est.5b05105).
- Yue, X., N. Unger, and Y. Zheng (2015). "Distinguishing the drivers of trends in land carbon fluxes and plant volatile emissions over the past 3 decades". In: *Atmospheric Chemistry and Physics* 15.20, pp. 11931–11948. ISSN: 16807324. DOI: [10.5194/acp-15-11931-2015](https://doi.org/10.5194/acp-15-11931-2015).
- Stavrakou, T. et al. (2014). "Isoprene emissions over Asia 1979–2012: Impact of climate and land-use changes". In: *Atmospheric Chemistry and Physics* 14.9. ISSN: 16807324. DOI: [10.5194/acp-14-4587-2014](https://doi.org/10.5194/acp-14-4587-2014).
- Kwon, Hyeong-Ahn et al. (2017). "Sensitivity of formaldehyde (HCHO) column measurements from a geostationary satellite to temporal variation of the air mass factor in East Asia". In: *Atmospheric Chemistry and Physics* 17.7, pp. 4673–4686. ISSN: 1680-7324. DOI: [10.5194/acp-17-4673-2017](https://doi.org/10.5194/acp-17-4673-2017). URL: <http://www.atmos-chem-phys.net/17/4673/2017/>.
- Kanakidou, M et al. (2005). "Physics Organic aerosol and global climate modelling : a review". In: *Atmospheric Chemistry and Physics* 5, pp. 1053–1123.
- Aksoyoglu, Sebnem et al. (2017). "Secondary inorganic aerosols in Europe: sources and the significant influence of biogenic VOC emissions, especially on ammonium nitrate". In: *Atmospheric Chemistry and Physics* 17.12, pp. 7757–7773. ISSN: 1680-7324. DOI: [10.5194/acp-17-7757-2017](https://doi.org/10.5194/acp-17-7757-2017). URL: <https://www.atmos-chem-phys.net/17/7757/2017/>.
- Forster, P. et al. (2007). *Changes in Atmospheric Constituents and in Radiative Forcing*. In: *Climate Change 2007: The Physical Science Basis. Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change* [Solomon, S., D. Qin, M. Man. URL: https://www.ipcc.ch/publications_and_data/ar4/wg1/en/ch2.html (visited on 01/14/2016).
- Lelieveld, J et al. (2015). "The contribution of outdoor air pollution sources to premature mortality on a global scale". In: *Nature* 525.7569, pp. 367–371. DOI: [10.1038/nature15371](https://doi.org/10.1038/nature15371). URL: <http://dx.doi.org/10.1038/nature15371>.
- Hoek, Gerard et al. (2013). "Long-term air pollution exposure and cardio-respiratory mortality: a review". In: *Environmental Health* 12.1, p. 43. DOI: [10.1186/1476-069x-12-43](https://doi.org/10.1186/1476-069x-12-43). URL: <http://dx.doi.org/10.1186/1476-069x-12-43>.

- Krewski, D et al. (2009). "Extended follow-up and spatial analysis of the American Cancer Society study linking particulate air pollution and mortality". In: *Res Rep Health Eff Inst* 140, pp. 5–36. ISSN: 1041-5505 (Print) 1041-5505 (Linking). URL: <http://www.ncbi.nlm.nih.gov/pubmed/19627030>.
- Silva, Raquel A et al. (2013). "Global premature mortality due to anthropogenic outdoor air pollution and the contribution of past climate change". In: *Environ. Res. Lett.* 8.3, p. 34005. DOI: [10.1088/1748-9326/8/3/034005](https://doi.org/10.1088/1748-9326/8/3/034005). URL: <http://dx.doi.org/10.1088/1748-9326/8/3/034005>.
- Kroll, Jesse H. and John H. Seinfeld (2008). "Chemistry of secondary organic aerosol: Formation and evolution of low-volatility organics in the atmosphere". In: *Atmospheric Environment* 42.16, pp. 3593–3624. ISSN: 13522310. DOI: [10.1016/j.atmosenv.2008.01.003](https://doi.org/10.1016/j.atmosenv.2008.01.003). URL: <http://www.sciencedirect.com.ezproxy.uow.edu.au/science/article/pii/S1352231008000253>.
- Guenther, A et al. (2006). "Estimates of global terrestrial isoprene emissions using MEGAN (Model of Emissions of Gases and Aerosols from Nature)". In: *Atmospheric Chemistry and Physics* 6.11, pp. 3181–3210. DOI: [10.5194/acp-6-3181-2006](https://doi.org/10.5194/acp-6-3181-2006). URL: <http://dx.doi.org/10.5194/acp-6-3181-2006>.
- Millet, Dylan B et al. (2006). "Formaldehyde distribution over North America: Implications for satellite retrievals of formaldehyde columns and isoprene emission". In: *J. Geophys. Res.* 111.D24. DOI: [10.1029/2005jd006853](https://doi.org/10.1029/2005jd006853). URL: [TODO](#).
- Arneth, A et al. (2008). "Why are estimates of global terrestrial isoprene emissions so similar (and why is this not so for monoterpenes)?" In: *Atmos. Chem. Phys* 8.x, pp. 4605–4620. ISSN: 1680-7375. DOI: [10.5194/acpd-8-7017-2008](https://doi.org/10.5194/acpd-8-7017-2008).
- Niinemets, U. et al. (2010). "The emission factor of volatile isoprenoids: Stress, acclimation, and developmental responses". In: *Biogeosciences* 7.7, pp. 2203–2223. ISSN: 17264170. DOI: [10.5194/bg-7-2203-2010](https://doi.org/10.5194/bg-7-2203-2010).
- Müller, Jean-François (1992). "Geographical distribution and seasonal variation of surface emissions and deposition velocities of atmospheric trace gases". In: *Journal of Geophysical Research: Atmospheres* 97.D4, pp. 3787–3804. ISSN: 2156-2202. DOI: [10.1029/91JD02757](https://doi.org/10.1029/91JD02757). URL: <http://dx.doi.org/10.1029/91JD02757>.
- Kefauver, Shawn C., Iolanda Filella, and Josep Peñuelas (2014). "Remote sensing of atmospheric biogenic volatile organic compounds (BVOCs) via satellite-based formaldehyde vertical column assessments". en. In: *International Journal of Remote Sensing*. URL: <http://www.tandfonline.com/doi/abs/10.1080/01431161.2014.968690#.VkjEubNM61M>.
- Brown, S. S. et al. (2009). "Nocturnal isoprene oxidation over the Northeast United States in summer and its impact on reactive nitrogen partitioning and secondary organic aerosol". In: *Atmospheric Chemistry and Physics* 9.9, pp. 3027–3042. ISSN: 16807316. DOI: [10.5194/acp-9-3027-2009](https://doi.org/10.5194/acp-9-3027-2009).
- Guenther, A. B. et al. (2012). "The model of emissions of gases and aerosols from nature version 2.1 (MEGAN2.1): An extended and updated framework for modeling biogenic emissions". In: *Geoscientific Model Development* 5.6, pp. 1471–1492. ISSN: 1991959X. DOI: [10.5194/gmd-5-1471-2012](https://doi.org/10.5194/gmd-5-1471-2012).
- Messina, Palmira et al. (2016). "Global biogenic volatile organic compound emissions in the ORCHIDEE and MEGAN models and sensitivity to key parameters". In: *Atmospheric Chemistry and Physics* 16.22, pp. 14169–14202. ISSN: 16807324. DOI: [10.5194/acp-16-14169-2016](https://doi.org/10.5194/acp-16-14169-2016).

- 5194/acp-16-14169-2016. URL: <http://www.atmos-chem-phys.net/16/14169/2016/acp-16-14169-2016.pdf>.
- Wagner, V (2002). "Are CH₂O measurements in the marine boundary layer suitable for testing the current understanding of CH₄ photooxidation?: A model study". In: *Journal of Geophysical Research* 107.D3, p. 4029. ISSN: 0148-0227. DOI: [10.1029/2001JD000722](https://doi.wiley.com/10.1029/2001JD000722). URL: <http://doi.wiley.com/10.1029/2001JD000722>.
- Nguyen, T. B. et al. (2014). "Overview of the Focused Isoprene eXperiment at the California Institute of Technology (FIXCIT): Mechanistic chamber studies on the oxidation of biogenic compounds". In: *Atmospheric Chemistry and Physics* 14.24. ISSN: 16807324. DOI: [10.5194/acp-14-13531-2014](https://doi.org/10.5194/acp-14-13531-2014).
- Crounse, John D et al. (2012). "Atmospheric Fate of Methacrolein. 1. Peroxy Radical Isomerization Following Addition of OH and O₂". In: *Physical Chemistry m.*
- Wolfe, G. M. et al. (2016). "Formaldehyde production from isoprene oxidation across NO_x regimes". In: *Atmospheric Chemistry and Physics* 16.x, pp. 2597–2610. ISSN: 16807324. DOI: [10.5194/acp-16-2597-2016](https://doi.org/10.5194/acp-16-2597-2016). URL: www.atmos-chem-phys.net/16/2597/2016/.
- Liu, Yingjun et al. (2016b). "Isoprene photochemistry over the Amazon rainforest". In: *Proceedings of the National Academy of Sciences* 113.22, pp. 6125–6130. ISSN: 0027-8424. DOI: [10.1073/pnas.1524136113](https://doi.org/10.1073/pnas.1524136113). URL: <http://www.pnas.org/content/113/22/6125.abstract>.
- PATCHEN, AMIE K. et al. (2007). "Direct Kinetics Study of the Product-Forming Channels of the Reaction of Isoprene-Derived Hydroxyperoxy Radicals with NO". In: *International journal of Chemical Kinetics* 31.5, pp. 493–499. ISSN: 13000527. DOI: [10.1002/kin.20248/full](https://doi.org/10.1002/kin.20248). URL: <http://onlinelibrary.wiley.com/wol1/doi/10.1002/kin.20248/full>.
- Mao, Jingqiu et al. (2013). "Ozone and organic nitrates over the eastern United States: Sensitivity to isoprene chemistry". In: *Journal of Geophysical Research Atmospheres* 118.19, pp. 11256–11268. ISSN: 21698996. DOI: [10.1002/jgrd.50817](https://doi.org/10.1002/jgrd.50817).
- Paulot, F. et al. (2009a). "Isoprene photooxidation: new insights into the production of acids and organic nitrates". In: *Atmospheric Chemistry and Physics* 9.4, pp. 1479–1501. ISSN: 1680-7324. DOI: [10.5194/acp-9-1479-2009](https://doi.org/10.5194/acp-9-1479-2009).
- Crounse, John D. et al. (2013). "Autoxidation of organic compounds in the atmosphere". In: *Journal of Physical Chemistry Letters* 4.20, pp. 3513–3520. ISSN: 19487185. DOI: [10.1021/jz4019207](https://doi.org/10.1021/jz4019207). URL: <http://pubs.acs.org/doi/abs/10.1021/jz4019207>.
- Lin, Ying-Hsuan et al. (2013). "Epoxide as a precursor to secondary organic aerosol formation from isoprene photooxidation in the presence of nitrogen oxides". In: DOI: [10.1073/pnas.1221150110](https://doi.org/10.1073/pnas.1221150110). URL: <http://www.pnas.org/content/110/17/6718.full.pdf>.
- Peeters, Jozef and Jean-Francis Muller (2010). "HO_x radical regeneration in isoprene oxidation via peroxy radical isomerisations. II: experimental evidence and global impact". In: *Physical Chemistry Chemical Physics* 12.42, p. 14227. ISSN: 1463-9076. DOI: [10.1039/c0cp00811g](https://doi.org/10.1039/c0cp00811g). URL: <http://pubs.rsc.org/en/content/articlepdf/2010/cp/c0cp00811g>.

- Mao, J. et al. (2012). "Insights into hydroxyl measurements and atmospheric oxidation in a California forest". In: *Atmospheric Chemistry and Physics* 12.17, pp. 8009–8020. ISSN: 16807316. DOI: [10.5194/acp-12-8009-2012](https://doi.org/10.5194/acp-12-8009-2012).
- Surratt, Jason D et al. "Reactive intermediates revealed in secondary organic aerosol formation from isoprene". In: DOI: [10.1073/pnas.0911114107](https://doi.org/10.1073/pnas.0911114107). URL: <http://www.pnas.org/content/107/15/6640.full.pdf>.
- Nguyen, Tran B. et al. (2016). "Atmospheric fates of Criegee intermediates in the ozonolysis of isoprene". In: *Phys. Chem. Chem. Phys.* 18.15, pp. 10241–10254. ISSN: 1463-9076. DOI: [10.1039/C6CP00053C](https://doi.org/10.1039/C6CP00053C). URL: <http://xlink.rsc.org/?DOI=C6CP00053C>.
- Rollins, A W et al. (2009). "Isoprene oxidation by nitrate radical: alkyl nitrate and secondary organic aerosol yields". In: *Atmos. Chem. Phys. Atmospheric Chemistry and Physics* 9, pp. 6685–6703. URL: www.atmos-chem-phys.net/9/6685/2009/.
- Intergovernmental Panel on Climate Change (IPCC): *Climate Change: The Scientific Basis* (2001). Tech. rep. Cambridge University Press. URL: <http://www.ipcc.ch/ipccreports/tar/>.
- Stocker, T.F. et al. *IPCC, 2013: Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*. Tech. rep. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA. DOI: [10.1017/CBO9781107415324](https://doi.org/10.1017/CBO9781107415324).
- Davenport, J. J. et al. (2015). "A measurement strategy for non-dispersive ultra-violet detection of formaldehyde in indoor air : spectral analysis and interferent gases". In: *Measurement Science and Technology* 015802. December 2015, p. 15802. ISSN: 0957-0233. DOI: [10.1088/0957-0233/27/1/015802](https://doi.org/10.1088/0957-0233/27/1/015802). URL: <http://dx.doi.org/10.1088/0957-0233/27/1/015802>.
- Palmer, Paul I (2003). "Mapping isoprene emissions over North America using formaldehyde column observations from space". In: *J. Geophys. Res.* 108.D6. DOI: [10.1029/2002jd002153](https://doi.org/10.1029/2002jd002153). URL: <http://dx.doi.org/10.1029/2002jd002153>.
- Shim, Changsub et al. (2005). "Constraining global isoprene emissions with Global Ozone Monitoring Experiment (GOME) formaldehyde column measurements". In: *Journal of Geophysical Research Atmospheres* 110.24, pp. 1–14. ISSN: 01480227. DOI: [10.1029/2004JD005629](https://doi.org/10.1029/2004JD005629).
- Andreae, M O (2001). "Emission of trace gases and aerosols from biomass burning". In: *Biogeochemistry* 15.4, pp. 955–966. URL: <http://onlinelibrary.wiley.com/doi/10.1029/2000GB001382/epdf>.
- Millet, Dylan B. et al. (2008). "Spatial distribution of isoprene emissions from North America derived from formaldehyde column measurements by the OMI satellite sensor". In: *Journal of Geophysical Research Atmospheres* 113.2, pp. 1–18. ISSN: 01480227. DOI: [10.1029/2007JD008950](https://doi.org/10.1029/2007JD008950).
- Zhu, Lei et al. (2014). "Anthropogenic emissions of highly reactive volatile organic compounds in eastern Texas inferred from oversampling of satellite (OMI) measurements of HCHO columns". In: *Environmental Research Letters* 9.11, p. 114004. ISSN: 1748-9326. DOI: [10.1088/1748-9326/9/11/114004](https://doi.org/10.1088/1748-9326/9/11/114004). URL: <http://stacks.iop.org/1748-9326/9/i=11/a=114004?key=crossref.3d2869ee02fd4f0792f831ac8cbec117>.
- Fu, Tzung-may et al. (2007). "Space-based formaldehyde measurements as constraints on volatile organic compound emissions in east and south Asia and implications for ozone". In: 112, pp. 1–15. DOI: [10.1029/2006JD007853](https://doi.org/10.1029/2006JD007853).

- Emmerson, Kathryn M. et al. (2016). "Current estimates of biogenic emissions from eucalypts uncertain for southeast Australia". In: *Atmospheric Chemistry and Physics* 16.11, pp. 6997–7011. ISSN: 1680-7324. DOI: [10.5194/acp-16-6997-2016](https://doi.org/10.5194/acp-16-6997-2016). URL: <http://www.atmos-chem-phys.net/16/6997/2016/>.
- CRUTZEN, PAUL J, MARK G LAWRENCE, and ULRICH PÖSCHL (1999). "On the background photochemistry of tropospheric ozone". In: *Tellus A* 51.1, pp. 123–146. ISSN: 1600-0870. DOI: [10.1034/j.1600-0870.1999.t01-1-00010.x](https://doi.org/10.1034/j.1600-0870.1999.t01-1-00010.x). URL: <http://dx.doi.org/10.1034/j.1600-0870.1999.t01-1-00010.x>.
- Levy, Hiram (1972). "Photochemistry of the lower troposphere". In: *Planetary and Space Science* 20.6, pp. 919–935. ISSN: 00320633. DOI: [10.1016/0032-0633\(72\)90177-8](https://doi.org/10.1016/0032-0633(72)90177-8).
- Franco, B. et al. (2015). "Retrievals of formaldehyde from ground-based FTIR and MAX-DOAS observations at the Jungfraujoch station and comparisons with GEOS-Chem and IMAGES model simulations". In: *Atmospheric Measurement Techniques* 8.4, pp. 1733–1756. ISSN: 18678548. DOI: [10.5194/amt-8-1733-2015](https://doi.org/10.5194/amt-8-1733-2015).
- Gonzalez Abad, G. et al. (2015). "Updated Smithsonian Astrophysical Observatory Ozone Monitoring Instrument (SAO OMI) formaldehyde retrieval". In: *Atmospheric Measurement Techniques* 8.1, pp. 19–32. ISSN: 18678548. DOI: [10.5194/amt-8-19-2015](https://doi.org/10.5194/amt-8-19-2015).
- Lee, Anita et al. (2006). "Gas-phase products and secondary aerosol yields from the ozonolysis of ten different terpenes". In: 111.Ci, pp. 1–18. DOI: [10.1029/2005JD006437](https://doi.org/10.1029/2005JD006437).
- Hak, C. et al. (2005). "Intercomparison of four different in-situ techniques for ambient formaldehyde measurements in urban air". In: *Atmospheric Chemistry and Physics Discussions* 5.3, pp. 2897–2945. ISSN: 1680-7316. DOI: [10.5194/acpd-5-2897-2005](https://doi.org/10.5194/acpd-5-2897-2005).
- Chevallier, F. et al. (2012). "The formaldehyde budget as seen by a global-scale multi-constraint and multi-species inversion system". In: *Atmospheric Chemistry and Physics* 12.15, pp. 6699–6721. ISSN: 16807316. DOI: [10.5194/acp-12-6699-2012](https://doi.org/10.5194/acp-12-6699-2012). URL: <http://www.atmos-chem-phys.net/12/6699/2012/acp-12-6699-2012.pdf>.
- EUMETSAT (2015). GOME2. URL: <http://www.eumetsat.int/website/home/Satellites/CurrentSatellites/Metop/MetopDesign/GOME2/index.html>.
- Dufour, G. et al. (2008). "SCIAMACHY formaldehyde observations: constraint for isoprene emissions over Europe?" In: *Atmospheric Chemistry and Physics* 8.6, pp. 19273–19312. ISSN: 1680-7324. DOI: [10.5194/acpd-8-19273-2008](https://doi.org/10.5194/acpd-8-19273-2008).
- Marais, E A et al. (2012). "Isoprene emissions in Africa inferred from OMI observations of formaldehyde columns". In: *Atmospheric Chemistry and Physics* 12.3, pp. 7475–7520. DOI: [10.5194/acp-12-6219-2012](https://doi.org/10.5194/acp-12-6219-2012). URL: <http://dx.doi.org/10.5194/acp-12-6219-2012>.
- Bauwens, M et al. (2013). "Satellite-based isoprene emission estimates (2007–2012) from the GlobEmission project". In: *Proceedings of the ACCENT-Plus Symposium, Atmospheric Composition Change-Policy Support and Science, Urbino*, pp. 17–20.
- Bauwens, Maite et al. (2016). "Nine years of global hydrocarbon emissions based on source inversion of OMI formaldehyde observations". In: *Atmospheric Chemistry and Physics Discussions* March, pp. 1–45. ISSN: 1680-7375. DOI: [10.5194/acp-2016-221](https://doi.org/10.5194/acp-2016-221). URL: <http://www.atmos-chem-phys-discuss.net/acp-2016-221/>.

- Zhang, Yang et al. (2012). "Impact of gas-phase mechanisms on Weather Research Forecasting Model with Chemistry (WRF/Chem) predictions: Mechanism implementation and comparative evaluation". In: *Journal of Geophysical Research: Atmospheres* 117.D1, n/a-n/a. DOI: [10.1029/2011JD015775](https://doi.wiley.com/10.1029/2011JD015775). URL: <http://doi.wiley.com/10.1029/2011JD015775>.
- Marais, E A et al. (2014). "Improved model of isoprene emissions in Africa using Ozone Monitoring Instrument (OMI) satellite observations of formaldehyde: implications for oxidants and particulate matter". In: *Atmospheric Chemistry and Physics* 14.15, pp. 7693–7703. DOI: [10.5194/acp-14-7693-2014](https://doi.org/10.5194/acp-14-7693-2014). URL: <http://dx.doi.org/10.5194/acp-14-7693-2014>.
- Miller, C. et al. (2014). "Glyoxal retrieval from the Ozone Monitoring Instrument". In: *Atmospheric Measurement Techniques* 7.11, pp. 3891–3907. ISSN: 1867-8548. DOI: [10.5194/amt-7-3891-2014](https://doi.org/10.5194/amt-7-3891-2014). URL: <http://www.atmos-meas-tech.net/7/3891/2014/>.
- Zeng, G. et al. (2015). "Multi-model simulation of CO and HCHO in the Southern Hemisphere: comparison with observations and impact of biogenic emissions". In: *Atmospheric Chemistry and Physics* 15.13, pp. 7217–7245. ISSN: 1680-7324. DOI: [10.5194/acp-15-7217-2015](https://doi.org/10.5194/acp-15-7217-2015). URL: <http://www.atmos-chem-phys.net/15/7217/2015/>.
- Niinemets, U. et al. (1999). "A model of isoprene emission based on energetic requirements for isoprene synthesis and leaf photosynthetic properties for Liquidambar and Quercus". In: *Plant, Cell and Environment* 22.11, pp. 1319–1335. ISSN: 01407791. DOI: [10.1046/j.1365-3040.1999.00505.x](https://doi.org/10.1046/j.1365-3040.1999.00505.x).
- Arneth, Almut et al. (2007). "CO₂ inhibition of global terrestrial isoprene emissions: Potential implications for atmospheric chemistry". In: *Geophysical Research Letters* 34.18, p. L18813. DOI: [10.1029/2007GL030615](https://doi.wiley.com/10.1029/2007GL030615). URL: <http://doi.wiley.com/10.1029/2007GL030615>.
- Hewitt, C N et al. (2011). "Ground-level ozone influenced by circadian control of isoprene emissions". In: *Nature Geoscience* 4.10, pp. 671–674. DOI: [10.1038/ngeo1271](https://doi.org/10.1038/ngeo1271). URL: <http://dx.doi.org/10.1038/ngeo1271>.
- Fan, Jiwen and Renyi Zhang (2004). "Atmospheric oxidation mechanism of isoprene". In: *Environmental Chemistry* 1.3, pp. 140–149. ISSN: 14482517. DOI: [10.1071/EN04045](https://doi.org/10.1071/EN04045). URL: <http://dx.doi.org/10.1071/en04045>.
- Müller, J.-F. et al. (2008). "Global isoprene emissions estimated using MEGAN ECMWF analyses and a detailed canopy environment model". In: *Atmospheric Chemistry and Physics Discussions* 7.6, pp. 15373–15407. DOI: [10.5194/acpd-7-15373-2007](https://doi.org/10.5194/acpd-7-15373-2007). URL: <http://dx.doi.org/10.5194/acpd-7-15373-2007>.
- Winters, Anthony J et al. (2009). "Emissions of isoprene, monoterpene and short-chained carbonyl compounds from Eucalyptus spp. in southern Australia". In: *Atmospheric Environment* 43.19, pp. 3035–3043. ISSN: 13522310. DOI: [10.1016/j.atmosenv.2009.03.026](https://doi.org/10.1016/j.atmosenv.2009.03.026).
- Wild, Oliver and Michael J. Prather (2006). "Global tropospheric ozone modeling: Quantifying errors due to grid resolution". In: *Journal of Geophysical Research Atmospheres* 111.11, pp. 1–14. ISSN: 01480227. DOI: [10.1029/2005JD006605](https://doi.org/10.1029/2005JD006605).
- Christian, Kenneth E, William H Brune, and Jingqiu Mao (2017). "Global sensitivity analysis of the GEOS-Chem chemical transport model: ozone and hydrogen oxides

- during ARCTAS (2008)". In: *Atmos. Chem. Phys* 17, pp. 3769–3784. DOI: [10.5194/acp-17-3769-2017](https://doi.org/10.5194/acp-17-3769-2017). URL: www.atmos-chem-phys.net/17/3769/2017/.
- Travis, Katherine R et al. (2016). "Why do models overestimate surface ozone in the Southeast United States?" In: *Atmos. Chem. Phys* 16, pp. 13561–13577. DOI: [10.5194/acp-16-13561-2016](https://doi.org/10.5194/acp-16-13561-2016). URL: www.atmos-chem-phys.net/16/13561/2016/.
- Rowntree, P. R. and J. A. Bolton (1983). "Simulation of the atmospheric response to soil moisture anomalies over Europe". In: *Quarterly Journal of the Royal Meteorological Society* 109.461, pp. 501–526. ISSN: 00359009. DOI: [10.1002/qj.49710946105](https://doi.org/10.1002/qj.49710946105). URL: <http://doi.wiley.com/10.1002/qj.49710946105>.
- Chen, Fei and Jimy Dudhia (2001). "Coupling an Advanced Land Surface–Hydrology Model with the Penn State–NCAR MM5 Modeling System. Part I: Model Implementation and Sensitivity". In: *Monthly Weather Review* 129. URL: <https://journals.ametsoc.org/doi/pdf/10.1175/1520-0493%282001%29129%3C0569%3ACAALSH%3E2.0.C0%3B2>.
- Wang, Yuxuan et al. (2017). "Adverse effects of increasing drought on air quality via natural processes". In: *Atmos. Chem. Phys* 175194, pp. 12827–12843. DOI: [10.5194/acp-17-12827-2017](https://doi.org/10.5194/acp-17-12827-2017). URL: <https://www.atmos-chem-phys.net/17/12827/2017/acp-17-12827-2017.pdf>.
- SPEI Drought Index. URL: <http://spei.csic.es/home.html> (visited on 12/19/2017).
- VanDerA, R J et al. (2008). "Trends seasonal variability and dominant NO x source derived from a ten year record of NO 2 measured from space". In: *J. Geophys. Res.* 113.D4. DOI: [10.1029/2007jd009021](https://doi.org/10.1029/2007jd009021). URL: <http://dx.doi.org/10.1029/2007jd009021>.
- Oltmans, J et al. (2001). "Ozone in the Pacific tropical troposphere from ozonesonde observations". In: *Journal of Geophysical Research* 106.D23, pp. 32503–32525.
- Gloudemans, Annemieke et al. (2007). "Evidence for long-range transport of carbon monoxide in the Southern Hemisphere from SCIAMACHY observations". In: *European Space Agency, (Special Publication)* 33.SP-636, pp. 1–5. ISSN: 03796566. DOI: [10.1029/2006GL026804](https://doi.org/10.1029/2006GL026804).
- Edwards, D. P. et al. (2006). "Satellite-observed pollution from Southern Hemisphere biomass burning". In: *Journal of Geophysical Research* 111.14, pp. 1–17. ISSN: 01480227. DOI: [10.1029/2005JD006655](https://doi.org/10.1029/2005JD006655).
- Pak, B.C.a et al. (2003). "Measurements of biomass burning influences in the troposphere over southeast Australia during the SAFARI 2000 dry season campaign". In: *Journal of Geophysical Research* 108.13, pp. 1–10. ISSN: 0148-0227. DOI: [10.1029/2002JD002343](https://doi.org/10.1029/2002JD002343). URL: <http://www.scopus.com/inward/record.url?eid=2-s2.0-0742322536&partnerID=40&md5=cafaf3b948fb456696583ed3ab9a5>.
- Liu, Junhua et al. (2016a). "Causes of interannual variability of tropospheric ozone over the Southern Ocean". In: *Atmospheric Chemistry and Physics Discussions* October, pp. 1–46. ISSN: 1680-7316. DOI: [10.5194/ACP-2016-692](https://doi.org/10.5194/ACP-2016-692).
- Alexander, S. P. et al. (2013). "High resolution VHF radar measurements of tropopause structure and variability at Davis, Antarctica (69 S, 78 E)". In: *Atmospheric Chemistry and Physics* 13.6, pp. 3121–3132. ISSN: 16807324. DOI: [10.5194/acp-13-3121-2013](https://doi.org/10.5194/acp-13-3121-2013). URL: <http://www.atmos-chem-phys.net/13/3121/2013/>.
- Sindelarova, K. et al. (2014). "Global data set of biogenic VOC emissions calculated by the MEGAN model over the last 30 years". In: *Atmospheric Chemistry and Physics*

- 14.17, pp. 9317–9341. ISSN: 16807324. DOI: [10.5194/acp-14-9317-2014](https://doi.org/10.5194/acp-14-9317-2014). arXiv: [arXiv:1011.1669v3](https://arxiv.org/abs/1011.1669v3).
- Stavrakou, T et al. (2009). “Evaluating the performance of pyrogenic and biogenic emission inventories against one decade of space-based formaldehyde columns”. In: *Atmospheric Chemistry and Physics* 9.3, pp. 1037–1060. DOI: [10.5194/acp-9-1037-2009](https://doi.org/10.5194/acp-9-1037-2009). URL: <http://dx.doi.org/10.5194/acp-9-1037-2009>.
- Brinksma, E. J. et al. (2002). “Five years of observations of ozone profiles over Lauder, New Zealand”. In: *Journal of Geophysical Research* 107.D14, pp. 1–11. ISSN: 0148-0227. DOI: [10.1029/2001JD000737](https://doi.org/10.1029/2001JD000737). URL: <http://doi.wiley.com/10.1029/2001JD000737>.
- Baray, Jean-Luc et al. (2012). “One year ozonesonde measurements at Kerguelen Island (49.2S, 70.1E): Influence of stratosphere-to-troposphere exchange and long-range transport of biomass burning plumes”. In: *Journal of Geophysical Research* 117.D6. ISSN: 2156-2202. DOI: [10.1029/2011JD016717](https://doi.org/10.1029/2011JD016717). URL: <http://dx.doi.org/10.1029/2011JD016717>.
- Mintz, Y (1982). “The sensitivity of numerically simulated climates to land surface conditions”. In: *Land Surface Processes in Atmospheric General Circulation Models*, pp. 109–111.
- Chen, D. et al. (2009). “Regional CO pollution in China simulated by the high-resolution nested-grid GEOS-Chem model”. In: *Atmospheric Chemistry and Physics Discussions* 9.2, pp. 5853–5887. ISSN: 1680-7324. DOI: [10.5194/acpd-9-5853-2009](https://doi.org/10.5194/acpd-9-5853-2009). URL: <http://dx.doi.org/10.5194/acpd-9-3825-2009>.
- Giglio, Louis, James T. Randerson, and Guido R. Van Der Werf (2013). “Analysis of daily, monthly, and annual burned area using the fourth-generation global fire emissions database (GFED4)”. In: *Journal of Geophysical Research* 118.1, pp. 317–328. ISSN: 21698961. DOI: [10.1002/jgrg.20042](https://doi.org/10.1002/jgrg.20042).
- Horowitz, Larry W. et al. (1998). “Export of reactive nitrogen from North America during summertime: Sensitivity to hydrocarbon chemistry”. In: *Journal of Geophysical Research* 103.D11, pp. 13451–13476. ISSN: 0148-0227. DOI: [10.1029/97JD03142](https://doi.org/10.1029/97JD03142). URL: <http://doi.wiley.com/10.1029/97JD03142> <http://www.agu.org/pubs/crossref/1998/97JD03142.shtml>.
- Crounse, John D et al. (2011). “Peroxy radical isomerization in the oxidation of isoprene”. In: *Physical Chemistry Chemical Physics* 13.30, pp. 13607–13613. ISSN: 1463-9076. DOI: [doi:10.1039/c1cp21330j](https://doi.org/10.1039/c1cp21330j). URL: <http://dx.doi.org/10.1039/C1CP21330J>.
- Jozef, Peeters et al. (2014). “Hydroxyl Radical Recycling in Isoprene Oxidation Driven by Hydrogen Bonding and Hydrogen Tunneling: The Upgraded LIM1 Mechanism”. In: *Journal of Physical Chemistry*.
- Guenther, Alex (2016). MEGAN. URL: <http://lar.wsu.edu/megan/>.
- Pegoraro, E. et al. (2004). “Effect of drought on isoprene emission rates from leaves of *Quercus virginiana* Mill.” In: *Atmospheric Environment* 38.36, pp. 6149–6156. ISSN: 13522310. DOI: [10.1016/j.atmosenv.2004.07.028](https://doi.org/10.1016/j.atmosenv.2004.07.028). URL: <http://linkinghub.elsevier.com/retrieve/pii/S1352231004007198>.
- Dunne, Erin et al. (2017). “Comparison of VOC measurements made by PTR-MS, Adsorbent Tube/GC-FID-MS and DNPH-derivatization/HPLC during the Sydney Particle Study, 2012: a contribution to the assessment of uncertainty in current

- atmospheric VOC measurements". In: *Atmospheric Measurement Techniques Discussions*, pp. 1–24. ISSN: 1867-8610. DOI: [10.5194/amt-2016-349](https://doi.org/10.5194/amt-2016-349). URL: <https://www.atmos-meas-tech-discuss.net/amt-2016-349/>.
- Lee, Hanlim et al. (2015). "Investigations of the Diurnal Variation of Vertical HCHO Profiles Based on MAX-DOAS Measurements in Beijing: Comparisons with OMI Vertical Column Data". In: *Atmosphere*. URL: [10.3390/atmos6111816](https://doi.org/10.3390/atmos6111816).
- Schreier, Stefan F. et al. (2016). "Estimates of free-Tropospheric NO₂ and HCHO mixing ratios derived from high-Altitude mountain MAX-DOAS observations at mid-latitudes and in the tropics". In: *Atmospheric Chemistry and Physics* 16.5. ISSN: 16807324. DOI: [10.5194/acp-16-2803-2016](https://doi.org/10.5194/acp-16-2803-2016).
- Leue, C et al. (2001). "Quantitative analysis of NO x emissions from Global Ozone Monitoring Experiment satellite image sequences". In: *J. Geophys. Res.* 106.D6, p. 5493. DOI: [10.1029/2000jd900572](https://doi.org/10.1029/2000jd900572). URL: <http://dx.doi.org/10.1029/2000jd900572>.
- Stavrakou, T. et al. (2015). "How consistent are top-down hydrocarbon emissions based on formaldehyde observations from GOME-2 and OMI?" English. In: *Atmospheric Chemistry and Physics* 15.20, pp. 11861–11884. ISSN: 1680-7324. DOI: [10.5194/acp-15-11861-2015](https://doi.org/10.5194/acp-15-11861-2015). URL: <http://www.atmos-chem-phys.net/15/11861/2015/acp-15-11861-2015.html>.
- Schenkeveld, V. M. Erik et al. (2017). "In-flight performance of the Ozone Monitoring Instrument". In: *Atmospheric Measurement Techniques* 10.5, pp. 1957–1986. ISSN: 1867-8548. DOI: [10.5194/amt-10-1957-2017](https://doi.org/10.5194/amt-10-1957-2017). URL: <http://www.atmos-meas-tech-discuss.net/amt-2016-420/> <https://www.atmos-meas-tech.net/10/1957/2017/>.
- Palmer, Paul I et al. (2001). "Air mass factor formulation for spectroscopic measurements from satellites' Application to formaldehyde retrievals from the Global Ozone Monitoring Experiment". In: *Journal of Geophysical Research* 106.D13.
- Lorente, Alba et al. (2017). "Structural uncertainty in air mass factor calculation for NO₂ and HCHO satellite retrievals". In: *Atmospheric Measurement Techniques* 2, pp. 1–35. ISSN: 1867-8610. DOI: [10.5194/amt-10-759-2017](https://doi.org/10.5194/amt-10-759-2017). URL: <https://www.atmos-meas-tech.net/10/759/2017/amt-10-759-2017.html> <http://www.atmos-meas-tech-discuss.net/amt-2016-306/>.
- Eskes, H J and K F Boersma (2003). "Averaging kernels for DOAS total-column satellite retrievals". In: *Atmospheric Chemistry and Physics* 3.1, pp. 1285–1291. ISSN: 1680-7324. DOI: [10.5194/acpd-3-895-2003](https://doi.org/10.5194/acpd-3-895-2003). URL: <http://dx.doi.org/10.5194/acpd-3-895-2003>.
- Lamsal, L N et al. (2014). "Evaluation of OMI operational standard NO₂ column retrievals using in situ and surface-based NO₂ observations". In: *Atmos. Chem. Phys* 14, pp. 11587–11609. DOI: [10.5194/acp-14-11587-2014](https://doi.org/10.5194/acp-14-11587-2014). URL: www.atmos-chem-phys.net/14/11587/2014/.
- Vasilkov, A et al. (2017). "Accounting for the effects of surface BRDF on satellite cloud and trace-gas retrievals: a new approach based on geometry-dependent Lambertian equivalent reflectivity applied to OMI algorithms". In: *Atmospheric Measurement Techniques* 10.1, pp. 333–349. DOI: [10.5194/amt-10-333-2017](https://doi.org/10.5194/amt-10-333-2017). URL: <http://www.atmos-meas-tech.net/10/333/2017/>.
- Vigouroux, C. et al. (2009). "Ground-based FTIR and MAX-DOAS observations of formaldehyde at Réunion Island and comparisons with satellite and model data".

- In: *Atmospheric Chemistry and Physics Discussions* 9, pp. 15891–15957. ISSN: 1680-7316. DOI: [10.5194/acpd-9-15891-2009](https://doi.org/10.5194/acpd-9-15891-2009).
- Instrument, O M I (2002). "OMI Algorithm Theoretical Basis Document Volume I". In: I.August, pp. 1–50.
- Zhu, Lei et al. (2016). "Observing atmospheric formaldehyde (HCHO) from space: validation and intercomparison of six retrievals from four satellites (OMI, GOME2A, GOME2B, OMPS) with SEAC4RS aircraft observations over the Southeast US". In: *Atmospheric Chemistry and Physics* 0, pp. 1–24. ISSN: 1680-7375. DOI: [10.5194/acp-2016-162](https://doi.org/10.5194/acp-2016-162). URL: <http://www.atmos-chem-phys.net/16/13477/2016/acp-16-13477-2016.pdf>.
- De Smedt, I. et al. (2015). "Diurnal, seasonal and long-term variations of global formaldehyde columns inferred from combined OMI and GOME-2 observations". In: *Atmospheric Chemistry and Physics* 15.21, pp. 12519–12545. ISSN: 16807324. DOI: [10.5194/acp-15-12519-2015](https://doi.org/10.5194/acp-15-12519-2015). URL: <http://www.atmos-chem-phys-discuss.net/15/12241/2015/5Cnhttp://www.atmos-chem-phys-discuss.net/15/12241/2015/acpd-15-12241-2015.pdf>.
- Barkley, Michael P. et al. (2013). "Top-down isoprene emissions over tropical South America inferred from SCIAMACHY and OMI formaldehyde columns". In: *Journal of Geophysical Research Atmospheres* 118.12, pp. 6849–6868. ISSN: 21698996. DOI: [10.1002/jgrd.50552](https://doi.org/10.1002/jgrd.50552). URL: <http://dx.doi.org/10.1002/jgrd.50552>.
- Abad, Gonzalo González et al. (2016). "Smithsonian Astrophysical Observatory Ozone Mapping and Profiler Suite (SAO OMPS) formaldehyde retrieval". In: *Atmospheric Measurement Techniques* 9.7, pp. 2797–2812. ISSN: 18678548. DOI: [10.5194/amt-9-2797-2016](https://doi.org/10.5194/amt-9-2797-2016).
- De Smedt, I. et al. (2012). "Improved retrieval of global tropospheric formaldehyde columns from GOME-2/MetOp-A addressing noise reduction and instrumental degradation issues". In: *Atmospheric Measurement Techniques* 5.11, pp. 2933–2949. ISSN: 18671381. DOI: [10.5194/amt-5-2933-2012](https://doi.org/10.5194/amt-5-2933-2012).
- Chance, K. et al. (2000). "Satellite observations of formaldehyde over North America from GOME". In: *Geophysical Research Letters* 27.21, pp. 3461–3464. ISSN: 00948276. DOI: [10.1029/2000GL011857](https://doi.org/10.1029/2000GL011857). URL: <http://dx.doi.org/10.1029/2000gl011857>.
- Zhu, Lei et al. (2013). "Variability of HCHO over the Southeastern United States observed from space : Implications for VOC emissions". In: vol. 1.