

## Variable Research

### QN43

The variable QN43 aims to observe whether students are binge drinkers. It defines a binge drinker as someone who has had four or more drinks of alcohol in a row if they were female or five or more drinks of alcohol in a row if they were male, within a couple of hours, on at least 1 day during the 30 days before the survey. Studies on drinking young adults indicate that drinking can cause emotional regulation which would have an impact on their suicidal ideation. Further studies have found adolescent who binge drinkers are 12.6x more likely to idolize suicide indicating a significant variable to consider when measuring cooccurring factors.

<https://pmc.ncbi.nlm.nih.gov/articles/PMC2607328/>

<https://www.sciencedirect.com/science/article/pii/S0165032722013477>

### QN48

The variable QN48 asks whether a student in the past 30 days, has used marijuana. Research on the significance of marijuana usage on ideations on suicide is rather mixed. With some papers citing marijuana as a significant factor, increasing by 50-65% in adolescence. Others cite marijuana as a significant factor only in heavy users of the substance. Thus, identifying it's significant in this survey would be useful to understand it's impact on adolescence.

<https://www.ncbi.nlm.nih.gov/books/NBK425748/>

<https://pubmed.ncbi.nlm.nih.gov/41145103/>

### QNTB4

QNTB4 measures whether currently smoked cigarettes or cigars or used smokeless tobacco or electronic vapor products. Studies have investigated the association between smoking status and found that smoking increased risk of suicidal ideation across age-groups generally. Within adolescent populations, evidence shows a positive direct relationship between cigarette smoking with suicidal attempts when modelling and assessing survey data in 28 countries. With publication backing this correlation, we've decided to look further into the relationship of adolescent suicide to see if the YRBS reflects these findings.

<https://pmc.ncbi.nlm.nih.gov/articles/PMC10082789/>

[A moderated mediation analysis of the association between smoking and suicide attempts among adolescents in 28 countries - PMC](#)

**QNOWT, QNOBESE**

Public research a person's BMI, particularly whether they are overweight and being obese showcases interesting. In adults, it has been used to show that it increased the likelihood of suicidal ideation, particularly in female professionals under the age 35. However, aren't significant over the age of 35. However, studies into young adults' weight and suicidal ideation show a strong correlation, particularly with young men.

<https://pubmed.ncbi.nlm.nih.gov/29957479/>

<https://pubmed.ncbi.nlm.nih.gov/41665369/>

<https://pmc.ncbi.nlm.nih.gov/articles/PMC9079098/>

**Bernard Kyaluzi**