

Top markers by original cluster
Top 5 per group: Bonferroni-adjusted P < 0.05; detection difference >= 10 percentage points
0: T
1: Epithelial
2: Macrophage
3: T/NK cells
4: Macrophage
5: Myeloid APCs
6: Epithelial
7: B
8: Plasma
9: Endothelial
10: Mast
11: Epithelial
12: Epithelial
13: Fibroblast
14: Pericyte
15: T
ICOS
CAMK4
BCL11B
IL7R
BICDL1
ZNF385B
PKHD1
GPC5
ABCA4
ACOXL
LYZ
APOC1
APOE
CCL18
SPP1
GNLY
NKG7
GZMB
GZMH
GZMA
RETN
MCEMP1
AQP9
MARCO
MMP19
CD1E
S100B
CD1C
HLA-DQB2
PADI2
SFTA2
S100A14
WFDC2
MDK
KRT18
MS4A1
VPREB3
PAX5
LINC00926
LINC01781
IGLC2
IGLC3
MZB1
ANKRD36BP2
IGHG4
EMCN
CCL14
CLEC14A
CYYR1
ACKR1
TPSB2
MS4A2
TPSAB1
CTSG
CPA3
LDLRAD1
CFAP144
HOATZ
C20orf85
MROH9
SERPINB3
BPIFB1
SCGB1A1
CXCL6
LCN2
MFAP4
ADH1B
FBLN1
SFRP2
ABCA8
FAM162B
COX4I2
HIGD1B
RGS5
EGFL6
MKI67
ASPM
H1-5
TOP2A
CENPF
Detected cells (%)
10
25
50
75
100
Relative mean RNA expression
2
1
0
-1
-2
All retained cells used. Size: raw count > 0; color: gene-wise z score of log1p(mean normalized expression).