**≮**Back to Week 2

**XLessons** 

Prev

Next

## Multiple Features

**Note:** [7:25 -  $\theta^T$  is a 1 by (n+1) matrix and not an (n+1) by 1 matrix]

Linear regression with multiple variables is also known as "multivariate linear regression".

We now introduce notation for equations where we can have any number of input variables.

 $x_i^{(i)} = \text{value of feature } j \text{ in the } i^{th} \text{ training example}$ 

 $x^{(i)} =$ the column vector of all the feature inputs of the  $i^{th}$  training example

m =the number of training examples

 $n = \left| x^{(i)} \right|; ext{(the number of features)}$ 

The multivariable form of the hypothesis function accommodating these multiple features is as follows:

$$h_{\theta}(x) = \theta_0 + \theta_1 x_1 + \theta_2 x_2 + \theta_3 x_3 + \dots + \theta_n x_n$$

In order to develop intuition about this function, we can think about  $\theta_0$  as the basic price of a house,  $\theta_1$  as the price per square meter,  $\theta_2$  as the price per floor, etc.  $x_1$  will be the number of square meters in the house,  $x_2$  the number of floors, etc.

Using the definition of matrix multiplication, our multivariable hypothesis function can be concisely represented as:

$$h_{ heta}(x) = \left[egin{array}{cccc} heta_0 & & heta_1 & & \dots & & heta_n 
ight] egin{bmatrix} x_0 \ x_1 \ dots \ x_n \end{bmatrix} = heta^T x$$

This is a vectorization of our hypothesis function for one training example; see the lessons on vectorization to learn more.

Remark: Note that for convenience reasons in this course we assume  $x_0^{(i)}=1$  for  $(i\in 1,\ldots,m)$ . This allows us to do matrix operations with theta and x. Hence making the two vectors ' $\theta$ ' and  $x^{(i)}$  match each other element-wise (that is, have the same number of elements: n+1).]

The following example shows us the reason behind setting  $x_0^{(i)}=1$  :

$$X = egin{bmatrix} x_0^{(1)} & x_0^{(2)} & x_0^{(3)} \ x_1^{(1)} & x_1^{(2)} & x_1^{(3)} \end{bmatrix} \; , heta = egin{bmatrix} heta_0 \ heta_1 \end{bmatrix}$$

As a result, you can calculate the hypothesis as a vector with:

$$h_{ heta}(X) = heta^T X$$

Mark as completed





