POTS Mission - Complete Report

Final Report and research files 01-17 by the Hermes Agent

POTS Mission — Hermes Agent research

PDF prepared 24 October 2026

Contents

POTS MISSION - FINAL REPORT
1. What post covid pots is
2. Master list remedies
3. Fluids salt electrolytes
4. Diet meals trigger foods
5. Compression physical methods
6. Exercise reconditioning
7. Vitamins minerals supplements
8. Sleep heat daily habits
9. Long covid specific research
10. Traditional medicine worldwide
11. Ongoing trials
12. Completed unpublished trials
13. Abandoned stopped trials
14. Recovery case reports
15. Otc products proven vs hype
16. Combination approaches
17. Own leads

POTS MISSION — FINAL REPORT

Natural/OTC help for post-COVID POTS. For a 32-year-old woman, ~6 months after COVID. Built from files 01-17. Every rating is honest; every link is real. Nothing here is a proven cure — POTS after COVID usually improves with months of consistent daily work.

TOP 10 NATURAL/OTC STEPS, RANKED BY PROOF

1. STRUCTURED RECUMBENT EXERCISE RECONDITIONING — MODERATE (multiple good trials). Recumbent bike/rowing/swimming first, start 10-20 min/day, build to 40-45 min, 4-6 days/week over 3 months; 71% of completers no longer met POTS criteria. https://pubmed.ncbi.nlm.nih.gov/33482385/

2. FLUIDS + SALT TOGETHER — MODERATE. 2-3 L fluid + 8-12 g salt/day (doctor's OK; never with kidney disease, heart failure or uncontrolled high BP). Works in days to weeks. https://www.sciencedirect.com/science/article/pii/S1566070222000108

3. ABDOMINAL/WAIST-HIGH COMPRESSION — MODERATE. 20-40 mmHg, abdomen included (beats socks-only), put on lying down before getting up. Works immediately. https://www.jacc.org/doi/10.1016/j.jacc.2020.11.040

4. ORAL REHYDRATION SOLUTION (ORS) — MODERATE. 1-2 packets/day (~2,000 mg sodium each) in ~500 mL water; matched IV saline in a controlled trial. Works in 30-60 min. https://www.jpeds.com/article/S0022-3476(19)30893-5/fulltext

5. ACUTE WATER BOLUS — MODERATE. ~500 mL plain water, drunk fairly fast, 20-30 min before standing/showering; blunts the heart-rate rise for ~60 min. Take salt separately, not in the bolus. https://www.ahajournals.org/doi/full/10.1161/01.cir.0000038921.64575.d0

6. MUSCLE COUNTER-MANEUVERS — MODERATE (lab/tilt evidence, no large trials). At first dizziness: leg-cross + tense, squat, or hard handgrip; BP rises in seconds. https://pmc.ncbi.nlm.nih.gov/articles/PMC9606335/

7. SMALL FREQUENT MEALS, LIMIT REFINED CARBS AND ALCOHOL — MODERATE. 4-6 meals of 300-500 kcal; avoid big sugar loads before standing; alcohol near-zero (it doubles the BP fall in controlled studies). https://www.ahajournals.org/doi/10.1161/HYPERTENSIONAHA.121.17852

8. EAR-CLIP VAGUS NERVE STIMULATION (tVNS) — MODERATE (one small blinded RCT). 1 hr/day for 2 months lowered standing HR and inflammatory markers; OTC ear-clip devices ~$50-150. https://pmc.ncbi.nlm.nih.gov/articles/PMC10932945/

9. IRON, ONLY IF FERRITIN IS LOW — EARLY. Get tested; if ferritin <~50 ng/mL, ferrous sulfate 325 mg every other day; retest in 8-12 weeks. Never without a confirmed deficiency. https://clinicaltrials.gov/study/NCT07197905

10. NICOTINAMIDE RIBOSIDE (NAD+) 2,000 mg/day — EARLY. In a long-COVID RCT it took 10+ weeks to show fatigue/sleep benefits. Expensive, unproven for POTS itself. https://www.thelancet.com/journals/eclinm/article/PIIS2589-5370(25)00567-X/fulltext

Also honest: melatonin 0.5-3 mg for sleep (EARLY), 4-week gluten-free trial (EARLY,

small study), head-of-bed raised 10-15 cm (EARLY). CoQ10 FAILED its long-COVID trial (https://www.thelancet.com/journals/lanepe/article/PIIS2666-7762(22)00235-6/fulltext). Loratadine 10 mg twice daily put ONE documented MCAS-overlap case (hives with episodes) into full remission — REPORTED, one person, do not copy without a doctor: https://europepmc.org/article/MED/38526146

BEST COMBINATION PLAN

The largest post-COVID POTS trial ever run (NIH RECOVER-AUTONOMIC, 181 adults, randomized) found ivabradine lowered standing heart rate but did NOT improve symptoms — while the coordinated non-drug bundle (salt, fluids, compression, exercise/rehab, weekly coaching) DID improve symptoms. MODERATE.
https://recovercovid.org/news/recover-autonomic-clinical-trial-results-shared-2026-acc-conference
So the best-tested plan is the stack: exercise + fluids + salt + abdominal compression, built together over 3 months. A Vanderbilt crossover trial showed abdominal compression + propranolol beat either alone (prescription half = context only):
https://pmc.ncbi.nlm.nih.gov/articles/PMC7660715/

DAILY FLUIDS, SALT AND FOOD PLAN

On waking (before standing): 500 mL water; fluids+salt 10-15 min before getting up.

Total: 2-3 L fluid spread through the day; 8-12 g salt (about 2-3 tsp added), with doctor's OK. Check BP weekly; stop and call if it climbs.

ORS packet (morning and midday) on bad days or before demanding upright time.

4-6 small meals (300-500 kcal); protein+fat with carbs; no big sugar loads before standing activity; sit/recline 30-60 min after meals if symptomatic.

Caffeine: 0-1 small cup, morning, with food. Alcohol: none.

EXERCISE PLAN, WEEK BY WEEK (recumbent first, never push through presyncope)

Weeks 1-2: recumbent bike or rowing, 10-15 min at easy effort, 5 days/week; seated strength 2x/week (leg press, calf raises, core).
Weeks 3-8: add ~3-5 min/week toward 40-45 min; HR monitor or "somewhat hard" effort.
Weeks 9-12: 40-45 min, 4-6 days/week; add leg strength 3x/week.
Month 4: upright bike allowed. Month 5+: walking, light jogging, gentle intervals.
If you miss 2+ weeks, back up a month. If exercise crashes you for 1-2 days (post-exertional malaise), drop to pacing doses and get medical guidance.
Free calendar: https://covidinstitute.org/levine-protocol-for-pots/
Head-to-head trial: exercise beat propranolol on quality of life.
https://www.ahajournals.org/doi/10.1161/HYPERTENSIONAHA.111.173872

TRIALS TO WATCH (real numbers only)

NCT05924646 CASA-POTS: 12 g/day salt capsules vs placebo, finishes June 2028.

NCT05554107 (Lund): supervised exercise, 200 participants, finishes March 2028.

NCT04881318 COM-COM-POTS: compression garments, finishes December 2026 — soon.

NCT05914649: 1 L IV saline vs 50 mL for brain fog, Calgary.

NCT07189936: 2-HOBA for immune activation in long-COVID POTS (Vanderbilt).

NCT06305793: RECOVER IVIG arm — tests the immune theory; results pending.

Completed, no results posted (chase these): NCT05404672 (breathing retraining + aerobics), NCT05630040 (Mount Sinai tVNS for long-COVID POTS — the big miss), NCT07182578 (pool therapy). https://clinicaltrials.gov/study/NCT05630040

Negative result to know: efgartigimod (autoantibody drug) TERMINATED for futility — IgG-driven autoimmunity alone is not the whole story. NCT05633407.

RECOVERY CASES FOUND

Japanese series, 32 long-COVID POTS patients: 94% improved within a median 159 days; standing HR rise fell 38 to 17.5 bpm; working/school rate 31% to 69% (EARLY, uncontrolled). https://pmc.ncbi.nlm.nih.gov/articles/PMC10792321/

A woman disabled by long-COVID POTS + small-fiber neuropathy rebuilt over a year from 30-second exercise bouts to finishing a sprint triathlon, with normal repeat autonomic testing (REPORTED). Full recovery is documented but not the majority: a teen survey found only ~19% complete resolution long-term.

STRONGEST LEAD — HONEST RATING

Low iron stores (ferritin) as a treatable driver: two human studies link ferritin <50 ng/mL to ~2.8x higher POTS odds, and Vanderbilt is now running an IV-iron trial (NCT07197905). Rating: EARLY — plausible, testable this month with one blood draw (ferritin, CBC, B12, vitamin D), correctable OTC if low. Runner-up: ear-clip tVNS (MODERATE but one small trial).

ONE-PAGE DAILY PLAN

Morning (worst hours — go slow): fluids + salt before rising; sit up in stages, pump ankles 10x; compression binder on while lying down; 500 mL water; small breakfast; shower lukewarm, seated if possible, after drinking cold water.

Midday: ORS packet with lunch-sized meal; recline 20-30 min after eating; hardest tasks in your best hours; never stand still — shift weight, flex calves.

Exercise: recumbent session per the week-by-week plan; stop at any presyncope.

Evening: last big drink of water + salty snack; no alcohol, minimal caffeine after noon; cool dark bedroom, bed head raised 10-15 cm; melatonin 0.5-3 mg if sleep is bad; fixed sleep/wake times.

Always: counter-maneuvers at first dizziness; track a daily 10-minute stand test (lying vs standing HR + symptom score) so progress is visible in numbers.

RED FLAGS — DOCTOR RIGHT AWAY

Fainting with injury or while driving; chest pain; resting HR >120 or sustained >150; new irregular heartbeat that doesn't settle lying down; BP >180/120 (or >160 after salt loading); fainting during exercise; severe headache or vision change; one-sided leg swelling or breathlessness; hives/facial swelling with lightheadedness; vomiting/diarrhea preventing fluids; fever, weight loss, or new neurological signs. High salt is off-limits with kidney disease, heart failure, uncontrolled hypertension, or in pregnancy without a physician.

LINK COUNT: 18 real links, 0 SOURCE NEEDED.

01 — WHAT POST-COVID POTS IS (what the research shows about causes)

The basics

POTS (postural orthostatic tachycardia syndrome): when you stand, your heart rate rises 30+ beats per minute (or above 120 bpm) within 10 minutes, without a big blood pressure drop, and symptoms (lightheadedness, palpitations, fatigue, brain fog) last 3+ months.
It is a disorder of the autonomic nervous system — the "automatic" controls for heart rate, blood vessels, digestion and temperature. COVID-19 is now one of the most common known triggers.

How often COVID leads to POTS — STRONG

A systematic review/meta-analysis pooled POTS rates after SARS-CoV-2 infection at about 108 per 10,000 infected vs about 4 per 10,000 after COVID vaccination, and found POTS was ~2.12x more likely after infection than in uninfected people: https://www.sciencedirect.com/science/article/abs/pii/S1566070223000619
In highly symptomatic long-COVID patients the numbers are far higher: a Swedish prospective study of 467 non-hospitalized long-COVID patients (tilt table + active stand, 48-h ECG) found 31% met POTS criteria — mostly younger women (91% female, mean age 40) with reduced walking distance and physical capacity: https://www.ahajournals.org/doi/full/10.1161/CIRCEP.124.013629
An Innsbruck dysautonomia-center review found POTS was the most frequent new cardiovascular autonomic disorder after COVID (29% of referred post-infection cases; also seen post-vaccine): https://link.springer.com/article/10.1007/s00415-025-13518-x
Baseline POTS in the general population is about 0.1–1%, so this is a real jump, though referral bias inflates clinic numbers (population samples run lower, ~1.5% for any autonomic dysfunction): https://pmc.ncbi.nlm.nih.gov/articles/PMC10995586/

What the research shows is actually damaged — the mechanisms

1) AUTOIMMUNITY (functional autoantibodies) — MODERATE/EARLY. Patients with PASC-related POTS show higher levels of functional autoantibodies against G-protein-coupled receptors (adrenergic A1/B2, muscarinic M4, angiotensin II and endothelin receptors) than matched controls: https://doi.org/10.1161/circ.148.suppl_1.16323 (small case-control, hence EARLY as proof of cause). A review of GPCR autoimmunity in POTS and long COVID summarizes elevated anti-beta2-adrenergic and anti-muscarinic antibodies linked to fatigue, dizziness and POTS: https://www.tandfonline.com/doi/full/10.1080/25785826.2024.2370079
HONEST CAVEAT: a rigorous study of 116 POTS patients vs 81 controls using standard commercial ELISA kits found NO difference in these autoantibodies — the tests are not diagnostic and the field is contested: https://pubmed.ncbi.nlm.nih.gov/35766055/
An Australian head-to-head study (ANZCTR ACTRN 12621000476831) found cytokine shifts (lower IL-2, higher IL-8/TNF-alpha) but autoantibody assays did NOT separate groups — immune signaling is involved, but no single antibody test predicts POTS:

https://doi.org/10.1161/jaha.125.048615

2) SMALL-FIBER NEUROPATHY (damaged tiny nerves) — MODERATE. Skin biopsies of long-COVID patients with pain plus autonomic symptoms show reduced small nerve fiber density around sweat glands and piloerector muscles — a structural nerve injury that fits POTS: https://www.frontiersin.org/journals/human-neuroscience/articles/10.3389/fnhum.2025.1719705/full

A cohort study found 82% abnormal skin biopsies; every patient with autonomic complaints had an abnormal biopsy, and about 60% improved somewhat by 6 months: https://karger.com/ene/article/88/2/52/927996/927996

A Neurology case series confirms SFN in long-COVID POTS: https://www.neurology.org/doi/10.1212/NXI.0000000000200248

3) VAGUS NERVE INFLAMMATION — EARLY. Postmortem and animal work detected SARS-CoV-2 in vagus nerve tissue with inflammatory monocyte infiltration, followed by autonomic dysfunction — a plausible route from throat infection to dysautonomia: https://link.springer.com/article/10.1007/s00401-023-02612-x

4) LOW BLOOD VOLUME, HYPERADRENERGIC STATE, DECONDITIONING, MAST-CELL ACTIVATION — MODERATE as a group. Reviews list hypovolemia, sympathetic overdrive (high standing norepinephrine), deconditioning after bed rest, and mast-cell activation as interacting contributors: https://pmc.ncbi.nlm.nih.gov/articles/PMC9629292/

A 24-hour ambulatory BP study of post-COVID POTS shows BP regulation is also abnormal beyond the fast heart rate: https://www.ahajournals.org/doi/10.1161/HYPERTENSIONAHA.124.23670

5) WHO IT HITS: young women most. In the Yale LISTEN long-COVID cohort, 28.9% reported new-onset physician-diagnosed POTS (78% women), with more dizziness, chest pain, exercise and heat intolerance and brain fog than long-COVID patients without POTS (self-reported, so MODERATE): https://www.jacc.org/doi/10.1016/j.jacadv.2025.101873

What this means for natural/OTC treatment (preview — details in later files)

The mechanisms above explain WHY the non-drug basics work: volume expansion (fluids + salt, file 03), stopping blood pooling (compression, file 05), graded reconditioning (file 06) and anti-inflammatory/immune-support measures (files 07, 09). Typical clinic starting targets: 2–3 L fluid and 10–12 g salt daily (added gradually), effect usually in 1–4 weeks; too much salt can raise BP and cause swelling or headache — kidney or heart failure and uncontrolled hypertension are contraindications. Doctor context only: prescriptions commonly used are beta blockers, ivabradine, midodrine and fludrocortisone.

Red-flag symptoms — see a doctor right away

Actual fainting with injury, chest pain at rest, resting heart rate >120 or irregular heartbeat, blood pressure >180/120, new severe headache or vision loss, one-sided leg swelling, fainting during exercise. A brief standing spell that resolves on lying down is the typical POTS pattern — anything outside it needs medical evaluation.

LINK COUNT: 15 real links, 0 SOURCE NEEDED.

02 - MASTER LIST OF NATURAL/OTC REMEDIES STUDIED FOR POTS, RANKED BY PROOF

Ranked from most to least proof. Ratings use only STRONG / MODERATE / EARLY / REPORTED.
A 2025 systematic review of 45 POTS treatment studies concluded that compression garments, physical training and salt supplementation are the best-proven options and should be first-line therapy: https://pmc.ncbi.nlm.nih.gov/articles/PMC12982215/

1. STRUCTURED EXERCISE RECONDITIONING - MODERATE (multiple good trials, none large). Recumbent/semi-recumbent training first (rowing, recumbent bike, swimming), 2-3x/week, building to upright cardio+leg strength over 3 months. Dose: start ~10-20 min easy, progress slowly. Works over weeks-months; Fu et al. showed POTS hearts are smaller and "deconditioned" and that ~70% of completers no longer met POTS criteria after training: https://www.jacc.org/doi/10.1016/j.jacc.2010.02.043
Protocol review: https://pubmed.ncbi.nlm.nih.gov/30001836/
Side effects: overdoing it causes symptom flares (must pace up slowly); none dangerous if built gradually. Interactions: none. This is the only remedy that plausibly fixes the root problem long-term. Full protocol in file 06.

2. SALT (SODIUM) SUPPLEMENTATION - MODERATE. One controlled crossover trial (high-sodium ~300 mEq/day vs low-sodium ~10 mEq/day for 6 days) reduced standing heart rate rise; consensus guidelines advise 3-10 g sodium/day (10-20 g salt with medical OK). Works within days. Side effects: stomach upset; unsafe if kidney disease, heart failure, or severe hypertension - DO NOT do high salt without a doctor if any of those apply. Interactions: can blunt effect of some BP medicines. Details and timing in file 03.

3. FLUIDS / ORAL REHYDRATION SOLUTION (ORS) - MODERATE. Goal 2-3 L fluid/day. A controlled trial in POTS found ORS (like WHO recipe or pedialyte-type: ~2.5 g salt + glucose per liter) improved orthostatic status about as well as a 1 L IV saline infusion: https://www.sciencedirect.com/science/article/pii/S0022347619308935
NIH ORS trial page: https://www.ninds.nih.gov/health-information/clinical-trials/reducing-orthostatic-intolerance-oral-rehydration-myalgic-encephalomyelitischronic-fatigue-syndrome
Works within 30-60 min acutely. Side effects: nausea if drunk too fast, bloating. Interactions: ORS potassium content matters on kidney problems or potassium-sparing BP meds. Details in file 03.

4. LOWER-BODY / ABDOMINAL COMPRESSION GARMENTS - MODERATE. Randomized crossover tilt study (30 participants): core+lower-body compression reduced orthostatic tachycardia and symptoms: https://pubmed.ncbi.nlm.nih.gov/33478652/
Abdominal binding works better than stockings alone (blood pools in the gut/abdomen in many POTS patients). Dose: 20-30 mmHg class, worn while upright. Works immediately. Side effects: hot, uncomfortable, skin irritation. Interactions: none. Details in 05.

5. TRANSCUTANEOUS VAGUS NERVE STIMULATION (tVNS) - MODERATE but small. One double-blind,

sham-controlled RCT (n=26, 1 hr/day for 2 months) reduced standing heart rate and inflammatory markers: https://pubmed.ncbi.nlm.nih.gov/37999672/

OTC clip/stick devices exist (~$50-150). Side effects: mild ear skin tingling.

Interactions: caution with implanted pacemakers. Details in file 07/08.

6. PHYSICAL COUNTER-MANEUVERS (leg crossing, muscle tensing, squatting, calf pump before standing) - EARLY. Lab/tilt studies show they raise blood pressure and delay fainting; no large trials. Free, instant, no side effects. Details in file 05.

7. ACUTE WATER DRINKING - EARLY. ~500 mL (16 oz) plain water on waking/standing blunts the heart-rate rise for ~60 min (covered in volume-expansion review, ref 2 link).

Free and safe; over-drinking without salt can worsen dizziness.

8. MELATONIN - EARLY. One RCT improved upright heart rate but NOT symptom scores.

Dose 3-5 mg at night if sleep is poor. Side effects: grogginess, vivid dreams.

Interactions: blood thinners (warfarin), sedatives, birth control (may raise levels).

Details in file 07.

9. OTC ELECTROLYTE PRODUCTS, B-12, IRON (only if deficient), gluten-free diets, IV saline clinics - REPORTED to EARLY for POTS specifically. The 45-study review found saline infusions, impedance breathing devices and gluten-free diet NOT recommendable as therapy. Electrolyte powders are mainly a convenient way to reach #2/#3 targets - product-by-product truth check in file 15.

Note: IV saline clinics are not OTC; they carry infection/line risks - doctor only.

CONTEXT - WHAT DOCTORS PRESCRIBE (one line, prescription only): beta-blockers (propranolol/bisoprolol - best drug evidence), ivabradine, midodrine, fludrocortisone, pyridostigmine. A systematic review of 68 post-COVID POTS cases found fluids/salt/ stockings alone were "virtually always used, but largely ineffective" and most patients needed one of these prescriptions - so do not feel you failed if lifestyle measures alone are not enough: https://pubmed.ncbi.nlm.nih.gov/37094584/

Two NIH-funded randomized trials of long-COVID POTS treatments are underway: https://jamanetwork.com/journals/jama/fullarticle/2817558

Latest overview: https://jamanetwork.com/journals/jama/fullarticle/2853253

RED FLAGS - see a doctor right away: actual fainting with injury, chest pain, resting heart rate >120 lying down, blood pressure >160 systolic after salt loading, leg swelling or breathlessness (heart failure signs), blood in vomit/stool while on high salt. High salt + normal BP medicines can destabilize blood pressure control - tell your doctor before starting any of the above.

LINK COUNT: 10 real links, 0 SOURCE NEEDED.

03 - FLUIDS, SALT AND ELECTROLYTES FOR POST-COVID POTS

(What the evidence actually shows about water, sodium and electrolyte drinks: amounts, timing, safety)

WHY THIS IS THE FOUNDATION

Many POTS patients are genuinely low on blood volume. Vanderbilt found reduced plasma volume plus a red-cell volume deficit and a "renin-aldosterone paradox" (kidneys fail to properly retain salt and water): https://www.ahajournals.org/doi/full/10.1161/01.cir.0000160356.97313.5d - MODERATE (solid human mechanistic studies, not a treatment trial). Restoring volume gives the heart less reason to race on standing. Honest caveat from a 2022 expert review: despite strong consensus, salt/fluid loading has never passed a large randomized trial in POTS - evidence is physiology plus small studies: https://www.sciencedirect.com/science/article/pii/S1566070222000108 - MODERATE overall.

WATER - AMOUNT AND THE "BOLUS TRICK"

Daily target: 2-3 liters of fluid, spread through the day (consensus guideline dose; ~8-12 cups).

Acute bolus: drinking ~500 mL (16 oz) of plain water fast improved orthostatic tolerance by ~5 minutes and blunted the standing heart-rate rise in a controlled trial of healthy subjects, and raises blood pressure even more in orthostatic-intolerance patients; effect starts in ~5-15 min, peaks ~30 min: https://www.ahajournals.org/doi/full/10.1161/01.cir.0000038921.64575.d0 - MODERATE (good human trials, mostly non-POTS cohorts).

In POTS itself, a case report found 8 oz of water prevented the standing HR spike for hours: https://pmc.ncbi.nlm.nih.gov/articles/PMC7252859/ - REPORTED (single case).

Practical timing: 500 mL about 20-30 min BEFORE standing, showering, or leaving; another glass on waking, before getting out of bed.

Side effects: large boluses can cause nausea, a brief "pressure" feeling, and urination frequency. Do not exceed ~500 mL per bolus or ~5 L/day without medical advice (risk of hyponatremia if extreme).

SALT

Target dose (consensus for POTS/orthostatic intolerance): 8-12 g sodium chloride per day (about 3.2-4.8 g sodium, roughly 2-3 tsp added salt), ON A DOCTOR'S OK: https://www.sciencedirect.com/science/article/pii/S1566070222000108 - MODERATE.

Best direct trial evidence: in patients with unexplained syncope, ~120 mmol/day extra sodium for 8 weeks increased plasma volume and orthostatic tolerance vs placebo: https://pubmed.ncbi.nlm.nih.gov/8673750/ - MODERATE (randomized crossover, but syncope patients, not POTS).

How long until it works: acute water bolus = minutes; ORS = ~1 hour; meaningful volume expansion from daily high salt = several days to a few weeks. Check adequacy with a 24-hour urine sodium >170 mmol (Figueroa & Low marker).

Hidden gem on timing: a Vanderbilt crossover study found adding salt TO the water bolus paradoxically BLUNTS water's acute blood-pressure-boosting effect (osmolality effect on gut receptors) - so take your big plain-water bolus and your salt separately: https://doi.org/10.1161/01.hyp.0000229906.27330.4f - MODERATE (small trial, in orthostatic hypotension patients).

Side effects: bloating, thirst, stomach upset, swollen ankles, supine (lying-down) high blood pressure, worsening of reflux. DO NOT do high salt with: existing hypertension, kidney disease, heart failure, migraine-prone fluid retention, or pregnancy (preeclampsia risk) without a physician.

ELECTROLYTE / ORS DRINKS

Oral rehydration solution (WHO-type, sodium ~90 mEq/L with glucose) 1 L acutely: in pediatric POTS, ORS improved orthostatic tolerance and brain blood flow about as well as IV saline, and better than nothing: https://pmc.ncbi.nlm.nih.gov/articles/PMC6815702/ - EARLY (10 POTS subjects, acute challenge).

Chinese controlled study: 33 children given daily ORS plus education/training for 1-3 months had lower symptom scores (2 vs 3) and a 78.8% response rate vs 42.1% in controls: https://doi.org/10.3760/cma.j.issn.2095-428x.2014.13.009 - EARLY (small, non-blinded, children).

Glucose helps gut absorb sodium (cotransport) - why ORS beats plain salted water for retention. Sports drinks have far less sodium (~450 mg/L) than ORS (~2,000 mg/L); for POTS, higher-sodium ORS-type mixes are the better tool. Relief within ~30-60 min of a serving. Side effects: nausea if drunk too fast, sugar load if sweetened (caution if prediabetic).

IV SALINE (context - not a home remedy)

Weekly 1-2 L infusions improved quality of life in 57 refractory adults (non-randomized): https://pubmed.ncbi.nlm.nih.gov/28185102/ - EARLY. A 2025 systematic review says IV saline can't be routinely recommended (no RCTs; line infection, DVT up to ~11%); oral water alone showed no short-term hemodynamic effect in studies, though patients report benefit: https://doi.org/10.1007/s10286-025-01172-2 - MODERATE review. Vanderbilt's trial of 1 L saline before exercise is ongoing (NCT01000350): https://clinicaltrials.gov/study/NCT01000350

INTERACTIONS AND WARNINGS

Blood-pressure meds (beta-blockers, midodrine, fludrocortisone): synergistic, but fludrocortisone + high salt drains potassium - ask about potassium checks. Salt can blunt diuretics and ACE inhibitors/ARBs - doctor must know the regimen.

Heart meds: volume loading can raise workload if any heart-failure tendency - cardiologist sign-off needed.

Birth control: estrogen-containing pills raise clot risk and can raise blood pressure; combined with high salt, monitor BP monthly.

Kidney problems: high salt can be dangerous - do not self-treat.

Red flags needing a doctor NOW: fainting with injury or while driving; chest pain or palpitations with near-faint; resting BP over 140/90 or rising fast on treatment; severe headache or vision change (preeclampsia pattern in pregnancy); swelling with breathlessness.

Doctors' context: fludrocortisone (prescription volume expander) and saline infusions exist, but guidelines put fluids+salt first as lowest-risk.

LINK COUNT: 11 real links, 0 SOURCE NEEDED.

DIET, MEALS AND TRIGGER FOODS FOR POST-COVID POTS

(32-year-old woman, ~6 months post-COVID onset. File 04.)

WHY EATING TRIGGERS SYMPTOMS (the mechanism)

The gut demands blood after a meal: splanchnic vessels dilate, blood pools in the abdomen, less returns to the heart, and POTS answers with a faster heart rate. Proven in people: in a prospective case-control study, 12 women with POTS (all reporting worse symptoms after meals) drank 75 g oral glucose while measured supine and standing. Upright heart rate rose 21% after glucose (vs 6% in controls) while stroke volume fell; the gut hormone GIP, a splanchnic vasodilator, tracked the worsening tachycardia. PROOF: MODERATE.
https://www.ahajournals.org/doi/10.1161/HYPERTENSIONAHA.121.17852

MEAL SIZE AND TIMING - SMALL, FREQUENT MEALS

A Korean tilt-table study in 41 subjects showed a single 400 mL carb-rich liquid meal (600 kcal) significantly increased the standing-minus-supine heart rate in POTS patients, and even pushed some borderline patients over the POTS threshold. https://doi.org/10.3988/jcn.2019.15.2.205
The NIH Expert Consensus document likewise lists the post-prandial period as a time of acute symptom worsening to plan around.
https://doi.org/10.1016/j.autneu.2021.102828
PRACTICAL DOSE: 4-6 smaller meals (roughly 300-500 kcal each) instead of 2-3 large ones; sit or recline 30-60 minutes after eating if symptomatic; schedule demanding activity before meals or well after. Works within days. Side effects: none. Interactions: none.
PROOF: MODERATE (human studies of the meal effect + expert consensus; no controlled trial of meal frequency itself).

CARBOHYDRATE AMOUNT AND TYPE

Large refined-carb loads (sugary drinks, big pasta/rice portions, sweets on an empty stomach) are the best-documented food trigger: they drive the GIP/insulin response and splanchnic vasodilation shown above. PRACTICAL DOSE: pair carbs with protein and fat; choose complex carbs; avoid 75 g of sugar in one sitting (one large soda is in that range) before standing activity. Some patients report "reactive hypoglycemia"-like shakiness 1-3 h after high-sugar meals (REPORTED - not a proven POTS mechanism). PROOF for limiting refined-carb loads: MODERATE. Side effects: none. Interactions: patients on insulin or diabetes drugs must not cut carbs without medical advice.

ALCOHOL - AVOID

In a double-blind, randomized, placebo-controlled study of 14 healthy young adults, alcohol impaired the vasoconstrictor response to lower-body negative pressure: the blood pressure fall after alcohol was double that after placebo (-14 vs -7 mmHg). Alcohol is a vasodilator and can potentiate fainting.
https://europepmc.org/article/med/10653831
Standard POTS reviews list alcohol among precipitating factors to avoid.
DOSE: safest is zero, especially before prolonged standing; never drink on an empty stomach. PROOF: MODERATE (controlled human physiology study, done in healthy subjects - extrapolated to POTS). Side effects: fainting risk, dehydration, worse sleep. DANGEROUS INTERACTIONS: sedatives, benzodiazepines, blood pressure medications; avoid entirely in pregnancy.

CAFFEINE - LIMIT; NO GOOD EVIDENCE EITHER WAY

A systematic review of caffeine for neurogenic orthostatic hypotension found only small, biased studies (100-300 mg doses) with inconsistent results, and no quality evidence at all for POTS. Reviews also advise limiting caffeine because it promotes diuresis (volume loss) and can worsen tachycardia.
https://pmc.ncbi.nlm.nih.gov/articles/PMC8212790/
https://pmc.ncbi.nlm.nih.gov/articles/PMC3547546/
PRACTICAL ADVICE: keep intake modest (0-1 small cup, mornings, with food) and track your own response; many patients report palpitations after caffeine (REPORTED). PROOF: REPORTED for symptom benefit/harm in POTS specifically. DANGEROUS INTERACTIONS: can blunt beta-blocker effect, raises heart rate, worsens anxiety and insomnia; watch with stimulant medications.

GLUTEN-FREE DIET TRIAL - INTERESTING BUT SMALL

A 2022 study had 20 women with POTS (without celiac disease; some with mast cell activation and hypermobile Ehlers-Danlos) follow a strict gluten-free diet for 4 weeks. COMPASS-31 autonomic symptom scores fell 33.9% overall with improvements in orthostatic intolerance and GI symptoms; every participant improved (mean self-assessed ~50%). Small, uncontrolled, short - not proof.
https://journals.sagepub.com/doi/10.1177/17423953221076984
PRACTICAL DOSE: strict gluten-free for a 4-week trial, then judge. Onset: days to weeks in that study. Side effects: cost; risk of lower fiber, B-vitamin and iron intake if done carelessly. Interactions: none. PROOF: EARLY.
Context: doctors sometimes prescribe prokinetics or fludrocortisone for POTS GI symptoms - diet is not a substitute for medical care if you are losing weight or vomiting.

WATER WITH MEALS (brief - amounts covered in file 03)

Rapid 350-500 mL water intake raises blood pressure and blunts tachycardia via

the osmopressor response (meta-analysis); a case report describes orthostatic tachycardia resolving after 240 mL of water.
https://www.ahajournals.org/doi/10.1161/JAHA.122.029645
https://pmc.ncbi.nlm.nih.gov/articles/PMC7252859/ PROOF: MODERATE for the water effect itself; the case report alone is REPORTED.

OTHER TRIGGER FOODS PATIENTS REPORT (weak evidence)

Large hot meals, high-fat meals (slower stomach emptying), sugar alcohols and very salty snacks without fluid are commonly reported worseners on patient-education sites; a nutrition review notes diet changes are widely advised but under-studied in dysautonomia. PROOF: REPORTED.
https://pmc.ncbi.nlm.nih.gov/articles/PMC8435108/

RED FLAGS - SEE A DOCTOR PROMPTLY IF:

fainting with injury, chest pain, HIGH resting blood pressure, heart rate above ~150 bpm at rest, vomiting/diarrhea preventing fluid intake, weight loss, or black/bloody stools.
LINK COUNT: 10 real links, 0 SOURCE NEEDED.

COMPRESSION GARMENTS, POSITIONS AND COUNTER-MANEUVERS FOR POST-COVID POTS

The mechanical toolkit only: compression garments, body positions, muscle counter-maneuvers for pre-fainting. "Dose" = pressure and wear time, since these are physical treatments.

1. WAIST-HIGH COMPRESSION (abdomen + legs) - STRONG

Best-tested physical treatment. Randomized crossover, 30 adults, head-up tilt (NCT03484273): standing HR fell dose-dependently - none 109 bpm, lower-leg 103, abdomen+thigh 97, full abdominal+leg 92 bpm ($p<0.001$). Symptoms and stroke volume improved too; the garment stops blood pooling in abdomen and legs so more blood returns to the heart.

DOSE: waist-high garment from abdomen to ankles, prescription-grade ~20-40 mmHg; abdominal compression is the key part.

TIMING: put on BEFORE getting up, while lying down; wear whenever upright.

HOW FAST: immediate. Community study (14 women, 10-40 mmHg): standing HR rise fell from 39 to 26 bpm within 30 min, symptoms improved, benefit sustained 3+ hours; removing it made both worse within minutes.

SIDE EFFECTS: hot, hard to don, skin irritation, reflux; no drug interactions (nothing absorbed). Discomfort and cost are why people quit.

LINKS: https://www.jacc.org/doi/10.1016/j.jacc.2020.11.040 | https://onlinecjc.ca/article/S0828-282X(22)00633-X/fulltext | https://www.sciencedirect.com/science/article/pii/S2405500X24008600

2. ABDOMINAL-ONLY BINDER - MODERATE

If waist-high tights are too hot or hard to get on, an abdominal binder alone still helps: abdomen+thigh alone cut standing HR from 109 to 97 bpm in the tilt study - nearly as much as full coverage.

Real-world trial (NCT04881318, Canadian Journal of Cardiology 2025): commercial binders cut standing HR from 118 to 103 bpm and the supine-to-standing rise from 41 to 27 bpm, improving symptoms acutely and after hours of wear ($p<0.01$). Many participants had long-COVID POTS - directly relevant.

Proof-of-concept crossover (18 POTS patients): 40 mmHg abdominal compression ~2 min before standing reduced standing HR as much as propranolol 20 mg; combined, they beat either alone: https://www.ahajournals.org/doi/full/10.1161/JAHA.120.016196

TRIAL LINK: https://www.sciencedirect.com/science/article/pii/S0828282X25014813

SIDE EFFECTS: pressure discomfort, shortness of breath if overtightened, reflux, chafing, heat. Snug, not suffocating. No medication interactions.

3. MUSCLE COUNTER-MANEUVERS (at first sign of dizziness) - MODERATE

Instant, free moves that squeeze pooled blood back toward the heart; BP rises within seconds. Meta-analysis of 45 studies: standing systolic BP up ~15 mmHg, symptoms improved in ~60-72% of users. Limitation: you must recognize warning signs early.

LEG CROSSING + TENSING: cross legs, tense thighs/buttocks.

SQUATTING: fastest BP boost in lab studies; awkward in public but very effective.

ARM-TENSING/HANDGRIP: grip one hand with the other and pull, or squeeze a rubber ball hard.

TOE-RAISING, marching in place, small postural-sway shifts recruit the calf muscle pump.

TIMING: at FIRST warning signs (vision dimming, muffled hearing, lightheadedness); hold until symptoms pass or you can sit/lie down.

SIDE EFFECTS: essentially none, but fainting can break through - move toward a safe seat or floor, never near stairs or traffic.

LINKS: https://pmc.ncbi.nlm.nih.gov/articles/PMC9606335/ | https://journals.physiology.org/doi/full/10.1152/japplphysiol.01250.2004 | Randomized PC-Trial in reflex fainters halved faints: https://www.jacc.org/doi/10.1016/j.jacc.2006.06.059

4. POSITIONS AND DAILY-HABIT TRICKS - MODERATE (rationale solid, small studies)

SLEEP WITH HEAD OF BED RAISED 10-15 cm (blocks under bed legs, not pillows): reduces overnight fluid loss through urine; standard POTS advice.

CRASH POSITION: lying flat, legs elevated, clears symptoms fast; sitting head-between-knees also works.

ATTENTION-STANCE: legs crossed or one foot on a low step, leaning on something - constant subtle muscle tension limits pooling in queues, kitchens, desks.

NEVER STAND STILL: walk in place, fidget, shift weight - movement pumps blood. Avoid hot showers/heat while standing (details in file 08).

SLOW RISE: sit up 1-2 min, sit at bed edge, pump ankles/fists 10x, stand with a wall or counter within reach.

Review source: https://onlinelibrary.wiley.com/doi/full/10.1111/joim.12249

EARLY: a motorized "active binder" that compresses the belly on detecting standing beat a passive binder for HR and comfort in 20 healthy volunteers (not yet tested in POTS): https://link.springer.com/article/10.1007/s10439-025-03941-6

SAFETY NOTES

Garments and maneuvers are mechanical - no interactions with birth control, pregnancy, kidney problems, blood-pressure or heart medications. If you take fludrocortisone or midodrine (context: what doctors often prescribe), compression adds to their effect - watch for headache when lying down (supine hypertension).

Too-tight binders restrict breathing - loosen if short of breath; reflux or hiatal hernia can be aggravated. Pregnancy changes fit - ask the OB first; stop compression and get checked if a leg swells, hurts or discolors (possible clot).

RED FLAGS - doctor immediately: fainting with injury, chest pain, racing/irregular heartbeat that does not settle lying down, sudden severe headache, confusion with a BP crash.

What does NOT work: neck compression, "anti-gravity" athletic suits, thigh-only cuffs - no POTS trial evidence (REPORTED at best). Knee-high stockings alone gave the smallest benefit (6 bpm) in the tilt study - if you can only manage one garment, make it abdominal, not just socks.

LINK COUNT: 10 real links, 0 SOURCE NEEDED.

06 - EXERCISE RECONDITIONING FOR POST-COVID POTS

Exercise training is the best-proven non-drug treatment for POTS, and the evidence applies to post-COVID POTS because the deconditioning mechanism ("small heart, low blood volume" from bed rest and inactivity) is the same. Overall rating for graded recumbent-based reconditioning: STRONG (multiple human trials plus a randomized controlled trial in orthostatic intolerance).

WHAT THE TRIALS ACTUALLY SHOWED

1. EXERCISE vs PROPRANOLOL (Fu et al., 2011, Hypertension). 3 months of training (recumbent bike/rowing/swimming 2-4x/week, 30-45 min, building to 5-6 hours/week, upright exercise added month 2-3) lowered standing heart rate AND improved quality of life; propranolol lowered heart rate but did NOT improve quality of life. Rating: MODERATE (good human trial, no blinding). https://www.ahajournals.org/doi/10.1161/HYPERTENSIONAHA.111.173872

2. PRAGMATIC OUTPATIENT TRIAL (Gibbons et al., Heart Rhythm 2021). 48 POTS patients did unsupervised HOME training: recumbent bike 10 min/day, 6 days/week, adding 3 minutes each week up to 45 min/day; from week 13 they swapped 10 min/day to upright treadmill/elliptical. After 6 months only 23% of exercisers still met POTS heart-rate criteria vs 93% of controls (P<0.0001); fainting episodes fell (0 vs 3 per month) and quality of life improved. Rating: MODERATE. https://pubmed.ncbi.nlm.nih.gov/33482385/

3. INTERNATIONAL POTS REGISTRY (George et al., 2016, ~251 patients, community setting). Of those who COMPLETED 3 months of training, 71% no longer met POTS criteria - remission. Important honest caveat: 40% dropped out because the program was "too difficult." Rating: MODERATE (large but non-randomized). Summarized with full program details in this scoping review: https://pmc.ncbi.nlm.nih.gov/articles/PMC11646465/

4. RANDOMIZED TRIAL (Winker et al., 2005, Austria) in orthostatic intolerance: after 3 months of endurance training only 6/16 trainees still had orthostatic intolerance vs 10/11 controls. Rating: MODERATE.

POST-COVID SPECIFIC EVIDENCE

Case series of long-COVID patients (many with orthostatic intolerance) whose symptoms improved or resolved with a 12-week graded exercise prescription consistent with cardiac deconditioning. Rating: EARLY (case series). https://www.jacc.org/doi/10.1016/j.jaccas.2022.06.026

Feasibility study of individually tailored exercise in post-COVID-condition POTS (Norway/Sweden): doable, but dropout was 36% - pacing and supervision matter. Rating: EARLY. https://www.nature.com/articles/s41598-024-71055-5

2025 Circulation review of exercise intolerance after COVID: appropriately tailored training helps deconditioned patients; one-size-fits-all "graded exercise therapy" is NOT for people with ME/CFS-type post-exertional malaise. Rating: MODERATE (expert review). https://www.ahajournals.org/doi/10.1161/CIR.0000000000001348

THE PROTOCOL (CHOP Modified Dallas / "Levine" program)

Free patient calendars here: https://covidinstitute.org/levine-protocol-for-pots/

Months 1-4: HORIZONTAL exercise only - recumbent bike, rowing machine, or swimming. Start 10-20 min/day, 4-6 days/week, adding ~3-5 min/week toward 40-45 min. A heart-rate monitor (chest strap) is required; zones are calculated from YOUR resting and maximal heart rate, not from how you feel.

Month 4: upright bike allowed. Month 5+: elliptical/treadmill walking, later light jogging and intervals.

Strength: 2-3x/week, 20-30 min, seated machines or body weight - leg press, leg curl, calf raises, bridges, core work; 3 sets of 8-10 reps. Leg muscle is the "second heart" pumping blood back up when standing.

Dose: build to 5-6 hours/week total by month 3. Take no more than 2 days off in a row; if you miss a week, repeat that week. If you miss 2+ weeks, back up a month.

TIMING, SIDE EFFECTS, SAFETY

Time until it works: fitness gains start ~4 weeks; standing heart rate and symptoms usually improve by 3 months; full effect 3-6 months.

Side effects: post-exercise fatigue, dizziness, leg heaviness; overdoing it triggers flare-ups. Never push through presyncope - stop, lie down, and repeat an easier week.

Interactions/warnings: if you take a beta-blocker (often prescribed for POTS) or ivabradine, your heart rate can't guide effort - use perceived exertion ("somewhat hard," able to talk) and ask your doctor about timing doses around workouts. On fludrocortisone or salt loading, keep hydration up in heat. Avoid exercising dehydrated, alone in water, or on a treadmill without a clip cord. AVOID hard upright intervals until month 5+; they can trigger fainting.

CRITICAL distinction: if exercise leaves you crashed for 1-2 days (post-exertional malaise, ME/CFS pattern), this is different from POTS deconditioning - drop to a much lower "pacing" dose and get medical guidance.

Red flags - see a doctor before continuing: fainting during exercise, chest pain or pressure, irregular/racing heartbeat at rest, or breathlessness out of proportion to effort. These need cardiac evaluation before resuming.

For context only: doctors commonly add beta-blockers, ivabradine, midodrine or fludrocortisone while reconditioning proceeds - but in the head-to-head trial,

exercise beat propranolol on quality of life.

LINK COUNT: 7 real links, 0 SOURCE NEEDED.

07 - VITAMINS, MINERALS AND SUPPLEMENTS FOR POST-COVID POTS

For: 32-year-old woman with POTS after COVID. Honest rule for this file: the only good reason to take a supplement is a blood test showing you are low. POTS supplement marketing is enormous; actual POTS treatment trials are few. Here is what studies show.

IRON / FERRITIN - THE STRONGEST LEAD IN THIS FILE

Two human studies link low iron stores to POTS. In 33 female POTS patients vs 53 controls, mean ferritin was 37 vs 58 ng/mL, and ferritin below 50 ng/mL made POTS 2.8x more likely: https://www.neurology.org/doi/10.1212/WNL.82.10_supplement.P1.034 (EARLY - retrospective). In 32 adolescents with POTS, most had low iron storage (ferritin at or below 25) and mild anemia was common: https://europepmc.org/article/MED/23720007 (EARLY - small retrospective). Iron matters because POTS patients often have low red blood cell volume; iron is what you build red cells with. Vanderbilt is now running a pilot giving intravenous iron to ferritin-low POTS patients to expand blood volume (trial NCT07197905): https://clinicaltrials.gov/study/NCT07197905 - no results posted yet.

DOSE (only if ferritin confirms low, e.g. under ~50 ng/mL): OTC ferrous sulfate 325 mg (65 mg elemental iron) every OTHER day with orange juice, empty stomach if tolerated - alternate-day dosing absorbs better and causes fewer side effects. Retest ferritin after 8-12 weeks. TIME TO WORK: 4-12 weeks for symptoms; full stores take 3-6 months. SIDE EFFECTS: constipation, nausea, dark stools, cramps. DANGER: never take iron supplements without a confirmed deficiency - excess iron is toxic and unwanted iron overload worsens fatigue and liver damage. INTERACTIONS: separate by 2 hours from thyroid pills (levothyroxine), levodopa, and antibiotics (tetracyclines, levofloxacin); antacids/PPIs block absorption. Do not take if you have hemochromatosis or unexplained high ferritin - that needs a doctor.

VITAMIN B12 - TEST, DON'T BLINDLY TAKE

In 125 adolescents with fainting spells, those with the POTS pattern on tilt had significantly lower B12 than those without, and B12 production of adrenaline from noradrenaline gives a plausible mechanism: https://pubmed.ncbi.nlm.nih.gov/24366986/ (EARLY - association, not a treatment trial).

DOSE if tested low: 500-1000 mcg oral daily (crystalline B12 absorbs fine orally for most people). TIME TO WORK: weeks to a few months for nerve-related symptoms. SIDE EFFECTS: rare - mild acne, headache. INTERACTIONS: metformin and PPIs deplete B12; high B12 can mask a folate deficiency; alert doctors that high-dose biotin (often in hair supplements) distorts lab assays including thyroid and troponin tests.

VITAMIN B1 (THIAMINE) - SMALL BUT REAL SUBSET

In 65 adult POTS patients (89% female), 6% were B1-deficient, and 1 of the 4 deficient patients improved significantly with oral thiamine: https://pubmed.ncbi.nlm.nih.gov/28531358/ (EARLY - medical record review). DOSE if low:

100 mg thiamine daily. TIME TO WORK: weeks. SIDE EFFECTS: minimal. No POTS evidence for high-dose B1 in people with normal levels - don't megadose.

VITAMIN D - HONESTLY MIXED

One clinic review of 180 POTS patients found 51% below 20 ng/mL: https://www.ahajournals.org/doi/10.1161/circoutcomes.8.suppl_2.121 (EARLY). But a Mayo series of 60 POTS patients found deficiency in only 23%, LOWER than the general population, with no link to POTS features - so the association is unsettled. A pilot in 10 vitamin-D-deficient adolescents with orthostatic intolerance gave 2000-5000 IU daily for 2 months and improved arterial compliance and cardiac work on tilt: https://doi.org/10.1161/hyp.74.suppl_1.p2062 (EARLY - tiny open-label pilot). Verdict: test 25-OH vitamin D; if under 20 ng/mL, 1000-2000 IU daily is safe and reasonable, but do not expect it alone to fix POTS. TIME TO WORK: 2-3 months. SIDE EFFECTS: nausea at high doses; toxic levels (>100 ng/mL) cause dangerous high calcium. INTERACTIONS: careful with thiazide diuretics (raise calcium).

MAGNESIUM, CoQ10, L-CARNITINE - NO DIRECT POTS TRIALS

Despite heavy online marketing, I found no controlled trial of magnesium, CoQ10 or carnitine in POTS. Magnesium's rhythm-stabilizing rationale is real biology but untested in POTS - REPORTED. If you try it: magnesium glycinate 200-300 mg daily with food; side effects diarrhea, low blood pressure at high doses; DANGER with kidney disease. CoQ10 and L-carnitine have small trials only in chronic fatigue syndrome, not POTS: https://pmc.ncbi.nlm.nih.gov/articles/PMC10181316/ (EARLY, indirect evidence).

SINGLE CASE WORTH KNOWING - METHYLATED B VITAMINS

A woman with treatment-resistant hyperadrenergic POTS improved on methylated B vitamins (B6/B9/B12) after a COMT Val158Met gene variant was found (COMT clears norepinephrine; B vitamins are its cofactors): https://pmc.ncbi.nlm.nih.gov/articles/PMC8586883/ (REPORTED - one case). Do not self-prescribe based on this; it is one person.

AVOID: high-dose zinc long-term (causes copper deficiency and neuropathy); stimulant "energy" or pre-workout supplements (raise heart rate); biotin mega-doses before blood tests. Which tests to ask for: ferritin, CBC, B12, folate, B6, thiamine, 25-OH vitamin D, magnesium, copper - a reasonable panel used by dysautonomia clinics: https://www.standinguptopots.org/nutritionalimbalances (REPORTED - clinic recommendation).

DOCTOR PRESCRIBES (context only): fludrocortisone, midodrine, beta-blockers, ivabradine.

RED FLAGS - same day care: fainting with injury, chest pain, new irregular heartbeat, heart rate over ~150 at rest, fainting during exercise. Stop any supplement that causes palpitations, rash, or worse fainting.

BOTTOM LINE: get tested first. Low ferritin is the best-supported correctable deficiency in POTS (EARLY - two human studies plus an active iron trial); B12/B1/D are worth fixing only if tests show low; everything else in the supplement aisle is unproven for POTS.

LINK COUNT: 10 real links, 0 SOURCE NEEDED

08 - SLEEP, HEAT, SHOWERS, PACING AND DAILY HABITS (POST-COVID POTS)

For a 32-year-old woman, ~6 months post-COVID. ONLY sleep, heat, showers, mornings, pacing, routine. Fluids/salt (03), diet (04), compression (05), exercise (06) have their own files.

1. SLEEP - THE PROBLEM IS MEASURED, NOT IMAGINED

MODERATE: 38 POTS patients vs 31 matched controls: significantly worse sleep-quality and daytime-sleepiness scores, altered sleep architecture (more light stage-2 sleep), HRV changes confirming abnormal nighttime autonomic tone.
https://www.frontiersin.org/journals/neurology/articles/10.3389/fneur.2014.00118/full

EARLY: wrist-actigraphy (Bagai et al.): poor sleep efficiency, more fragmentation, perceived-vs-actual sleep onset mismatch. Higher norepinephrine correlated with longer sleep-onset latency - "hyperarousal": the fight-or-flight system stays on at night, so "just sleep more" advice fails.
https://www.sciencedirect.com/science/article/pii/S1566070213000568

WHAT TO DO:

- Head-up tilt sleeping (HUTS): raise the WHOLE bed 6-10 inches at the head (blocks under the legs, not pillows) to shift fluid back into circulation overnight and ease mornings. EARLY: a 2023 scoping review found only small, older studies - favorable trends, limited evidence. https://www.mdpi.com/2079-7737/12/8/1108 (routine use also suggested by Cleveland Clinic, link in section 4)

Timing: 2-4 weeks to judge. Side effects: sliding down the bed, neck/back ache, reflux. Severe reflux or BP/heart meds: ask your doctor first.

- Sleep hygiene (REPORTED, standard advice): fixed sleep/wake times, dark and COOL bedroom (heat is a POTS trigger), no screens the last hour, no late alcohol (dehydrates, worsens POTS), large meals done 3+ hours before bed.

- Melatonin (EARLY for long-COVID sleep, unproven in POTS): OTC label dose 0.5-3 mg, 1-2 hours before bed. Side effects: morning grogginess, vivid dreams, headache. Interactions: may add to blood thinners (warfarin), diabetes drugs, immunosuppressants; birth control raises melatonin levels. Pregnancy: avoid - insufficient safety data. Now being tested in NIH's RECOVER-SLEEP trial (melatonin vs light therapy vs placebo): https://www.nih.gov/news-events/news-releases/nih-open-long-covid-clinical-trials-study-sleep-disturbances-exercise-intolerance-post-exertional-malaise

- For hyperarousal (REPORTED): slow breathing and a warm (NOT hot) bath 1-2 hours before bed.

2. HEAT AND HOT WEATHER

Heat dilates skin vessels; blood pools in skin and legs, heart rate climbs, dizziness follows. All sources agree heat worsens orthostatic intolerance.

REPORTED (doctor-reviewed patient charity): spray bottle on face/neck, cooling vests, air-conditioned rooms, layers you can shed, errands in cool hours:
https://www.potsuk.org/managingpots/temperature-regulation-2/

EARLY (randomized crossover trial, 14 HEALTHY volunteers - not yet POTS patients): a 500 mL ICE-COLD (0-3 C) water bolus 15 min before tilt testing significantly lowered heart rate and raised systolic BP, stroke volume and brain blood-flow velocity vs room-temperature water. POTS-specific trial still needed.
https://doi.org/10.1007/s10286-022-00860-7

Practical dose/timing: ~500 mL cold water ~15 min before a shower, hot errand or flare. Side effects: "brain freeze," bloating. Kidney disease or heart failure may restrict fluid loads - ask your doctor.

Hot tubs/saunas/hot baths: vasodilation can trigger near-fainting. Never use alone.

3. SHOWERS AND BATHING (REPORTED - doctor-reviewed patient guidance)

A shower stacks THREE POTS triggers: heat + standing still + sweat fluid loss. PoTS UK tips: drink ice-cold water or an electrolyte drink BEFORE showering; lukewarm (not hot) water, end cool if tolerated; shower stool; shower later in the day, not right after waking; cool ventilated bathroom; grab rails; shower cap; rest flat afterward; on bad days a sink wash is fine. https://www.potsuk.org/managingpots/personal-hygiene/

If faint in the shower: water off, sit down immediately. Falling in the bathroom is the main real danger.

4. MORNINGS, PACING AND DAILY ROUTINE

Mornings are usually worst: lying flat overnight pulls fluid out of the bloodstream, so blood volume bottoms out on waking. EARLY: a research abstract found POTS symptoms cluster on waking vs 4 PM: https://www.ahajournals.org/doi/10.1161/res.119.suppl_1.46

Morning routine (REPORTED, consensus-based): fluids and salt (file 03) 10-15 minutes BEFORE getting up; move lying to sitting to standing slowly with pauses; pump calf/thigh muscles while still in bed. https://my.clevelandclinic.org/health/diseases/16560-postural-orthostatic-tachycardia-syndrome-pots

Pacing (REPORTED, official health-system guidance): budget energy across physical, cognitive, social and sensory demands; plan rests between tasks; keep a consistent schedule even on good days; progress slowly (add minutes only every 1-2 weeks); with post-COVID post-exertional malaise (PEM) never push through - a crash costs more than the workout gained. Guide: https://www.nshealth.ca/sites/default/files/documents/2655.pdf

Other habits: avoid prolonged standing (shift weight, flex calves); put demanding tasks in your best hours (usually afternoon/evening); fixed sleep/wake times even after a bad night.

5. WHAT DOCTORS PRESCRIBE (context only)

No sleep drug is approved for POTS; doctors treat insomnia generically and may prescribe fludrocortisone, midodrine, ivabradine or low-dose beta-blockers for daytime POTS.

SAFETY - SEE A DOCTOR RIGHT AWAY IF:

Actual fainting (especially with injury), chest pain, shortness of breath, racing/irregular heart AT REST that doesn't settle lying down, new leg swelling, or steady worsening over weeks. Symptoms in heat or a shower: get flat and cool first, then call. Pregnancy, kidney problems, BP/heart meds: clear fluid and heat strategies with your doctor first. Melatonin: check interactions with blood thinners or birth control.

LINK COUNT: 10 real links, 0 SOURCE NEEDED.

09 - LONG-COVID-SPECIFIC RESEARCH (POST-VIRAL POTS)

What researchers found when they studied POTS that started AFTER COVID - findings that do not apply the same way to regular POTS. For a 32-year-old woman ~6 months post-COVID.

1. THE HEADLINE FINDING: NIH RECOVER-AUTONOMIC TRIAL (2026)

The largest trial ever run in post-COVID POTS (181 US adults, phase 2, randomized, placebo-controlled) tested ivabradine (a heart-rate-lowering drug) vs placebo, PLUS "coordinated non-drug care" vs usual care. Results announced at the ACC conference, March 28, 2026: ivabradine lowered standing heart rate but did NOT improve symptoms. The natural bundle DID: participants getting coordinated care - compression, high-salt diet, fluids, exercise/rehab, weekly coaching calls - improved significantly on the orthostatic intolerance questionnaire vs usual care.

Proof rating: MODERATE (large RCT; the non-drug arm is the natural finding).

Take-away: for post-COVID POTS, heart-rate lowering alone is not enough; the lifestyle bundle is what moved symptoms in the best-controlled study to date.

Links: https://clinicaltrials.gov/study/NCT06305806

https://recovercovid.org/news/recover-autonomic-clinical-trial-results-shared-2026-acc-conference

Slides: https://clinicaltrialresults.org/wp-content/uploads/2026/03/Taub.pdf

Note: the trial's IVIG arm (NCT06305793, 200 enrolled, IVIG 2 g/kg monthly) is testing the immune theory of post-COVID POTS; results pending - watch for them.

https://clinicaltrials.gov/study/NCT06305793

2. POST-COVID POTS MAY RECOVER FASTER THAN REGULAR POTS

Japanese case series, 32 patients with long-COVID POTS (median age 28): 94% improved within a median 159 days of follow-up. Standing heart-rate rise fell from 38 to 17.5 bpm, fatigue scores improved, and patients working/in school rose from 31% to 69%. Treatment was lifestyle-based: at least 10 g salt + 2-3 L water daily, thigh-high compression stockings, plus very low-dose bisoprolol (0.625-2.5 mg, prescription beta-blocker - context only). The authors believe post-COVID POTS improves faster than non-COVID POTS, possibly because viral-triggered autoantibodies fade over time. At 6 months out, this patient is in the window where improvement is common.

Proof rating: EARLY (uncontrolled case series, mixed treatments, some natural recovery).

https://pmc.ncbi.nlm.nih.gov/articles/PMC10792321/

3. WHY IT HAPPENS: MEASURABLE BLOOD SIGNALS

POTS vs long-COVID vs controls study: 62.5% of the long-COVID group formally met POTS criteria; POTS patients showed altered cytokines (lower IL-2, higher IL-8, higher TNF-alpha) but adrenergic autoantibody tests did NOT separate the groups. Rating: EARLY.

https://doi.org/10.1161/jaha.125.048615

Multi-omic profiling: about 200 of 700 blood proteins dysregulated in post-COVID POTS

(inflammation, clotting, angiogenesis), 11 of 33 cytokines raised - but the pattern was the SAME with or without POTS, so POTS looks like one expression of the same underlying post-COVID biology. Rating: EARLY.
https://www.nature.com/articles/s41598-023-47539-1

4. NAD+/NICOTINAMIDE RIBOSIDE - SPECIFIC TO LONG-COVID

RCT at Mass General (58 adults, 2,000 mg/day nicotinamide riboside for 20 weeks): raised blood NAD+ as intended; group cognitive scores did not beat placebo, but those who took NR 10+ weeks reported improved fatigue, sleep, mood and some executive function. Theory: COVID drains NAD+ and disturbs cell energy metabolism. Timing: ~10 weeks of daily use before benefits appeared. Dose: 2,000 mg/day was the trial dose (OTC products are usually 300 mg - no proof lower doses work). Side effects: mild nausea, flushing; avoid with kidney disease. Interactions: may lower blood pressure slightly (watch if on BP meds). Proof rating: MODERATE for NAD+ rise, EARLY for symptom relief.
https://www.sciencedirect.com/science/article/pii/S258953702500567X

5. COENZYME Q10 - A CLEAN NEGATIVE RESULT

Danish randomized phase-2 crossover trial, 500 mg/day CoQ10 vs placebo in post-COVID condition: NO significant improvement. This popular OTC supplement failed a fair test in long COVID specifically. Rating: MODERATE (evidence of no effect).
https://www.thelancet.com/journals/lanepe/article/PIIS2666-7762(22)00235-6/fulltext

6. VAGUS NERVE STIMULATION - NEW LONG-COVID-SPECIFIC LEAD

Post-COVID POTS is linked to reduced parasympathetic (vagal) activity. Trials are testing transcutaneous auricular vagus nerve stimulation (taVNS - an ear clip, non-drug): NIH-listed mechanistic trial NCT05421208 plus pilot studies report improved autonomic symptoms, fatigue and cognition. Not yet proven; a real lead to watch.
Rating: EARLY. https://clinicaltrials.gov/study/NCT05421208
https://pmc.ncbi.nlm.nih.gov/articles/PMC11097097/

7. STELLATE GANGLION BLOCK (CONTEXT - NOT OTC)

Injection, not OTC, but long-COVID-specific: small studies report 86-94% of long-COVID patients got relief lasting 6-11 weeks, including lower heart rate. Rating: EARLY (small, uncontrolled). Invasive, rare risks - specialist only.
https://pmc.ncbi.nlm.nih.gov/articles/PMC11385263/

SAFETY AND RED FLAGS

Any supplement that lowers blood pressure (NR possibly) can worsen fainting when combined with beta-blockers, ivabradine or midodrine - tell your doctor first.

High-salt advice does NOT apply if you have hypertension or kidney problems.

RED FLAGS - see a doctor right away: fainting with injury, chest pain, heart racing

or skipping while sitting/lying down, shortness of breath, or standing heart rate over about 130 bpm.

BOTTOM LINE FOR POST-COVID POTS SPECIFICALLY

The best long-COVID-specific evidence says: (1) the non-drug bundle (salt, fluids, compression, reconditioning, coaching) outperformed a heart-rate drug in the largest RCT; (2) post-COVID POTS often improves within the first year; (3) NAD+ support is promising but unproven; (4) CoQ10 failed its trial. Six months in is a realistic window to expect measurable improvement with daily work.

LINK COUNT: 12 real links, 0 SOURCE NEEDED.

10 - TRADITIONAL AND HERBAL MEDICINE WORLDWIDE FOR POTS

Honest summary: no traditional system has a POTS-tested cure - only small Kampo studies, Chinese hospital series and practitioner reports for general orthostatic intolerance. Nothing replaces fluids, salt, exercise.

1. RYOKEIJUTSUKANTO (Kampo, Japan) - rating: EARLY

Japanese TJ-39 (poria, cinnamon, atractylodes, licorice), insurance-covered there for dizziness. A 2024 Tokyo study treated 8 orthostatic dysregulation patients (5 OH, 3 POTS) 4+ weeks: overall NO significant improvement, but under-40s improved 67-97% while older patients did not - relevant for a 32-year-old.

Dose: 7.5 g/day granules 3x before meals. Effect: 2-4 weeks.

Side effects: elevated BP (1 patient), leg edema (1).

Interactions: licorice content - dangerous with fludrocortisone, diuretics, digoxin.

Links: https://doi.org/10.1002/tkm2.1419 https://pubmed.ncbi.nlm.nih.gov/32335006/

2. SHAKANZOTO (Kampo, Japan) - rating: REPORTED

Two Japanese teens with orthostatic dysregulation (morning fatigue, dizziness, palpitations, school absence) resolved fully on shakanzoto (peony + licorice), one by day 14, one by day 4. Case-level only; pattern overlaps POTS.

Dose: granules 7.5 g/day split 3x. Effect: days to 2 weeks. Side effects: mild GI upset; licorice warnings under #4.

Link: https://www.jstage.jst.go.jp/article/kampomed/75/2/75_132/_article/

3. TCM ACUPUNCTURE - rating: REPORTED

A 2025 Clinical Pearl describes twice-weekly 40-min acupuncture for POTS (PC6, ST36, BL15, BL20, KI3, SP6, GV20; 2 Hz electroacupuncture), rechecked after 6-8 sessions; patients "frequently report" less fatigue, palpitations and better orthostatic tolerance. An Indonesian 12-session protocol (GV20, PC6, ST36, LR3) targets the same symptoms. No controlled POTS trials, but an RCT review found acupuncture reliably raises vagal tone (HRV) - relevant since POTS is high-sympathetic.

Dose: 2 sessions/week, 15-40 min, 6-12 sessions; benefit after 4-6.

Side effects: minor (soreness, bruising). SAFETY: needling can trigger vasovagal fainting - lie down for treatment, never standing.

Links: https://pmc.ncbi.nlm.nih.gov/articles/PMC12820681/ https://pmc.ncbi.nlm.nih.gov/articles/PMC12820665/ https://pubmed.ncbi.nlm.nih.gov/36494036/ (HRV review)

4. LICORICE ROOT (GANCAO/KANZO) - MODERATE for orthostatic hypotension, NOT tested

in POTS; serious safety limits

Works like weak fludrocortisone: glycyrrhizin blocks 11-beta-HSD2, extending cortisol's salt-water retention, raising blood volume and standing BP. A human study found it helped postural hypotension in diabetic autonomic neuropathy; a review supports it for low-aldosterone orthostatic hypotension - the closest traditional equivalent of the #1 prescription POTS drug, with the same risks.

Dose: 1-2 g root/day traditional; toxicity risk rises above ~100 mg/day glycyrrhizin (crisis documented at 1,000 mg/day extract). Limit trials to 4-6 weeks with BP and potassium checks. Effect: 1-2 weeks.

Side effects: hypertension, low potassium (weakness, cramps, palpitations), leg swelling, alkalosis, headache.

DANGEROUS interactions: fludrocortisone/steroids (documented severe hypokalemia), diuretics, digoxin, laxatives. Avoid in pregnancy, kidney/heart failure, hyperadrenergic POTS.

SAFETY FLAG: severe hypertension and arrhythmias from potassium loss - the one traditional remedy that can directly harm a POTS patient.
https://pmc.ncbi.nlm.nih.gov/articles/PMC7769495/ (licorice + fludrocortisone)

5. ASTRAGALUS / HUANGQI (China) - rating: EARLY, honest negative

Classic TCM "raise the falling yang qi" tonic theorized for orthostatic hypotension; a Chinese RCT (NCT05647473, 10-20 g/day) tested it for that. But the only double-masked crossover study in humans (10 men, 14 days, head-up tilt + lower-body negative pressure) found NO improvement in standing time, BP or heart rate.

Dose in trials: 10-20 g/day (OTC capsules lower). Side effects: mild GI upset; avoid with immunosuppressants, pregnancy.

Link: https://doi.org/10.3357/asem.2193.2008 (negative controlled study)

6. SHENGMAI SAN (China) - rating: EARLY

Ginseng + ophiopogon + schisandra, the classic "Qi and Yin deficiency" formula - the TCM pattern assigned to POTS. Chinese hospital series report shengmai INJECTION improving orthostatic hypotension (49 cases) and vasovagal syncope (60 cases, 90% efficacy) - small, unreplicated, IV-only; no oral POTS trials.
Review: https://lcm.amegroups.org/article/view/4731/html

Dose: 2-3 g granules 2-3x daily. Time: unknown. Side effects: ginseng insomnia/palpitations; avoid with warfarin, MAOIs, uncontrolled hypertension; schisandra with CYP3A4 drugs.

7. AYURVEDA (India) - rating: REPORTED, no studies in POTS

Ayurveda frames POTS-like states as vata aggravation (rapid heart, dizziness) and recommends warm, salty, oily, grounding regimens - matching standard POTS advice (salt, fluids). Ashwagandha and brahmi are often suggested, but zero POTS studies exist, and ashwagandha can LOWER blood pressure and alter thyroid function -

potentially worsening orthostatic intolerance. Such claims are unproven; "cure" pages are hype.

Link: https://www.easyayurveda.com/dysautonomia-ayurveda-understanding/

SOURCE NEEDED: any Ayurvedic clinical study in POTS (none found).

BOTTOM LINE FOR A 32-YEAR-OLD WITH POST-COVID POTS

Kampo and TCM hold the only traditional data touching actual POTS patients, all small. Least-risk first step: acupuncture (vagal-toning evidence, minimal side effects, done lying down). Kampo formulas contain licorice - her doctor must know, especially with fludrocortisone; pure licorice needs potassium monitoring. Astragalus failed its one controlled test. RED FLAGS needing a doctor now: fainting with injury, chest pain, resting HR over 120, BP 160+/100+, severe headache, weakness/cramps (low K+), palpitations with lightheadedness seated.

LINK COUNT: 10 real links, 1 SOURCE NEEDED.

ONGOING TRIALS FOR POTS (recruiting or running as of October 2026)

Searched: ClinicalTrials.gov API (live query for recruiting POTS studies), EU CTIS, ChiCTR and WHO ICTRP via web search. ChiCTR showed no POTS treatment trials. Below are the trials most worth watching for a 32-year-old woman with post-COVID POTS. Statuses verified against the live registry October 2026.

1. SALT CAPSULES - CAlgary SAlt for POTS (CASA-POTS) - Trial NCT05924646

Remedy: sodium chloride (salt). DOSE: 6 g salt in slow-release capsules PLUS 6 g salt in the diet (12 g/day total) vs placebo capsules. University of Calgary, randomized double-blind crossover, 50 participants.

Status: RECRUITING (verified 2026-04). Started 2024. Expected finish: June 2028 (primary), December 2028 (full).

Why it matters: tests the "go high on salt" advice in a blinded way; slow-release capsules may be kinder to the stomach than plain salt.

Side effects/interactions: high salt can raise blood pressure, swelling, headache; risky with kidney disease or if on lithium (salt changes lithium levels); check with your doctor if on blood pressure medicines. Proof for salt approach so far: MODERATE.

Link: https://clinicaltrials.gov/study/NCT05924646

2. SUPERVISED EXERCISE TRAINING - Lund University, Sweden - Trial NCT05554107

Remedy: supervised structured exercise program vs self-guided activity. Randomized crossover, 200 participants (large for POTS).

Status: RECRUITING (verified 2026-08). Expected finish: March 2028 (primary), October 2028.

Side effects/interactions: overdoing exercise can trigger flares; must be built up gradually, ideally recumbent (rowing, cycling, swimming) at first. Proof for exercise in POTS: MODERATE. Link: https://clinicaltrials.gov/study/NCT05554107

3. COMPRESSION GARMENTS + MEDICATIONS (COM-COM-POTS) - Trial NCT04881318

Remedy: waist-high compression tights, worn with and without usual POTS medications; optional abdominal-compression sub-study. Calgary, crossover, 30 participants.

Status: RECRUITING (verified 2026-04). Expected finish: December 2026 - results soon.

Side effects: garments are safe; too-tight wear can cause skin irritation; heat intolerance worse in summer. Proof for compression: MODERATE.

Link: https://clinicaltrials.gov/study/NCT04881318

4. COUNTERPRESSURE MANEUVERS + BIOFEEDBACK - Innsbruck - Trial NCT06936319

Remedy: leg-crossing, muscle-tensing "counter-maneuvers" taught with a lab biofeedback

session vs counseling alone. Randomized, 40 participants.

Status: RECRUITING (verified 2025-03; registry not updated recently). Primary completion Dec 2025 - check for results. Side effects: none expected; free and safe.

Proof: EARLY. Link: https://clinicaltrials.gov/study/NCT06936319

5. BREATHING RETRAINING IN PREGNANCY - Trial NCT07579026

Remedy: diaphragmatic breathing exercises (slow nasal inhale, controlled exhale) in pregnant women with POTS. Pakistan (Riphah Univ.), randomized double-blind, 44 participants. DOSE: daily breathing sessions per protocol.

Status: RECRUITING. Expected finish: September 2026 - results soon. Safe in pregnancy; no known interactions. Proof: EARLY (small, one site).

Link: https://clinicaltrials.gov/study/NCT07579026

6. IV SALINE AND BRAIN FOG (1 L vs 50 mL) - Calgary - Trial NCT05914649

Remedy: 999 mL IV normal saline vs 50 mL (placebo dose), testing neurocognitive (brain fog) improvement. Randomized double-blind crossover, 100 participants.

Status: RECRUITING. Expected finish: December 2030 (dates often move earlier).

Side effects/interactions: IV saline needs a clinic and IV access; risks include vein irritation and, rarely, fluid overload (a concern with heart failure or kidney disease).

Proof: EARLY. Link: https://clinicaltrials.gov/study/NCT05914649

7. 2-HOBA (2-hydroxybenzylamine) FOR IMMUNE ACTIVATION - Vanderbilt - Trial NCT07189936

Remedy: 2-HOBA, a scavenger of reactive "isolevuglandin" molecules tied to persistent immune activation in long-COVID POTS. Investigational capsule (not OTC); dose set by protocol, not published on the registry. Phase 2, randomized, quadruple-blind, 50 participants. Status: RECRUITING (started Dec 2025). Expected finish: June 2029.

Proof: EARLY. Link: https://clinicaltrials.gov/study/NCT07189936

8. MEDICATION CONTEXT (not OTC): PROPRANOLOL vs IVABRADINE - Trial NCT04186286

Calgary crossover study comparing 4-week courses of propranolol, ivabradine and placebo, 20 participants. Still RECRUITING; finish December 2026. Prescription drugs; interactions with blood-pressure and heart-rhythm medicines; do not self-start.

Link: https://clinicaltrials.gov/study/NCT04186286

9. WORTH WATCHING (observational): DEEP PHENOTYPING OF POTS - Trial NCT06292104

UT Southwestern (NHLBI-funded), 350 participants. Not a treatment trial, but it measures blood volume, skin nerves and biomarkers that will shape which remedies fit which patients. RECRUITING, finish January 2029. Link: https://clinicaltrials.gov/study/NCT06292104

JUST FINISHED - RESULTS TO WATCH (context)

RECOVER-AUTONOMIC (NCT06305780, NIH): ivabradine arm finished March 2026 - it lowered standing heart rate but did NOT improve POTS symptoms; the COORDINATED CARE arm (coaching + diet/lifestyle recommendations) DID improve symptoms - the strongest signal yet that structured non-drug care matters. Proof for coordinated care: MODERATE.
Links: https://clinicaltrials.gov/study/NCT06305780 and https://recovercovid.org/news/recover-autonomic-clinical-trial-results-shared-2026-acc-conference

HOW TO JOIN

Search each link above for enrollment contact under "Contacts/Locations". Never join two treatment trials at once, and tell the trial team every supplement you take.
SAFETY NOTE: see a doctor right away for chest pain, fainting with injury, heart racing while lying down, or new shortness of breath.
LINK COUNT: 11 real links, 0 SOURCE NEEDED.

12 - COMPLETED POTS TRIALS WITH NO RESULTS POSTED

Checked ClinicalTrials.gov on 2026-10-21. It lists 54 POTS trials with status COMPLETED; only 17 of 54 have results posted on the registry. The other 37 are finished studies whose data never appeared there - a few were later published in journals (noted below), the rest vanished from public view. That gap is where answers may sit. Doses/timings below are what the trial PROTOCOLS tested - not proven cures.

NO RESULTS ANYWHERE (registry empty, no paper found)

1. BREATHING RETRAINING + AEROBICS (Pakistan). NCT05404672. RCT, 28 women, done Jan 2023: breathing retraining (nasal breathing + controlled pause, 15 min twice daily, later 4x/day) plus cycling/treadmill 30 min, 5 days/ week x 1 month, vs aerobics alone. NO results posted; no paper found. Rating: EARLY. Safe; mild lightheadedness during breath holds; no interactions. A related UK cohort (not this trial) improved breathing-symptom scores: https://pubmed.ncbi.nlm.nih.gov/31743851/
Registry: https://clinicaltrials.gov/study/NCT05404672

2. THERMAL WRIST DEVICE "Embr Wave 2" (Stanford). NCT04943276. Single-arm pilot, 23 adults, done Nov 2023: warming/cooling wrist watch, daily 4 weeks, for heat intolerance.
Registry: https://clinicaltrials.gov/study/NCT04943276
Not on the registry, but an Apr 2026 paper reports improved temperature-related symptoms: https://www.sciencedirect.com/science/article/pii/S1566070226000202
Rating: EARLY (no control). OTC (~$300); safe, no drug interactions; skin irritation possible.

3. AQUATIC (POOL) THERAPY (Cal State Dominguez Hills). NCT07182578. Single-group pilot, done Feb 2026: 3 pool sessions/week, 60 min, 12 weeks.
Registry: https://clinicaltrials.gov/study/NCT07182578
NO results posted; too new, watch for the paper. Water pressure on the legs is a mild compression effect - plausible but unproven. Rating: EARLY. Safe; fainting risk if alone in water - use supervised sessions.

4. NON-INVASIVE VAGUS NERVE STIMULATION FOR LONG COVID + POTS (Mount Sinai). NCT05630040. Sham-controlled crossover RCT, 40 adults, daily 6-week protocol, done Oct 2024. Registry: https://clinicaltrials.gov/study/NCT05630040
NO results posted and no publication found as of Oct 2026 - the biggest miss here, since POTS was a named condition and it was placebo-controlled. Consumer tVNS devices are OTC but the dose used is unknown. Rating: EARLY. Side effects: skin irritation, mild headache; avoid with

pacemakers. A related 24-women pilot (10 days, 30 min twice daily) improved cognition/sleep/fatigue: https://pmc.ncbi.nlm.nih.gov/articles/PMC11097097/

REGISTRY EMPTY, BUT A JOURNAL PAPER APPEARED LATER

5. ABDOMINAL + LEG COMPRESSION (Nationwide Children's). NCT01795469, done 2016, no registry results, but published 2014: abdominal plus leg compression (Zoex garment) cut standing heart-rate rise and symptoms vs none during tilt; abdominal-only worked nearly as well.
Paper: https://www.sciencedirect.com/science/article/pii/S0022347614003424
Registry: https://clinicaltrials.gov/study/NCT01795469
Rating: MODERATE. Dose equivalent: an abdominal binder or waist-high compression shorts is the key piece. Side effects: discomfort, heat. Safe with all medications.

6. EAR-CLIP VAGUS NERVE STIMULATION (U. Oklahoma). NCT03930914, done Dec 2021, NO registry results - but a double-blind sham-controlled RCT was published 2023: 2 months of daily low-level tragus (ear-clip) tVNS improved standing heart rate and HRV vs sham, with a drop in adrenergic autoantibodies and inflammatory cytokines.
Paper: https://pmc.ncbi.nlm.nih.gov/articles/PMC10932945/
Registry: https://clinicaltrials.gov/study/NCT03930914
Rating: MODERATE (small but well-designed). Ear stimulators are sold OTC; this model is investigational. Side effects: ear tingling.

7. SEMI-SUPERVISED REMOTE EXERCISE (Mayo Clinic). NCT04603157, done Feb 2022, NO registry results - but published Aug 2023: a 3-month personalized strength + aerobic program beat standard care for fitness, orthostatic symptoms and exercise tolerance.
Paper: https://pubmed.ncbi.nlm.nih.gov/37598401/
Registry: https://clinicaltrials.gov/study/NCT04603157
Rating: MODERATE. Dose: 3 months, progressive, mostly home-based. Flare risk if progressed too fast; start recumbent.

8. ANTIVIRAL NIRMATRELVIR/RITONAVIR FOR LONG COVID (Karolinska). NCT05823896 ("PROLIFIC"), 219 adults, done Nov 2024, POTS listed as a condition, NO results posted yet. Two sibling trials (STOP-PASC, PAX LC) already found 15 days of nirmatrelvir/ritonavir 300/100 mg twice daily did NOT beat placebo for long-COVID symptoms.
https://jamanetwork.com/journals/jamainternalmedicine/fullarticle/2819901
Registry: https://clinicaltrials.gov/study/NCT05823896
Rating: MODERATE (against benefit). Ritonavir interacts dangerously with statins, blood thinners, some heart-rhythm and BP meds, birth-

control hormones. Prescription-only; never self-source.

WHAT THIS MEANS

The unpublished pile is richest in our exact target areas: breathing retraining, pool therapy, vagus stimulation, compression, home exercise.

"Completed, no results posted" often means negative, boring or never-analyzed data - do not assume hidden cures. But where a paper later appeared (compression, tVNS, exercise) it was positive, so the unpublished ones are worth chasing via the study contacts on each registry page.

SAFETY: all non-prescription except nirmatrelvir. Red flags needing a doctor NOW: fainting, chest pain, racing heart that will not settle, GI bleeding, new leg swelling.

LINK COUNT: 15 real links, 0 SOURCE NEEDED.

Abandoned, Stopped, and Terminated Trials for POTS

This document tracks clinical trials for Postural Orthostatic Tachycardia Syndrome (POTS) that were terminated, withdrawn, or suspended. Understanding why a trial stopped is critical: some fail due to "futility" (the drug did not work), while others stop due to recruitment failures, funding gaps, or external crises (like the COVID-19 pandemic).

1. Efgartigimod (FcRn Antagonist)

Status: TERMINATED (Futility)

Trial IDs: NCT05918978 (Extension), NCT05633407 (Main Phase 2)

Target: Post-COVID-19 POTS (PC-POTS)

Intervention: Efgartigimod IV at a dose of 10 mg/kg.

Findings: The trial was terminated because the Phase 2 results (ALPHA study) showed no clinically meaningful improvement compared to placebo on the Malmö POTS symptom (MaPS) score and COMPASS31.

Proof Rating: STRONG (The failure to meet endpoints in a randomized placebo-controlled trial provides strong evidence against its efficacy for PC-POTS).

Timeline: Treatment period was designed for up to 48 weeks.

Side Effects/Interactions: Generally well-tolerated in other indications, but for PC-POTS, it failed to provide benefit. Potential for increased risk of infections due to lowered IgG levels.

Link: https://clinicaltrials.gov/study/NCT05918978

2. Intravenous Iron Sucrose

Status: TERMINATED / WITHDRAWN (Recruitment Failure)

Trial IDs: NCT01978535, NCT04855266

Target: POTS patients with iron deficiency (even if non-anemic).

Intervention: Intravenous iron sucrose (Venofer) at 5 mg/kg, maximum dose 200 mg, diluted in 0.9% NaCl.

Findings: These trials were stopped primarily due to difficulty in recruiting enough participants. No final efficacy data was published to prove or disprove the hypothesis that correcting iron stores improves cardiovascular indices in POTS.

Proof Rating: EARLY (The hypothesis remains plausible, but these specific trials failed to provide a conclusion).

Timeline: Infusions were typically single-event or short-course.

Side Effects/Interactions: Risk of hypersensitivity/anaphylaxis (rare but serious). Interaction with certain oral antibiotics or antidepressants that inhibit iron absorption (though less relevant for IV).

Link: https://clinicaltrials.gov/study/NCT01978535

3. High Sodium Diet & External Abdominal Compression

Status: TERMINATED (Funding/Recruitment)

Trial ID: NCT01771484

Target: General POTS population.

Intervention:

High Sodium: 300 mEq/day of sodium for 4-5 days.

Device: External automated abdominal binder (compression).

Findings: Terminated because enough volunteers were not recruited before funding was exhausted. It did not provide a definitive verdict on the synergy between extreme sodium loading and automated abdominal compression.

Proof Rating: REPORTED (Based on trial design only; no results published).

Timeline: 4-5 day dietary loading phase.

Side Effects/Interactions: Very high sodium can cause severe edema, hypertension, and is dangerous for those with kidney disease or heart failure.

Link: https://clinicaltrials.gov/study/NCT01771484

4. Northera (Droxidopa) - Specific POTS Studies

Status: TERMINATED (Recruitment/COVID)

Trial ID: NCT02558972

Target: POTS and Postural Vasovagal Syncope (VVS).

Intervention: Droxidopa titrated from 100 mg three times a day (TID) up to 600 mg TID.

Findings: Terminated because adequate numbers could not be recruited, exacerbated by COVID-19 pandemic risks. While Droxidopa is FDA-approved for neurogenic orthostatic hypotension (nOH), its specific efficacy in the general POTS population remains less settled than in nOH.

Proof Rating: MODERATE (Droxidopa works for some orthostatic issues, but this specific POTS trial failed to conclude).

Timeline: Two-week titration and monitoring.

Side Effects/Interactions: Supine hypertension (dangerously high blood pressure when lying down), headache, nausea.

Link: https://clinicaltrials.gov/study/NCT02558972

5. Human Albumin Infusions

Status: WITHDRAWN (Product Support)

Trial ID: NCT03365414

Target: Severe POTS.

Intervention: Intravenous Albumin (Human) 5%, USP.

Findings: The study was withdrawn because the support for the product required for the trial was withdrawn. No clinical data on whether albumin improves blood volume and symptoms in POTS was generated.

Proof Rating: REPORTED (Hypothesis only).

Timeline: Multi-phase administration.

Side Effects/Interactions: Fluid overload, allergic reaction to foreign proteins.

Link: https://clinicaltrials.gov/study/NCT03365414

6. IgPro20 (Subcutaneous IgG)

Status: TERMINATED (Sponsor Decision)

Trial ID: NCT06524739

Target: Post-COVID-19 POTS.

Intervention: IgPro20 (20% liquid polyvalent human IgG) administered subcutaneously.

Findings: Terminated by sponsor decision. The documentation explicitly states this was not for safety reasons, but the trial did not reach completion.

Proof Rating: EARLY (No data provided).

Timeline: 2024-2025 study window.

Side Effects/Interactions: Local injection site reactions, potential systemic immune modulation.

Link: https://clinicaltrials.gov/study/NCT06524739

7. Light Swimming for ME/CFS and POTS

Status: SUSPENDED (Ethics/Safety Concern)

Trial ID: NCT07454395

Target: ME/CFS and related conditions (often including POTS).

Intervention: Light swimming at a self-selected intensity (RPE < 5/10).

Findings: Suspended by the Research Ethics Board following a complaint regarding the safety of exercise in this population (specifically the risk of Post-Exertional Malaise, or PEM). While no participants complained, the board felt the risk of triggering crashes was too high.

Proof Rating: EARLY (Study was suspended before data collection).

Timeline: Baseline tracking followed by light swimming.

Side Effects/Interactions: Risk of Post-Exertional Malaise (PEM), severe fatigue, and symptom exacerbation.

Link: https://clinicaltrials.gov/study/NCT07454395

Summary of Lessons from Stopped Trials

Many POTS trials fail not because the treatment is "wrong," but because the patient population is difficult to recruit or the funding is unstable. However, the failure of Efgartigimod (NCT05633407/NCT05918978) is a significant marker, suggesting that for many with post-COVID POTS, the mechanism may not be purely driven by the IgG-mediated pathways that Efgartigimod targets.

LINK COUNT: 7 real links, 0 SOURCE NEEDED.

14 - RECOVERY AND REMISSION CASE REPORTS (post-COVID POTS)

Documented cases where people with POTS after COVID got substantially better or fully recovered. Bottom line: FULL remission is documented but uncommon; MAJOR improvement within 6-12 months is usual.

1. THE MEDICAL EVIDENCE ON WHO GETS BETTER

Majority improve within months - MODERATE (multiple human cohort studies; observational, not drug trials).

Japanese case series, 32 patients with long-COVID POTS (median age 28, ~5 months follow-up): 94% improved. Standing heart-rate jump fell from a median of 38 bpm to 17.5 bpm; fatigue score improved from 4 to 1; patients able to work or attend school rose from 31% to 69%. Treatment: salt/fluid counseling (10 g salt, 2-3 L water), compression stockings, low-dose bisoprolol (0.625-2.5 mg). Some improvement was spontaneous. Link: https://pmc.ncbi.nlm.nih.gov/articles/PMC10792321/

Mayo Clinic prospective 1-year follow-up (58 POTS patients, pre-COVID era): orthostatic symptoms improved and over one-third no longer met tilt-table criteria for POTS at 1 year. Link: https://www.sciencedirect.com/science/article/pii/S0025619612004697

Chinese pediatric cohort (121 patients, conventional non-drug treatment): cumulative symptom-free rate rose steadily over years, best when POTS was diagnosed and treated early. Link: https://www.frontiersin.org/journals/pediatrics/articles/10.3389/fped.2019.00261/full

2026 characterization study: COVID-triggered POTS cases started with somewhat worse symptoms but showed GREATER improvement over follow-up than non-COVID POTS. Link: https://www.sciencedirect.com/science/article/pii/S1566070225001407

A survey cited in the Japanese series (172 teens, ~5.4 years): only 19% had COMPLETE resolution, 51% improved-but-persistent, 16% intermittent - full recovery happens, but is not the majority. (Same PMC link above.)

2. PUBLISHED REMISSION CASE REPORTS

Loratadine (OTC antihistamine) - complete remission of hyperadrenergic post-COVID POTS - REPORTED (single peer-reviewed case, n=1; Journal of Hypertension 2024, PMID 38526146). A healthy 26-year-old man had 1 year of palpitations, presyncope, dizziness, brain fog and fatigue starting 2 months after mild COVID. Tilt test showed orthostatic tachycardia (85 to ~150 bpm) plus a dangerous standing blood pressure spike (to ~183 mmHg). Beta-blockers WORSENED

his presyncope and were stopped. Because hives/angioedema appeared alongside his orthostatic episodes, doctors suspected mast cell activation syndrome (MCAS) and tried loratadine 10 mg every 12 hours (twice the usual OTC dose). Result: full symptom remission within a few weeks, confirmed by a repeat tilt test 2 weeks in showing normal standing blood pressure and heart rate. Dose: 10 mg twice daily. Time to work: weeks. Side effects: headache, drowsiness, dry mouth. Interactions: adds to sedating meds/alcohol; erythromycin and ketoconazole raise loratadine levels; reduce dose with liver or kidney disease; ask a doctor if pregnant or breastfeeding. SAFETY: do not copy the double dose without a doctor - this applies only to the MCAS-overlap subgroup (rash, flushing, hives with your POTS episodes). Link: https://europepmc.org/article/MED/38526146

Epipharyngeal abrasive therapy (EAT, Japan) - adolescent long-COVID POTS - REPORTED (single peer-reviewed case, Medicine 2025). An adolescent with long-COVID POTS unresponsive to usual treatment recovered enough to attend school normally after EAT - zinc chloride applied to the inflamed nasopharynx by an ENT doctor, a Japanese treatment for chronic epipharyngitis. Not a home remedy. Side effects are mainly local (pain, minor bleeding). Link: https://www.ovid.com/jnls/md-journal/fulltext/10.1097/md.0000000000043333~successful-treatment-of-long-covid-postural-tachycardia

3. PATIENT-REPORTED RECOVERIES (all REPORTED - personal stories, no study)

A woman disabled by long-COVID POTS plus small-fiber neuropathy reports a year-long comeback using pacing and recumbent exercise started at 30 seconds; repeat autonomic testing at Stanford later showed normal standing heart rate and blood pressure. She finished a sprint triathlon. Link: https://medium.com/@kirsten.lundgren/healing-from-long-covid-post-vaccine-pots-small-fiber-neuropathy-its-possible-a5716b5b03a7

Newsweek first-person story: a teen girl's POTS resolved about two years after onset during a slower-paced, low-stress year abroad; symptom-free 10 years since. Link: https://www.newsweek.com/my-sudden-symptoms-doctors-stumped-rare-diagnosis-i-1912336

A lifelong runner with POTS rebuilt from 3 minutes of exercise tolerance to running 20 minutes three times a week over ~4 years of gradual walk/run and strength training; her POTS improved as conditioning grew. Link: https://dropitlikeitspots.com/running-with-pots/

4. WHAT THE RECOVERY CASES HAVE IN COMMON

Gradual, never-sudden reconditioning; consistent fluids/electrolytes/salt; months-to-years timelines (weeks only in the loratadine MCAS case); treating overlapping problems (sleep, anxiety, mast-cell symptoms); early diagnosis.

No case report found of a supplement or single food curing post-COVID POTS. Reddit long-haulers also carry multiple self-reported full recoveries from post-COVID POTS, typically within about a year. Link: https://www.reddit.com/r/covidlonghaulers/comments/vquigv/had_anyone_fully_recovered_from_pots/

5. RED FLAGS - SEE A DOCTOR NOW

Fainting with injury or while driving; chest pain or breathlessness on standing; heart rate sustained over 150 bpm; standing blood pressure over 180 mmHg (as in the loratadine case, this is dangerous); new hives, facial swelling or throat tightness with lightheadedness (possible MCAS/anaphylaxis - call emergency services); POTS with fever, weight loss, or new neurological signs. Never stop prescribed heart or BP medication without your doctor.

LINK COUNT: 10 real links, 0 SOURCE NEEDED.

15 - OTC PRODUCTS FOR POST-COVID POTS: CHECKED vs HYPE

What store-bought products actually deliver for POTS, versus what is marketing.
Ratings apply to the PRODUCT as sold (the generic salt/fluid/compression therapy behind them is covered in files 03 and 05).

CHECKED - worth considering

1. ORAL REHYDRATION POWDERS (NormaLyte type, ~1,700-2,300 mg sodium/pack) - MODERATE. A real randomized trial in pediatric POTS found an oral rehydration solution worked about as well as IV saline for orthostatic intolerance (https://www.sciencedirect.com/science/article/pii/S0022347619308935). DOSE: 1-2 packs/day mixed in ~500 mL water each, spread morning and midday. TIMING: effect on standing heart rate within 30-60 minutes of drinking; fuller benefit after 1-2 weeks of consistent use. SIDE EFFECTS: bloating, nausea if chugged; loose stools. INTERACTIONS: dangerous with kidney disease, heart failure, or fluid-restricted; potassium content matters on kidney meds or potassium-sparing diuretics. Manufacturer page is marketing, but the ORS category itself has a human POTS trial: (https://normalyte.com/pages/journal-of-pediatric-medicine-clinical-study-of-iv-normal-saline-vs-oral-rehydration-salts)

2. MEDICAL-GRADE COMPRESSION (20-30 mmHg socks; abdominal binders) - MODERATE. Two independent human crossover studies: waist-high compression cut standing HR from 109 to ~92 bpm (https://www.jacc.org/doi/10.1016/j.jacc.2020.11.040), and commercial abdominal-only garments lowered standing HR (118 to 103 bpm) and symptoms, with benefit persisting after hours of wear (https://pubmed.ncbi.nlm.nih.gov/41338488/). Abdominal compression beat socks-only in both studies. A 330-patient survey found most patients still use leg socks even though abdominal/shapewear may work better (https://www.mdpi.com/2077-0383/13/23/7304). DOSE: 20-40 mmHg, put on before getting up, wear through upright hours. TIMING: same-day effect. SIDE EFFECTS: heat, discomfort, skin irritation. INTERACTIONS: none drug-based, but too-tight garments can numb legs. Brands like CEP or PotsSocks are store-level choices (https://www.potssocks.com/) - the therapy is proven, the brand is not.

3. HIGH-SODIUM POWDERS: LMNT (~1,000 mg sodium/pack, sugar-free) vs Pedialyte - REPORTED as products (therapy behind them is proven). LMNT costs ~$1.50 per 1,000 mg sodium; Pedialyte Classic ~$5.08 per 1,000 mg and only ~1,080 mg/L - too dilute and too costly for a POTS salt target (3,000-10,000 mg/day). Label comparison: (https://sportingdrinks.com/pedialyte-vs-lmnt/). DOSE: 1-3 packs/day in 16 oz water. TIMING: 30-60 min before long upright periods. SIDE EFFECTS: nausea if taken without enough water. INTERACTIONS: same kidney/heart-failure cautions; skip caffeinated LMNT flavors if palpitations.

Dietitian guide with a free DIY recipe (1/2 tsp salt in 16-20 oz water): (https://eds.clinic/articles/electrolyte-supplements-for-pots-eds-and-mcas-guide)

4. SODIUM CHLORIDE TABLETS (Nutricost, Vitamatic, ~1,000 mg NaCl = ~390 mg sodium/tablet) - REPORTED. Cheapest per mg of sodium (about 5 cents per 390 mg sodium) but easy to overdose; label: (https://www.walmart.com/c/kp/sodium-chloride-tablet). DOSE: 1-2 tablets with a full glass of water, up to 3-4 times daily within the doctor-set sodium cap. TIMING: same-day. SIDE EFFECTS: stomach irritation, vomiting if taken dry. INTERACTIONS: never with salt-sensitive hypertension, kidney disease, or corticosteroids without doctor sign-off.

5. NICOTINAMIDE RIBOSIDE (NR, NAD+ precursor) for long-COVID brain fog - EARLY. A double-blind trial (2,000 mg/day, 12 weeks) in long-COVID raised NAD+ and showed signals on cognition/fatigue, but it was not a POTS trial and effects were modest: (https://www.thelancet.com/journals/eclinm/article/PIIS2589-5370(25)00567-X/fulltext). DOSE: 1,000-2,000 mg/day morning. TIMING: 6-12 weeks trial. SIDE EFFECTS: flushing, nausea, mostly mild. INTERACTIONS: theoretical with blood thinners. Expensive (~$50/month) for an early-evidence product.

REJECTED - hype, no proof, or bad value

"ADRENAL COCKTAILS" (orange juice + coconut water + cream of tartar + salt): Cleveland Clinic states there is no evidence "adrenal fatigue" exists or that these drinks support the adrenal glands; the potassium load is risky in kidney disease: (https://health.clevelandclinic.org/what-to-know-about-adrenal-cocktails). REPORTED - and the mechanism claim is false.

COCONUT water sold as a "POTS hydration cure": no POTS trials found; it is low-sodium, high-potassium - the opposite of what POTS salt loading needs. REPORTED.

PEDIALYTE CLASSIC as a POTS loader: ~$5 per 1,000 mg sodium with 13 g sugar per liter - you would need 3+ liters and 40 g sugar daily to reach 3,000 mg. Bad value, not dangerous. REPORTED.

"POTS cure" pills and Amazon supplements with review-inflated claims: the FDA/AARP scam pattern list applies - no POTS product is FDA-approved (https://www.aarp.org/money/scams-fraud/supplements/). REPORTED.

SAFETY: any product causing fainting, chest pain, racing heart while seated, or BP spikes needs a doctor right away. All high-sodium products are unsafe with kidney failure, heart failure, or uncontrolled high blood pressure. Prescription context only: doctors often add beta-blockers, ivabradine, midodrine or fludrocortisone on top of salt/fluid/compression.

LINK COUNT: 12 real links, 0 SOURCE NEEDED.

16 - COMBINATION APPROACHES (things tested TOGETHER in people)

Multi-part treatments studied as a COMBINATION in humans. Single remedies are in files 03-08. Combinations often beat single treatments head-to-head.

1. Exercise training + volume expansion (salt and fluid) - PROOF: MODERATE

The Dallas combination. JACC study, 19 POTS patients: a small heart plus low blood volume drove the syndrome; ~3 months of structured exercise (with salt and fluid loading) improved or resolved it in most.

Dose: exercise built over ~3 months, 4-5x/week, recumbent first (rower, recumbent bike, swimming) before upright work, plus ~2-3 L fluid and ~10 g sodium/day.

Timing: HR benefits in 4-6 weeks; full effect ~3 months. Side effects: flare if you progress too fast; early upright exercise can trigger fainting.

Interactions: high salt unsafe with hypertension, kidney disease, heart failure; beta-blockers blunt exercise heart rate (use perceived effort).

https://www.jacc.org/doi/10.1016/j.jacc.2010.02.043

2. Exercise training vs propranolol, same patients - PROOF: MODERATE

Vanderbilt: 19 POTS patients, 4 weeks double-blind propranolol (20-80 mg) or placebo, then 3 months exercise training in all. Exercise was superior for upright BP control, kidney-hormone responses and quality of life; the pill alone was not. Lesson: pair short-term beta-blockers with reconditioning, don't replace exercise with a drug. (Doctors also use midodrine, ivabradine, pyridostigmine.)

Side effects/interactions: propranolol can drop standing BP and worsen dizziness; interacts with asthma, diabetes meds, pregnancy.

https://pmc.ncbi.nlm.nih.gov/articles/PMC3142863/

3. Abdominal compression + propranolol - clearest "combo wins" trial - PROOF: MODERATE

Vanderbilt, 18 POTS patients, placebo-controlled crossover (NCT00262470). Abdominal binder at 40 mm Hg, applied ~2 min before standing, ALONE did nothing. Propranolol 20 mg alone lowered standing HR (81 vs 98 bpm) but dropped standing BP (93 vs 100 mm Hg) and didn't help symptoms. The COMBINATION kept the HR benefit, prevented the BP drop (98 vs 93 mm Hg, P=0.029) and cut symptom burden (-6 vs -1 points, P=0.041).

Dose: binder ~40 mm Hg before/while standing; propranolol 20 mg (prescription - context only). Timing: minutes (binder), ~2 h (propranolol). Side effects: too-tight binders cause discomfort, reflux. Interactions: beta-blockers.

https://pmc.ncbi.nlm.nih.gov/articles/PMC7660715/

4. PULSE: online group exercise + behavioural support + home recumbent bike - PROOF: EARLY

UK feasibility RCT (ISRCTN45323485; 44 adults, 98% female, mean age 30): online consult, 12 weeks supervised live online group exercise, 6 weeks behavioural support, recumbent bike + bands at home. Attendance 15 of 18 sessions median. COMPASS-31 dysautonomia fell ~6 points at 4 months (usual care +1) and ~9 at 7 months (usual care -2); EQ-5D VAS +12. Not powered to prove effect - but safe, no serious adverse events. Side effects: short-term post-exercise flares; some dropped out as "too unwell".

https://bmjopen.bmj.com/content/15/2/e090197

5. Water + salt together (oral rehydration solution) vs IV saline - PROOF: MODERATE

Controlled study (10 POTS patients, 15 controls): on separate days, 1 L oral ORS or 1 L IV saline over 30 min. BOTH prolonged time to fainting under orthostatic stress ($P<0.05$ oral, $P<0.001$ IV) - the cheap home version works.

Acute dose: ~500 mL water within ~15-30 min; maintenance 2-3 L fluid + 10-12 g sodium daily (details in file 03). Side effects: bloating, nausea if chugged.

Interactions: salt unsafe with hypertension, heart or kidney failure - ask a doctor first. Long-term IV saline carries line infection/clot risks.

https://www.jpeds.com/article/S0022-3476(19)30893-5/fulltext

6. Tailored endurance + strength training with weekly physiotherapist support (post-COVID POTS) - PROOF: EARLY

Swedish feasibility study (Sci Rep 2024): 26 post-COVID POTS patients (81% female, median 41), 12 weeks of tailored endurance AND strength training plus weekly physiotherapist support. 76% adherence, 96% completion, safe. Symptoms improved (VOSS 41.4 to 32.9; PHQ-9 12.5 to 9; walk +19 m). The tailoring + weekly human support may be the active ingredient.

https://www.nature.com/articles/s41598-024-71055-5

7. Group exercise + psychological support (REGAIN, post-COVID, not POTS-only) - PROOF: MODERATE (for post-COVID condition)

Multicenter RCT, n=585 (ISRCTN11466448): 8 weeks of weekly live online supervised group exercise + 6 group psychological sessions improved quality of life at 3 AND 12 months vs usual care. Broad post-COVID population, not POTS-only - apply with caution.

https://doi.org/10.1136/bmj-2023-076506

8. Creatine + glycerol + water hyperhydration stack - PROOF: EARLY

Lab study, 19 HEALTHY volunteers (not POTS): 20 g creatine + 2 g/kg glycerol + 2 L water daily for 6 days raised tilt BP (+11 mmHg systolic) and cut

presyncopal events. No POTS trial yet - a lead, not a recommendation. Side effects: weight gain, GI upset, headache; kidney patients must avoid creatine. https://pubmed.ncbi.nlm.nih.gov/19142656/

9. Multimodal conventional care over years - PROOF: EARLY

Mayo Clinic follow-up of children/adolescents on the standard multimodal package (fluids, salt, exercise, sleep ± medication): outcomes benign, symptom-free rate rose steadily over time - the whole stack, patiently applied, leads to remission in a growing share. Observational. https://pmc.ncbi.nlm.nih.gov/articles/PMC6610301/

NOT FOUND

No trial tested the full "everything stack" as one package vs placebo - evidence is piece-by-piece.

SAFETY - SEE A DOCTOR RIGHT AWAY IF:

fainting with injury or while driving; chest pain; irregular or resting palpitations; resting HR over ~120; BP over 160/100 or under 90/60 with symptoms; new severe headache, vision change or trouble speaking; fainting during exercise. High salt can dangerously raise BP in hypertension, kidney or heart failure.

17 - OWN LEADS (worth chasing, not covered by any other file)

Leads found while researching this mission that do not fit files 01-16: new mechanisms, cheap targeting tools, prognosis facts, and non-drug directions. Ratings are honest; most are EARLY or REPORTED.

LEAD 1 - THE GUT MICROBIOME (post-COVID specific) - EARLY

A 2024 controlled study (24 POTS patients, 18 post-COVID, vs 21 matched controls) found gut-microbiota changes in POTS: GI symptoms correlated with altered flora, with reduced microbial diversity and altered short-chain-fatty-acid patterns. This is the first human evidence that the gut bug mix differs in post-COVID POTS - a mechanism distinct from diet (file 04): https://www.nature.com/articles/s41598-024-53784-9

HOW TO CHASE IT: no probiotic has been tested in POTS (that is exactly why it is a lead, not a remedy). Reasonable low-risk self-experiment: a standard multi-strain probiotic (e.g. Lactobacillus/Bifidobacterium blend, label dose, 4-8 week trial) plus fermented food daily. Time to judge: 4-8 weeks. Side effects: bloating, gas the first weeks; rare infection risk if severely immunocompromised. Interactions: none dangerous known; may shift warfarin response slightly - tell the doctor if on blood thinners.

LEAD 2 - HUPERZINE A: the OTC cousin of a proven POTS drug - REPORTED for POTS

The prescription drug pyridostigmine lowered standing heart rate from 119 to 104 bpm and beat placebo in a randomized crossover trial in 17 POTS patients (MODERATE - it works by boosting acetylcholine, shifting autonomic balance): https://www.ahajournals.org/doi/full/10.1161/circulationaha.104.497594

Huperzine A (Chinese club-moss extract, sold OTC as a memory supplement) is the same kind of acetylcholinesterase inhibitor, with meta-analyses of 20 randomized trials in other diseases: https://pubmed.ncbi.nlm.nih.gov/24086396/

HONEST CAVEAT: ZERO trials of huperzine A in POTS exist. This is a lead only.

IF her doctor agrees to a supervised trial: 50-100 mcg once daily with food, morning; effect on alertness within 1-2 hours if any. Side effects: nausea, GI cramps, sweating, drooling, dizziness, vivid dreams. DANGER: can slow the heart rate (bradycardia) and trigger rhythm problems; worsens asthma, ulcers, urinary obstruction. INTERACTIONS: never combine with pyridostigmine, donepezil, beta-blockers, digoxin, or other cholinergic drugs; avoid in pregnancy (no safety data). This is the one supplement in this file that can genuinely harm - do not try it without a doctor who knows her ECG.

LEAD 3 - CHEAP TESTS THAT PREDICT WHO RESPONDS TO SALT/FLUIDS - EARLY

Chinese pediatric POTS research found responders to oral rehydration salts can be predicted in advance: 24-hour urine sodium under 124 mmol (sensitivity 77%,

specificity 93%), low BMI, and high MCHC (a red-blood-cell index on any CBC).
A 24-hour urine jug plus a basic CBC are cheap, and can tell HER whether volume expansion (file 03) is likely to work before weeks of guessing: https://doi.org/10.1002/pdi3.2509
Done in children only - adults untested. Not a remedy: a targeting tool.
Dose/timing: none; test once, repeat after 8-12 weeks of treatment. Side effects: none. Interactions: none.
LEAD 4 - THE PROGNOSIS LEAD: post-COVID POTS often improves by itself - EARLY
A clinic series of long-COVID POTS patients found 94% improved within a median 159 days: https://pmc.ncbi.nlm.nih.gov/articles/PMC10792321/
Single-center, non-randomized, treatments not standardized - so treat the number with caution. PRACTICAL MEANING: at ~6 months in, she is inside the window where many improve; track a simple daily 10-minute stand test (heart rate lying vs standing + symptom score) so a real trajectory is visible.
Improvement is likeliest with the fundamentals (files 03-06) in place.
LEAD 5 - BRAIN-RETRAINING (acoustic neurotechnology) - EARLY
7 adolescents with tilt-confirmed POTS used HIRREM (closed-loop acoustic stimulation that mirrors brain electrical activity back as sound): heart-rate variability rose 51%, baroreflex sensitivity 65%, symptoms trended down, and all 4 who had been on fludrocortisone were able to stop it. Open-label, tiny: https://pmc.ncbi.nlm.nih.gov/articles/PMC4751215/
Clinic-delivered, not OTC - a research lead. A home cousin is slow-paced breathing / HRV-biofeedback apps (6 breaths/min, 10-15 min twice daily; effect if any in 2-4 weeks; side effects none; harmless): no POTS trials, REPORTED.
Interactions: none. Pregnancy/kidney/heart-meds: no restrictions for the breathing version.
LEAD 6 - STELLATE GANGLION BLOCK - watch, do not buy - EARLY
Injection (not natural, not OTC - chased here only because it targets the autonomic ganglion driving the standing heart-rate response). A 41-patient cohort reported 86% relief of long-COVID symptoms including tachycardia and dizziness: https://pmc.ncbi.nlm.nih.gov/articles/PMC10498998/ ; a 2024 retrospective review found sustained symptom reduction in most patients: https://www.sciencedirect.com/science/article/pii/S1566070224000493
NIH's RECOVER-TLC trial is now testing bilateral SGB against placebo in long COVID - the results will answer whether this is real: https://recover-tlc.org/documents/RECOVER-TLC_SGB_Protocol_Synopsis_For_Public_Comment.pdf
Risks: temporary droopy eyelid/hoarseness, infection, very rarely carotid or vertebral artery injury. Wait for the trial results or join the trial.
RED FLAGS (whole mission): fainting with injury, chest pain, resting heart

rate over 120, new irregular heartbeat, blood pressure over 180/120, leg swelling with breathlessness - doctor same day. Lead 2 additionally: any slow pulse, dizziness or palpitations on huperzine A means stop immediately.
LINK COUNT: 9 real links, 0 SOURCE NEEDED.

Claude's notes

These are my comments, not part of Hermes's research. The text above is Hermes's work, copied word for word - I only changed the layout and added this page.

Points worth flagging:

High salt (8-12 g/day) is central to this plan but is not safe for everyone. Hermes says this too: it is off-limits with kidney disease, heart failure, uncontrolled high blood pressure, or in pregnancy. A doctor should OK the salt target before it is used.

Iron should only be taken if a blood test (ferritin) confirms it is low. Taking iron without a proven deficiency can be harmful.

One item in the research (huperzine A, file 17) can genuinely slow the heart and interact with common medicines. Hermes flags it as dangerous; I agree - never try it without a doctor who knows your ECG.

The loratadine remission case (file 14) is one single person. Do not copy the double dose without a doctor.

Ratings in the reports (STRONG / MODERATE / EARLY / REPORTED) are Hermes's honest labels for how good the evidence is. 'REPORTED' or 'EARLY' means mostly unproven.

None of this is medical advice, and nothing here is a proven cure. Please check anything you want to try with your doctor first - especially anything involving salt, supplements, or changes to prescribed medicines.