

CHAPTER 5

Section 5.3

1. (a) From the given end points $(0, 0)$, $(2, 2)$ it follows that we can represent the curve C in the form $y = x$, $0 \leq x \leq 2$. Hence, by (5.6) we find

$$\int_{(0,0)}^{(2,2)} y^2 dx = \int_0^2 x^2 dx = \left. \frac{x^3}{3} \right|_0^2 = \frac{8}{3}$$

- (b) Given the end points $(2, 1)$, $(1, 2)$ we will parameterise the curve C according to: $x = 2 - t$, $y = 1 + t$, $0 \leq t \leq 1$. Then by (5.4) we find

$$\int_{(2,1)}^{(1,2)} y dx = - \int_0^1 (1 + t) dt = - \left[t + \frac{t^2}{2} \right]_0^1 = -\frac{3}{2}$$

- (c) Given the end points $(1, 1)$, $(2, 1)$ we will parameterise the curve C according to $x = 1 + t$, $y = 1$, $0 \leq t \leq 1$. Then by (5.5) we find

$$\int_{(1,1)}^{(2,1)} x dy = \int_0^1 (1 + t) (0) dt = 0$$

2. (a) Let us represent the curve $C : x = \sqrt{1 - y^2}$ in the form $x = \cos t$, $y = \sin t$, $-\pi/2 \leq t \leq \pi/2$. Then by (5.4) and (5.5)

$$\begin{aligned} \int_{(0,-1)}^{(0,1)} y^2 dx + x^2 dy &= \int_{-\pi/2}^{\pi/2} -\sin^3 t dt + \cos^3 t dt \\ &= \int_{-\pi/2}^{\pi/2} -(1 - \cos^2 t) \sin t + (1 - \sin^2 t) \cos t dt \\ &= \left[\cos t - \frac{\cos^3 t}{3} + \sin t - \frac{\sin^3 t}{3} \right]_{-\pi/2}^{\pi/2} = \frac{4}{3} \end{aligned}$$

- (b) Let C be the parabola $y = x^2$. Then by (5.6) and (5.7) we find

$$\int_{(0,0)}^{(2,4)} y dx + x dy = \int_0^2 (x^2 + 2x^2) dx = \left[\frac{x^3}{3} + \frac{2}{3}x^3 \right]_0^2 = 8$$

- (c) Let C be the curve $x = \cos^3 t$, $y = \sin^3 t$, $0 \leq t \leq \pi/2$ and let us use the substitution $u = \tan^3 t$. Then by (5.4) and (5.5) we can rewrite the integral as

$$\begin{aligned} \int_{(1,0)}^{(0,1)} \frac{y dx - x dy}{x^2 + y^2} &= -3 \int_0^{\pi/2} \frac{\sin^4 t \cos^2 t + \sin^2 t \cos^4 t}{\cos^6 t + \sin^6 t} dt = \int_0^{\pi/2} \frac{-3 \sin^2 t \cos^2 t}{\cos^6 t + \sin^6 t} dt \\ &= - \int_0^{\infty} \frac{\cos^6 t}{\cos^6 t + \sin^6 t} du = - \int_0^{\infty} \frac{du}{1 + u^2} = \lim_{b \rightarrow \infty} - \int_0^b \frac{du}{1 + u^2} \\ &= \lim_{b \rightarrow \infty} -\tan^{-1} u \Big|_0^b = \lim_{b \rightarrow \infty} -\tan^{-1} b = -\frac{\pi}{2} \end{aligned}$$

3. (a) Let C be the square with vertices $(1, 1)$, $(-1, 1)$, $(-1, -1)$, $(1, -1)$. Then the integral

$$\oint_C y^2 dx + xy dy$$

can be evaluated by computing the sum of the four integrals

$$\underbrace{\int_{(1,1)}^{(-1,1)} y^2 dx}_{dy=0} \quad \underbrace{\int_{(-1,1)}^{(-1,-1)} xy dy}_{dx=0} \quad \underbrace{\int_{(-1,-1)}^{(1,-1)} y^2 dx}_{dy=0} \quad \underbrace{\int_{(1,-1)}^{(1,1)} xy dy}_{dx=0}$$

Hence,

$$\begin{aligned} \oint_C y^2 dx + xy dy &= \int_1^{-1} dx - \int_1^{-1} y dy + \int_{-1}^1 dx + \int_{-1}^1 y dy \\ &= x \Big|_1^{-1} - \frac{y^2}{2} \Big|_1^{-1} + x \Big|_{-1}^1 + \frac{y^2}{2} \Big|_{-1}^1 = 0 \end{aligned}$$

- (b) Let C be the circle $x^2 + y^2 = 1$. Using the parameterization $x = \cos t$, $y = \sin t$ where $0 \leq t \leq 2\pi$, then by (5.4) and (5.5) the integral

$$\oint_C y dx - x dy$$

may be written as

$$\begin{aligned} \oint_C y dx - x dy &= \int_0^{2\pi} -\sin^2 t dt - \cos^2 t dt = - \int_0^{2\pi} (\sin^2 t + \cos^2 t) dt = - \int_0^{2\pi} dt \\ &= -2\pi \end{aligned}$$

- (c) Let C be the triangle with vertices $(0, 0)$, $(1, 0)$, $(1, 1)$. Then the integral

$$\oint_C x^2 y^2 dx - xy^3 dy$$

can be evaluated by computing the sum of the three integrals

$$\underbrace{\int_{(0,0)}^{(1,0)} x^2 y^2 dx}_{dy=0} = 0 \quad \underbrace{- \int_{(1,0)}^{(1,1)} xy^3 dy}_{dx=0} \quad \int_{(1,1)}^{(0,0)} x^2 y^2 dx - xy^3 dy$$

Hence,

$$\begin{aligned} \oint_C x^2 y^2 dx - xy^3 dy &= - \int_0^1 y^3 dy + \int_0^1 x^4 dx - \int_0^1 y^4 dy \\ &= - \frac{y^4}{4} \Big|_0^1 + \frac{x^5}{5} \Big|_0^1 - \frac{y^5}{5} \Big|_0^1 = -\frac{1}{4} \end{aligned}$$

4. (a) Let C be the circle $x^2 + y^2 = 4$. Then using the parametrisation $x = 4 \cos t$, $y = 4 \sin t$, where $0 \leq t \leq 2\pi$ and (5.12) the integral

$$\oint_C (x^2 - y^2) ds$$

may be written as

$$\oint_C (x^2 - y^2) ds = 64 \int_0^{2\pi} (\cos^2 t - \sin^2 t) dt = 64 \int_0^{2\pi} \cos 2t dt = 32 \sin 2t \Big|_0^{2\pi} = 0$$

- (b) Let C be the line $y = x$ with endpoints $(0, 0)$, $(1, 1)$. Then by (5.14) the integral

$$\int_{(0,0)}^{(1,1)} x ds$$

may be written as

$$\int_{(0,0)}^{(1,1)} x ds = \sqrt{2} \int_0^1 x dx = \frac{\sqrt{2}}{2} x^2 \Big|_0^1 = \frac{1}{\sqrt{2}}$$

- (c) Let C be the parabola $y = x^2$ with endpoints $(0, 0)$, $(1, 1)$. Then by (5.14) and using the substitution $x = (1/2) \tan u$, such that $dx = (1/2) \sec^2 u du$ the integral

$$\int_{(0,0)}^{(1,1)} ds$$

may be written as

$$\int_{(0,0)}^{(1,1)} ds = \int_0^1 \sqrt{1 + 4x^2} dx = \frac{1}{2} \int_0^{\tan^{-1} 2} \sec^3 u du$$

In order to solve the integral on the right hand side, let us solve the indefinite integral

$$\begin{aligned} \int \sec^3 x dx &= \int_0^x \sec^2 x \sec x dx = \sec x \tan x - \int \sec x \tan^2 x dx + C \\ &= \sec x \tan x - \int \sec x (\sec^2 x - 1) dx + C \\ &= \sec x \tan x - \int \sec^3 x dx + \int \sec x dx + C \end{aligned}$$

Adding the term $\int \sec^3 x dx$ to both sides and dividing by two then gives

$$\begin{aligned} \int \sec^3 x dx &= \frac{1}{2} \sec x \tan x + \frac{1}{2} \int \sec x dx + C \\ &= \frac{1}{2} \sec x \tan x + \frac{1}{2} \ln |\sec x + \tan x| + C \end{aligned}$$

Substituting in the original equation then gives

$$\begin{aligned}
\int_{(0,0)}^{(1,1)} ds &= \int_0^1 \sqrt{1+4x^2} dx = \frac{1}{2} \int_0^{\tan^{-1} 2} \sec^3 u du \\
&= \frac{1}{4} \sec u \tan u \Big|_0^{\tan^{-1} 2} + \frac{1}{4} \ln |\sec u + \tan u| \Big|_0^{\tan^{-1} 2} \\
&= \frac{\sqrt{5}}{2} + \frac{\ln(\sqrt{5}+2)}{4}
\end{aligned}$$

5. Let a path $x = \phi(t)$, $y = \psi(t)$, $h \leq t \leq k$, where x and y are continuous and have continuous derivatives for $h \leq t \leq k$ like (5.1) be given. Next, let us make a change of parameter by the equation $t = g(\tau)$, $\alpha \leq \tau \leq \beta$, where $g'(\tau)$ is continuous and positive in the interval and $g(\alpha) = h$, $g(\beta) = k$. Then by (5.4) the line integral $\int_C f(x, y) dx$ on the path $x = \phi(g(\tau))$, $y = \psi(g(\tau))$, such that $dx = (d/d\tau)\phi(g(\tau)) d\tau$, is given by

$$\begin{aligned}
\int_C f(x, y) dx &= \int_\alpha^\beta f[\phi(g(\tau)), \psi(g(\tau))] \frac{d}{d\tau} \phi(g(\tau)) d\tau \\
&= \int_\alpha^\beta f[\phi(g(\tau)), \psi(g(\tau))] \frac{d\phi}{dt} \frac{d}{d\tau} g(\tau) d\tau \\
&= \int_h^k f[\phi(t), \psi(t)] \frac{d\phi}{dt} \frac{dt}{d\tau} d\tau = \int_h^k f[\phi(t), \psi(t)] \phi'(t) dt
\end{aligned}$$

6. (a) Using (a), the integral $\int P dx + Q dy$ along the path $C \rightarrow ABFG$ may be approximated as

$$\begin{aligned}
\int_C P dx + Q dy &\sim \left[\frac{1}{2} (0+3) \cdot 1 + \frac{1}{2} (1+2) \cdot 0 \right] + \left[\frac{1}{2} (3+0) \cdot 0 + \frac{1}{2} (2+4) \cdot 1 \right] \\
&\quad + \left[\frac{1}{2} (0+5) \cdot 1 + \frac{1}{2} (4+6) \cdot 0 \right] = 7
\end{aligned}$$

- (b) Using (a), the integral $\int P dx + Q dy$ along the path $C \rightarrow AFGKH$ may be approximated as

$$\begin{aligned}
\int_C P dx + Q dy &\sim \left[\frac{1}{2} (0+0) \cdot 1 + \frac{1}{2} (1+4) \cdot 1 \right] + \left[\frac{1}{2} (0+5) \cdot 1 + \frac{1}{2} (4+6) \cdot 0 \right] \\
&\quad + \left[\frac{1}{2} (5+0) \cdot 0 + \frac{1}{2} (6+9) \cdot 1 \right] + \left[\frac{1}{2} (0+2) \cdot 1 + \frac{1}{2} (9+8) \cdot -1 \right] \\
&= 5
\end{aligned}$$

- (c) Using (a), the integral $\int P dx + Q dy$ along the path $C \rightarrow ABCDHLSONMIEA$ may be approximated as

$$\begin{aligned}
\int_C P dx + Q dy &\sim \left[\frac{1}{2} (0 + 3) \cdot 1 + \frac{1}{2} (1 + 2) \cdot 0 \right] + \left[\frac{1}{2} (3 + 8) \cdot 1 + \frac{1}{2} (2 + 3) \cdot 0 \right] \\
&+ \left[\frac{1}{2} (8 + 5) \cdot 1 + \frac{1}{2} (3 + 4) \cdot 0 \right] + \left[\frac{1}{2} (5 + 2) \cdot 0 + \frac{1}{2} (4 + 8) \cdot 1 \right] \\
&+ \left[\frac{1}{2} (2 + 1) \cdot 0 + \frac{1}{2} (8 + 2) \cdot 1 \right] + \left[\frac{1}{2} (1 + 4) \cdot 0 + \frac{1}{2} (2 + 6) \cdot 1 \right] \\
&+ \left[\frac{1}{2} (4 + 3) \cdot -1 + \frac{1}{2} (6 + 2) \cdot 0 \right] + \left[\frac{1}{2} (3 + 7) \cdot -1 + \frac{1}{2} (2 + 8) \cdot 0 \right] \\
&+ \left[\frac{1}{2} (7 + 2) \cdot -1 + \frac{1}{2} (8 + 4) \cdot 0 \right] + \left[\frac{1}{2} (2 + 8) \cdot 0 + \frac{1}{2} (4 + 3) \cdot -1 \right] \\
&+ \left[\frac{1}{2} (8 + 3) \cdot 0 + \frac{1}{2} (3 + 2) \cdot -1 \right] + \left[\frac{1}{2} (3 + 0) \cdot 0 + \frac{1}{2} (2 + 1) \cdot -1 \right] \\
&= 8
\end{aligned}$$

- (d) Using (a), the integral $\int P dx + Q dy$ along the path $C \rightarrow AFJNMIJFA$ may be approximated as

$$\begin{aligned}
\int_C P dx + Q dy &\sim \left[\frac{1}{2} (0 + 0) \cdot 1 + \frac{1}{2} (4 + 1) \cdot 1 \right] + \left[\frac{1}{2} (0 + 5) \cdot 0 + \frac{1}{2} (4 + 6) \cdot 1 \right] \\
&+ \left[\frac{1}{2} (5 + 7) \cdot 0 + \frac{1}{2} (6 + 8) \cdot 1 \right] + \left[\frac{1}{2} (7 + 2) \cdot -1 + \frac{1}{2} (8 + 4) \cdot 0 \right] \\
&+ \left[\frac{1}{2} (2 + 8) \cdot 0 + \frac{1}{2} (4 + 3) \cdot -1 \right] + \left[\frac{1}{2} (8 + 5) \cdot 1 + \frac{1}{2} (3 + 6) \cdot 0 \right] \\
&+ \left[\frac{1}{2} (5 + 0) \cdot 0 + \frac{1}{2} (6 + 4) \cdot -1 \right] + \left[\frac{1}{2} (0 + 0) \cdot -1 + \frac{1}{2} (4 + 1) \cdot -1 \right] \\
&= \frac{11}{2}
\end{aligned}$$

- (e) Using (a), the integral $\int P dx + Q dy$ along the path $C \rightarrow ABFEAEFBA$ may

be approximated as

$$\begin{aligned}
\int_C P dx + Q dy &\sim \left[\frac{1}{2} (0+3) \cdot 1 + \frac{1}{2} (1+2) \cdot 0 \right] + \left[\frac{1}{2} (3+0) \cdot 0 + \frac{1}{2} (2+4) \cdot 1 \right] \\
&\quad + \left[\frac{1}{2} (0+3) \cdot -1 + \frac{1}{2} (4+2) \cdot 0 \right] + \left[\frac{1}{2} (3+0) \cdot 0 + \frac{1}{2} (2+1) \cdot -1 \right] \\
&\quad + \left[\frac{1}{2} (0+3) \cdot 0 + \frac{1}{2} (1+2) \cdot 1 \right] + \left[\frac{1}{2} (3+0) \cdot 1 + \frac{1}{2} (2+4) \cdot 0 \right] \\
&\quad + \left[\frac{1}{2} (0+3) \cdot 0 + \frac{1}{2} (4+2) \cdot -1 \right] + \left[\frac{1}{2} (3+0) \cdot -1 + \frac{1}{2} (2+1) \cdot 0 \right] \\
&= 0
\end{aligned}$$

7. Let C be a smooth curve in the xy -plane and let $f(x, y) > 0$ be a continuous function defined over a region of the xy -plane containing the curve C . The equation $z = f(x, y)$ then is the equation of a surface that lies above the region of the xy -plane containing the curve C . Next, we imagine moving a straight line along C perpendicular to the xy -plane, effectively tracing out a "wall" standing on C , orthogonal to the xy -plane. This "wall" cuts the surface $z = f(x, y)$, forming a curve on it that lies above the curve C . In fact, the curve C may be interpreted as the projection of the surface curve onto the xy -plane. Using (5.11), the line integral

$$\int_C f(x, y) ds = \lim_{\substack{n \rightarrow \infty \\ \max \Delta_i s \rightarrow 0}} \sum_{i=1}^n f(x_i^*, y_i^*) \Delta_i s$$

then may be interpreted as an infinite sum of the length of each straight line directed from C to the surface curve lying above it in the limit where the distance Δs between each subsequent line becomes infinitely small, effectively tracing out a "wall" with height at each point (x, y) given by $f(x, y)$. This may be interpreted the as the area of the cylindrical surface $0 \leq z \leq f(x, y)$, (x, y) on C .

Section 5.5

1. Let the vector $v = (x^2 + y^2)\mathbf{i} + 2xy\mathbf{j}$ be given. Then by (5.25) and (5.29)

- (a) The integral $\int_C v_T ds$ along the path $C \rightarrow y = x$ from $(0, 0)$ to $(1, 1)$ may be evaluated as

$$\int_C v_T ds = \int_C (x^2 + y^2) dx + 2xy dy \stackrel{(5.6)(5.9)}{=} \int_0^1 2x^2 dx + \int_0^1 2y^2 dy = \frac{4}{3}$$

- (b) The integral $\int_C v_T ds$ along the path $C \rightarrow y = x^2$ from $(0, 0)$ to $(1, 1)$ may be evaluated as

$$\int_C v_T ds = \int_C (x^2 + y^2) dx + 2xy dy \stackrel{(5.6)(5.7)}{=} \int_0^1 (x^2 + 5x^4) dx = \frac{4}{3}$$

- (c) The integral $\int_C v_T ds$ along the broken line from $(0,0)$ to $(1,1)$ with corner at $(1,0)$ may be evaluated as

$$\begin{aligned}\int_C v_T ds &= \int_C (x^2 + y^2) dx + 2xy dy \\ &= \int_{(0,0)}^{(1,0)} (x^2 + y^2) dx + 2xy dy + \int_{(1,0)}^{(1,1)} (x^2 + y^2) dx + 2xy dy \\ &= \int_0^1 x^2 dx + \int_0^1 2y dy = \frac{4}{3}\end{aligned}$$

2. Let $\mathbf{v} = P(x, y)\mathbf{i} + Q(x, y)\mathbf{j}$ be the same vector as given in Problem 1, and let \mathbf{n} be the unit normal vector 90° behind the tangent vector \mathbf{T} as given by (5.37). Then the normal component of \mathbf{v} is given by $v_n = \mathbf{v} \cdot \mathbf{n} = (P\mathbf{i} + Q\mathbf{j}) \cdot (y_s\mathbf{i} - x_s\mathbf{j}) = -Qx_s + Py_s$. Then by (5.25) and (5.29)

- (a) The integral $\int_C v_n ds$ along the path $C \rightarrow y = x$ from $(0,0)$ to $(1,1)$ may be evaluated as

$$\int_C v_n ds = \int_C -2xy dx + (x^2 + y^2) dy \stackrel{(5.6)(5.9)}{=} \int_0^1 -2x^2 dx + \int_0^1 2y^2 dy = 0$$

- (b) The integral $\int_C v_n ds$ along the path $C \rightarrow y = x^2$ from $(0,0)$ to $(1,1)$ may be evaluated as

$$\int_C v_n ds = \int_C -2xy dx + (x^2 + y^2) dy \stackrel{(5.6)(5.7)}{=} \int_0^1 2x^5 dx = \frac{1}{3}$$

- (c) The integral $\int_C v_n ds$ along the broken line from $(0,0)$ to $(1,1)$ with corner at $(1,0)$ may be evaluated as

$$\begin{aligned}\int_C v_n ds &= \int_C -2xy dx + (x^2 + y^2) dy \\ &= \int_{(0,0)}^{(1,0)} -2xy dx + (x^2 + y^2) dy + \int_{(1,0)}^{(1,1)} -2xy dx + (x^2 + y^2) dy \\ &= \int_0^1 (1 + y^2) dy = \frac{4}{3}\end{aligned}$$

3. Let the gravitational force near a point on the earth's surface be represented approximately by the vector $\mathbf{F} = P(x, y)\mathbf{i} + Q(x, y)\mathbf{j} = -mg\mathbf{j}$, where the y -axis points upwards. Then by (5.29) and the fact that $P(x, y) = 0$ the work done by the force \mathbf{F} on a body moving in a vertical plane from height h_1 to height h_2 along any path is equal to

$$\int_C F_T ds = \int_C (P \cos \alpha + Q \sin \alpha) ds = \int_C Q dy = - \int_{h_1}^{h_2} mg dy = -mgy \Big|_{h_1}^{h_2} = mg(h_1 - h_2)$$

4. Let the gravitational force \mathbf{F} be given by $\mathbf{F} = -(kMm/r^2)(\mathbf{r}/r)$. Then in order to compute the work by the gravitational force \mathbf{F} in bringing a particle to its present position r from infinite distance along the ray through the earth's center, we will represent the curve C in terms of parameter t and then use (5.34) to get

$$\begin{aligned}\int_C \mathbf{F} \cdot d\mathbf{r} &= \int_{-\infty}^r \left(\mathbf{F} \cdot \frac{d\mathbf{r}}{dt} \right) dt = \int_{-\infty}^r \left(-\frac{kMm}{t^2} \frac{d\mathbf{r}}{dt} \cdot \frac{d\mathbf{r}}{dt} \right) dt = \int_{-\infty}^r -\frac{kMm}{t^2} dt = \frac{kMm}{t} \Big|_{-\infty}^r \\ &= kMm \left(\frac{1}{r} - \frac{1}{\infty} \right) = \frac{kMm}{r} \\ &= -U\end{aligned}$$

where $(d\mathbf{r}/dr) \cdot (d\mathbf{r}/dr) = 1$ follows from the fact that $d\mathbf{r}/dr$ is a unit vector.

5. (a) By (5.40) the integral $\oint_C ay \, dx + bx \, dy$ may be written as

$$\oint_C ay \, dx + bx \, dy = \iint_R (b - a) \, dx \, dy = (b - a) A$$

where A is the area enclosed by the curve C .

- (b) By (5.40) the integral $\oint e^x \sin y \, dx + e^x \cos y \, dy$ around the rectangle with vertices $(0, 0), (1, 0), (1, \pi/2), (0, \pi/2)$ may be written as

$$\oint e^x \sin y \, dx + e^x \cos y \, dy = \int_0^{\pi/2} \int_0^1 (e^x \cos y - e^x \cos y) \, dx \, dy = 0$$

- (c) By (5.40) and (4.61) the integral $\oint (2x^3 - y^3) \, dx + (x^3 + y^3) \, dy$ around the circle $x^2 + y^2 = 1$ may be written as

$$\oint (2x^3 - y^3) \, dx + (x^3 + y^3) \, dy = 3 \int_0^1 \int_0^{2\pi} r^3 \, d\theta \, dr = 6\pi \int_0^1 r^3 \, dr = \frac{3\pi}{2}$$

- (d) By (5.43) and (3.31) the integral $\oint_C u_T \, ds$, where $\mathbf{u} = \text{grad}(x^2y)$ and C is the circle $x^2 + y^2 = 1$ may be written as

$$\oint_C u_T \, ds = \iint_R \text{curl}_z \mathbf{u} \, dx \, dy = \iint_R \text{curl}_z \text{grad}(x^2y) \, dx \, dy = 0$$

- (e) By (5.44) the integral $\oint_C v_n \, ds$, where $\mathbf{v} = (x^2 + y^2)\mathbf{i} - 2xy\mathbf{j}$ and C is the circle $x^2 + y^2 = 1$ (\mathbf{n} being the outer normal) may be written as

$$\begin{aligned}\oint_C v_n \, ds &= \iint_R \text{div} \mathbf{v} \, dx \, dy = \iint_R \text{div} [(x^2 + y^2)\mathbf{i} - 2xy\mathbf{j}] \, dx \, dy = \iint_R (2x - 2x) \, dx \, dy \\ &= 0\end{aligned}$$

- (f) Let $F = (x - 2)^2 + y^2$. Then by (2.117) $\partial F / \partial n = \nabla F \cdot \mathbf{n} = \mathbf{v} \cdot \mathbf{n}$ and since $\oint_C \mathbf{v} \cdot \mathbf{n} ds = \oint_C v_n ds$ we find by (5.44) and (4.64)

$$\begin{aligned} \oint_C v_n ds &= \iint_R \operatorname{div} (\nabla F) dx dy = \iint_R \nabla \cdot \nabla F dx dy = \iint_R \nabla^2 F dx dy \\ &= \iint_R \nabla^2 [(x - 2)^2 + y^2] dx dy \\ &= 4 \int_0^{2\pi} \int_0^1 r dr d\theta = 4\pi \end{aligned}$$

- (g) Let $F = \ln[(x - 2)^2 + y^2]^{-1}$. Then by (2.117) $\partial F / \partial n = \nabla F \cdot \mathbf{n} = \mathbf{v} \cdot \mathbf{n}$ and since $\oint_C \mathbf{v} \cdot \mathbf{n} ds = \oint_C v_n ds$ we find by (5.44)

$$\begin{aligned} \oint_C v_n ds &= \iint_R \operatorname{div} (\nabla F) dx dy = \iint_R \nabla \cdot \nabla F dx dy = \iint_R \nabla^2 F dx dy \\ &= \iint_R \nabla \ln \frac{1}{(x - 2)^2 + y^2} dx dy \\ &= 2 \iint_R \frac{x^2 - 4x + 4 - y^2 - (x - 2)^2 + y^2}{[(x - 2)^2 + y^2]^2} dx dy = 0 \end{aligned}$$

- (h) By (5.40) the integral $\oint_C f(x) dx + g(y) dy$ may be written as

$$\oint_C f(x) dx + g(y) dy = \iint_R \left[\frac{\partial}{\partial x} g(y) - \frac{\partial}{\partial y} f(x) \right] dx dy = 0$$

6. Let $\mathbf{r} = x\mathbf{i} + y\mathbf{j}$ be the position vector of an arbitrary point (x, y) and let \mathbf{n} be the outer normal to some arbitrary closed curve C . Then by (5.44)

$$\begin{aligned} \frac{1}{2} \oint_C r_n ds &= \frac{1}{2} \oint_C \mathbf{r} \cdot \mathbf{n} ds = \frac{1}{2} \iint_R \operatorname{div} \mathbf{r} dx dy = \frac{1}{2} \iint_R \nabla \cdot (x\mathbf{i} + y\mathbf{j}) dx dy \\ &= \frac{1}{2} \iint_R \left(\frac{\partial}{\partial x} x + \frac{\partial}{\partial y} y \right) dx dy \\ &= \iint_R dx dy = A \end{aligned}$$

7. As for Problem 2(a), let the line integral $\int_{(0,-1)}^{(0,1)} y^2 dx + x^2 dy$, where C is the semi-circle

$x = \sqrt{1-y^2}$ be given. Then by (5.40) and (4.64)

$$\begin{aligned}
\oint_C y^2 dx + x^2 dy &= \iint_R \left(\frac{\partial}{\partial x} x^2 - \frac{\partial}{\partial y} y^2 \right) dx dy = 2 \iint_R (x - y) dx dy \\
&= 2 \int_{-1}^1 \int_0^{\sqrt{1-y^2}} (x - y) dx dy \\
&= 2 \int_{-\pi/2}^{\pi/2} \int_0^1 (\cos \theta - \sin \theta) r^2 dr d\theta \\
&= \frac{2}{3} \int_{\pi/2}^{\pi/2} (\cos \theta - \sin \theta) d\theta = \frac{4}{3}
\end{aligned}$$

As for Problem 3(a), let the line integral $\oint_C y^2 dx + xy dy$, where C is the square with vertices $(1, 1)$, $(-1, 1)$, $(-1, -1)$, $(1, -1)$ be given. Then by (5.40)

$$\begin{aligned}
\oint_C y^2 dx + xy dy &= \iint_R \left[\frac{\partial}{\partial x} (xy) - \frac{\partial}{\partial y} y^2 \right] dx dy = - \iint_R y dx dy = - \int_{-1}^1 \int_{-1}^1 y dx dy \\
&= - \int_{-1}^1 xy \Big|_{-1}^1 dy = -2 \int_{-1}^1 y dy \\
&= -y^2 \Big|_{-1}^1 = 0
\end{aligned}$$

As for Problem 3(b), let the line integral $\oint_C y dx - x dy$, where C is the circle $x^2 + y^2 = 1$ be given. Then by (5.40) and (4.64)

$$\begin{aligned}
\oint_C y dx - x dy &= - \iint_R \left(\frac{\partial}{\partial x} x + \frac{\partial}{\partial y} y \right) dx dy = -2 \iint_R dx dy = -2 \int_0^{2\pi} \int_0^1 r dr d\theta \\
&= - \int_0^{2\pi} d\theta = -2\pi
\end{aligned}$$

As for Problem 3(c), let the line integral $\oint_C x^2 y^2 dx - xy^3 dy$, where C is the triangle with vertices $(0, 0)$, $(1, 0)$, $(1, 1)$ be given. Then by (5.40)

$$\begin{aligned}
\oint_C x^2 y^2 dx - xy^3 dy &= - \iint_R \left[\frac{\partial}{\partial x} (xy^3) + \frac{\partial}{\partial y} (x^2 y^2) \right] dx dy \\
&= - \iint_R (y^3 + 2x^2 y) dx dy = - \int_0^1 \int_0^x (y^3 + 2x^2 y) dy dx \\
&= - \int_0^1 \left[\frac{y^4}{4} + x^2 y^2 \right]_0^x dx = -\frac{5}{4} \int_0^1 x^4 dx = -\frac{1}{4}
\end{aligned}$$

As for Problem 4(a), let the line integral $\oint_C (x^2 - y^2) ds$, where C is the circle $x^2 + y^2 = 4$ be given. Then by (5.44) and the fact that \mathbf{n} may be written as $\mathbf{n} = (x\mathbf{i} + y\mathbf{j})/|x + y|$

$$\oint_C (x^2 - y^2) ds = a \iint_R \operatorname{div} (x\mathbf{i} - y\mathbf{j}) dx dy = a \iint_R \left(\frac{\partial}{\partial x} \mathbf{i} + \frac{\partial}{\partial y} \mathbf{j} \right) \cdot (x\mathbf{i} - y\mathbf{j}) dx dy = 0$$

Section 5.7

1.