

QM Lecture 13



HARMONIC OSC.

$$\Delta x \Delta p_x = (n + \frac{1}{2}) \hbar$$

SO GROUND ($n=0$) HAS $\Delta x \Delta p_x = \frac{\hbar}{2}$, MIN. ALLOWED!

(Q: why is this not surprising?)

A: $\psi_0(x) \sim e^{-x^2/\hbar}$ GAUSSIAN \Rightarrow MIN. $\Delta x \Delta p_x$

FULL SOLUTION OF ONE-DIM. SHO

(Q: why do this? A: more gen soln ted)

GOAL SOLVE $\langle x | \hat{H} | E \rangle = E \langle x | E \rangle$ (what does "solve" mean?)

$$\langle x | \left(\frac{\hat{p}_x^2}{2m} + \frac{1}{2} m \omega^2 \hat{x}^2 \right) | E \rangle = E \langle x | E \rangle$$

$$-\frac{\hbar^2}{2m} \frac{d^2}{dx^2} \langle x | E \rangle + \frac{1}{2} m \omega^2 x^2 \langle x | E \rangle = E \langle x | E \rangle \quad \text{S.EQN.}$$

SIMP. $y := \sqrt{\frac{m\omega}{\hbar}} x \quad \epsilon := \frac{2E}{\hbar\omega}$

$$\langle x | E \rangle = \psi(x) = \psi(y) \quad \frac{d}{dx} \frac{d}{dx} = \left(\frac{dy}{dx} \right)^2 \frac{d}{dy} \frac{d}{dy} = \frac{m\omega}{\hbar}$$

$$-\frac{\hbar^2}{2m} \frac{m\omega}{\hbar} \frac{d^2}{dy^2} \psi(y) + \frac{1}{2} m \omega^2 \frac{\hbar}{m\omega} y^2 \psi(y) = \frac{\hbar\omega\epsilon}{2} \psi(y)$$

$$\Rightarrow \left(\frac{d^2}{dy^2} + \epsilon - y^2 \right) \psi(y) = 0 \quad (*)$$

as $y \rightarrow \infty$ HAVE $\frac{d^2 \psi}{dy^2} - y^2 \psi = 0$

$$\Rightarrow \psi(y) \Big|_{y \rightarrow \infty} = A e^{-y^2/2} \quad (\text{note diff from book})$$

∴ ANSATZ ("educated guess")

$$\psi(y) = A(y) e^{-y^2/2} \quad (**)$$

(**) INTO (*)

$$\frac{d\psi}{dy} = \frac{dA}{dy} e^{-y^2/2} - y A e^{-y^2/2}$$

$$\frac{d^2\psi}{dy^2} = \frac{d^2A}{dy^2} e^{-y^2/2} - y \frac{dA}{dy} e^{-y^2/2} - A e^{-y^2/2} - y \frac{dA}{dy} e^{-y^2/2} + \underbrace{y^2 A e^{-y^2/2}}_{\text{CANC. by } -y^2\psi}$$

$$\therefore \boxed{\frac{d^2A(y)}{dy^2} - 2y \frac{dA(y)}{dy} + (\epsilon - 1) A(y) = 0}$$

↳ HERMITE'S DIFF EQ.

TRICK $A(y) = \sum_{k=0}^{\infty} a_k y^k$

$$\Rightarrow \sum_{k=0}^{\infty} k(k-1) a_k y^{k-2} - 2 \sum_{k=0}^{\infty} k a_k y^k + (\epsilon - 1) \sum_{k=0}^{\infty} a_k y^k = 0$$

$k=0,1$ TERMS VANISH! $\sum_{k=0}^{\infty} k(k-1) a_k y^{k-2} = \sum_{k'=0}^{\infty} (k'+2)(k'+1) a_{k'+2} y^{k'}$

$$= \sum_{k=0}^{\infty} (k+2)(k+1) a_{k+2} y^k$$

$$\Rightarrow \sum_{k=0}^{\infty} [(k+2)(k+1) a_{k+2} - 2k a_k + (\epsilon - 1) a_k] y^k = 0$$

$$\Rightarrow \frac{a_{k+2}}{a_k} = \frac{2k+1-\epsilon}{(k+1)(k+2)}$$

- RECURSION REL.

- SPEC. a_0, a_1 GET ALL OTHERS

ASYMPTOTICS

$$\frac{a_{k+2}}{a_k} \xrightarrow{k \rightarrow \infty} \frac{2k}{k^2} = \frac{2}{k}$$

COMPARE $e^{y^2} = \sum_{n=0}^{\infty} \frac{y^{2n}}{n!} = \sum_{k=0}^{\infty} b_k y^k$

$$\Rightarrow \frac{b_{k+2}}{b_k} = \frac{\left(\frac{k}{2}\right)!}{\left[\left(\frac{k}{2}+1\right)! \right]} = \frac{1}{\frac{k}{2}+1} \xrightarrow{k \rightarrow \infty} \frac{2}{k}$$

so $A(y) \sim e^{-y^2/2}$ @ LARGE k NOT GOOD!

CUTOFF: IF \exists SOME $k=n$ w/ $a_{n+2}=0$ THEN $a_{n+2+i}=0 \forall i>0$

SOLVE $2n+1-\varepsilon=0 \Rightarrow \varepsilon=2n+1$

$$\Rightarrow \boxed{\frac{a_{k+2}}{a_k} = \frac{2(k-n)}{(k+2)(k+1)}}$$

* SOL. DEP NOW ON a_0, a_1, n .

* BUT n EVEN REQ $a_1=0 \Rightarrow \psi(y)$ EVEN

n ODD REQ $a_0=0 \Rightarrow \psi(y)$ ODD

EXAMPLES

$n=0$ | a_0 CONST $a_1=0$ $a_2=0 \dots$

$n=1$ | $a_0=0$ $a_1=\text{CONST}$ $a_2=0 \dots a_3=0 \dots$

$n=2$ | $a_0=\text{CONST}$ $a_1=0$ $a_2=-2a_0$ $a_3=a_4=\dots=0$

$n=3$ | $a_1=\text{CONST}$ $a_0=0$ $a_2=0$ $a_3=-\frac{2}{3}a_1$ $a_4=a_5=\dots=0$

NORMALIZATION FIXES a_0, a_1

$$\therefore \Psi_n(x) = \langle x | E_n \rangle = \left(\frac{m\omega}{\pi \hbar} \right)^{\frac{1}{4}} \frac{1}{\sqrt{2^n n!}} A_n(y) e^{-y^2/2}$$
$$y = \sqrt{\frac{m\omega}{\hbar}} x$$

FIRST FEW A'_n

$$A_0(y) = 1$$

$$A_1(y) = 2y$$

$$A_2(y) = 4y^2 - 2$$

$$A_3(y) = 8y^3 - 12y$$

$$A_4(y) = 16y^4 - 48y^2 + 12$$

$$A_5 = 32y^5 - 160y^3 + 120y$$

LESSONS

- 1) a_n TERM. \Rightarrow ONLY EVEN OR ODD POWERS \Rightarrow IF EVEN OR ODD
- 2) THAT REQ PICKING $n \in \mathbb{Z}$ $n \geq 0$
- 3) THEN $E = \frac{2E}{\hbar\omega} = 2n+1 \Rightarrow E = (n + \frac{1}{2})\hbar\omega$ ✓ QUANTIZED

NOTE 2) 3) ALREADY KNEW FROM QTA TREATMENT.

(was much easier that way. There are many ways to explain a fact!)

