

LABORATORY REPORT

Report Status: Final

NATIONAL LABS 3948 TRUST WAY HAYWARD, CA 94545 Ph.: (800) 800-6060

 Patient Name:
 SMITH, BRENDA
 Accession:
 1708210231

 Patient ID:
 07131957
 Collect Date:
 08/10/2017

 DOB, Age, Sex:
 7/13/1957 60 Female
 Receive Date:
 08/21/2017

 Client:
 (1028) Lags Spine and Sportscare
 Release Date:
 08/21/2017

Requested By: Hector Sanchez (1104251933)

Medications: Lidocaine, Norco, Nortriptyline, Flexeril, Aspirin, Levothyroxine, Metformin

Test Not Performed (TNP). Quantity Not Sufficient (QNS) for Custom Pain Panel testing.

*** Comments: Specimens received more than 3 days post collection may be compromised and not reflect actual concentration. This may limit detection of low levels. This test is not approved by the FDA for the treatment, diagnosis, or prevention, of any medical condition. It is performed for information and is intended for educational and research purposes only. Only a physician familiar with a Patient() 's individual medical history can make medical judgments and give that patient specific medical advice. results are not intended to be a substitute for consultation with a licensed health care provider.

Those analytes' test results, reported as detected by LC/MS, have not been approved by the FDA. The performance characteristics of the assays have been determined by National Labs in accordance with Clinical Laboratory Improvement Act of 1988 (CLIA '88) guidelines.

False positive and false negative test results do occur. A licensed health care provider must provide any interpretation or recommendation for treatment based upon test results. The results are not for use as evidence in civil or criminal cases. Specimens are not obtained under the guidelines required for secure collection and, therefore, cannot be used for employment purposes. The results are based upon the method used for testing and the sensitivity of that method. A negative result indicates that the drugs tested for were not detected at the sensitivity level used. Positive results may be possible with the same or subsequent specimens if testing is performed using a greater degree of sensitivity. Other factors that may affect the results include the type of drug; the amount, duration and time of last drug use; the body type and weight of the individual being testing; and, in the case of urine drug screens, the concentration, acidity, or alkalinity of the urine.