



Norwegian University of  
Science and Technology

# **TMA4300 - COMPUTER INTENSIVE STATISTICAL METHODS**

Exercise 1: Problem C


Jim Totland, Martin Tufte

March 1, 2022

# Problem C: Monte Carlo integration and variance reduction

Aim: Learn about variance reduction techniques in Monte Carlo integration.

Problem: Find  $\theta = \text{Prob}(X > 4)$  when  $X \sim N(0, 1)$ .



## C.1: Monte Carlo integration

Estimate  $\theta$  by Monte Carlo integration using  $n = 100000$  samples from the standard normal distribution. Compute also a 95% confidence interval for  $\theta$  based on these  $n$  samples.

## C.1: Monte Carlo integration

Estimate  $\theta$  by Monte Carlo integration using  $n = 100000$  samples from the standard normal distribution. Compute also a 95% confidence interval for  $\theta$  based on these  $n$  samples.

Rewrite  $\theta$  using expected value:

$$\theta = \Pr\{X > 4\} = \int_4^{\infty} \phi(x) \, dx = \int_{-\infty}^{\infty} \mathbb{I}(x > 4) \phi(x) \, dx = \mathbb{E}[\mathbb{I}(X > 4)]$$

## C.1: Monte Carlo integration

Estimate  $\theta$  by Monte Carlo integration using  $n = 100000$  samples from the standard normal distribution. Compute also a 95% confidence interval for  $\theta$  based on these  $n$  samples.

Rewrite  $\theta$  using expected value:

$$\theta = \Pr\{X > 4\} = \int_4^{\infty} \phi(x) \, dx = \int_{-\infty}^{\infty} \mathbb{I}(x > 4) \phi(x) \, dx = E[\mathbb{I}(X > 4)]$$

Use sample mean as estimator:

$$\hat{\theta} = \frac{1}{n} \sum_{i=1}^n \underbrace{\mathbb{I}(x_i > 4)}_{h(x_i)}$$

## C.1: Confidence Intervals

Estimate variance:

$$\text{Var}[\hat{\theta}] = \frac{1}{n^2} \sum_{i=1}^n \text{Var}_f[h(X_i)] = \frac{1}{n} \text{Var}_f[h(X)] \approx \frac{1}{n} \left( \frac{1}{n-1} \sum_{i=1}^n (h(x_i) - \hat{\theta})^2 \right) =: \hat{\sigma}^2$$

## C.1: Confidence Intervals

Estimate variance:

$$\text{Var}[\hat{\theta}] = \frac{1}{n^2} \sum_{i=1}^n \text{Var}_f[h(X_i)] = \frac{1}{n} \text{Var}_f[h(X)] \approx \frac{1}{n} \left( \frac{1}{n-1} \sum_{i=1}^n (h(x_i) - \hat{\theta})^2 \right) =: \hat{\sigma}^2$$

Using the central limit theorem:  $\hat{\theta} \sim \mathcal{N}(\theta, \text{Var}[\hat{\theta}]) \implies \frac{\hat{\theta} - \theta}{\hat{\sigma}} \sim t_{n-1}$ .

A 95% confidence interval is therefore

$$[\hat{\theta} - t_{n-1, 0.025} \cdot \hat{\sigma}, \hat{\theta} + t_{n-1, 0.025} \cdot \hat{\sigma}].$$

## C.2: Importance sampling

$$\hat{\theta}_{IS} = \frac{1}{n} \sum_{i=1}^n \frac{h(X_i)f(X_i)}{g(X_i)} = \frac{1}{n} \sum_{i=1}^n h(X_i)w(X_i).$$




## C.2: Importance sampling

$$\hat{\theta}_{IS} = \frac{1}{n} \sum_{i=1}^n \frac{h(X_i)f(X_i)}{g(X_i)} = \frac{1}{n} \sum_{i=1}^n h(X_i)w(X_i).$$

Need to sample from proposal distribution  $g(x)$ .

## C.2: Importance sampling

$$g(x) = \begin{cases} cx \exp -\frac{1}{2}x^2 & , x > 4 \\ 0 & , \text{else} \end{cases}$$



## C.2: Importance sampling

$\hat{\theta}_{IS} = \frac{1}{n} \sum_{i=1}^n \frac{h(X_i)f(X_i)}{g(X_i)} = \frac{1}{n} \sum_{i=1}^n h(X_i)w(X_i)$ . Find variance:

$$\begin{aligned}\text{Var}[\hat{\theta}_{IS}] &= \frac{1}{n^2} \sum_{i=1}^n \text{Var}_g[h(X_i)w(X_i)] = \frac{1}{n} \text{Var}_g[h(X)w(X)] \\ &\approx \frac{1}{n} \left( \frac{1}{n-1} \sum_{i=1}^n (h(x_i)w(x_i) - \hat{\theta}_{IS})^2 \right) =: \hat{\sigma}_{IS}^2.\end{aligned}$$

## C.2: Importance sampling

$\hat{\theta}_{IS} = \frac{1}{n} \sum_{i=1}^n \frac{h(X_i)f(X_i)}{g(X_i)} = \frac{1}{n} \sum_{i=1}^n h(X_i)w(X_i)$ . Find variance:

$$\begin{aligned}\text{Var}[\hat{\theta}_{IS}] &= \frac{1}{n^2} \sum_{i=1}^n \text{Var}_g[h(X_i)w(X_i)] = \frac{1}{n} \text{Var}_g[h(X)w(X)] \\ &\approx \frac{1}{n} \left( \frac{1}{n-1} \sum_{i=1}^n (h(x_i)w(x_i) - \hat{\theta}_{IS})^2 \right) =: \hat{\sigma}_{IS}^2.\end{aligned}$$

95% Confidence interval:  $\left[ \hat{\theta}_{IS} - t_{n-1,0.025} \cdot \hat{\sigma}_{IS}, \hat{\theta}_{IS} + t_{n-1,0.025} \cdot \hat{\sigma}_{IS} \right]$

## C.3: Antithetic sampling

$$\hat{\theta}_{AS} = \frac{1}{2} \left( \hat{\theta}_{IS}^{(1)} + \hat{\theta}_{IS}^{(2)} \right)$$

## C.3: Antithetic sampling

$$\hat{\theta}_{AS} = \frac{1}{2} \left( \hat{\theta}_{IS}^{(1)} + \hat{\theta}_{IS}^{(2)} \right)$$

$$\hat{\theta}_{IS}^{(1)} = \frac{1}{n} \sum_{i=1}^n h \left( X_i^{(1)} \right) w \left( X_i^{(1)} \right) \text{ and } \hat{\theta}_{IS}^{(2)} = \frac{1}{n} \sum_{i=1}^n h \left( X_i^{(2)} \right) w \left( X_i^{(2)} \right).$$

The  $X_j^{(1)}$ 's are based on  $u$  while the  $X_j^{(2)}$ 's are based on  $u - 1$  in the inversion sampling of  $g(x)$ .

## C.3: Antithetic sampling

$$\hat{\theta}_{AS} = \frac{1}{2} \left( \hat{\theta}_{IS}^{(1)} + \hat{\theta}_{IS}^{(2)} \right)$$

$$\hat{\theta}_{IS}^{(1)} = \frac{1}{n} \sum_{i=1}^n h \left( X_i^{(1)} \right) w \left( X_i^{(1)} \right) \quad \text{and} \quad \hat{\theta}_{IS}^{(2)} = \frac{1}{n} \sum_{i=1}^n h \left( X_i^{(2)} \right) w \left( X_i^{(2)} \right).$$

The  $X_j^{(1)}$ 's are based on  $u$  while the  $X_j^{(2)}$ 's are based on  $u - 1$  in the inversion sampling of  $g(x)$ .

$\implies$  Only need  $n = 50000$  to make  $2n$  (antithetic) samples from  $g(x)$ .

## C.3: Confidence intervals part 1

$$\hat{\theta}_{AS} = \frac{1}{2} \left( \hat{\theta}_{IS}^{(1)} + \hat{\theta}_{IS}^{(2)} \right).$$



## C.3: Confidence intervals part 1

$$\hat{\theta}_{AS} = \frac{1}{2} \left( \hat{\theta}_{IS}^{(1)} + \hat{\theta}_{IS}^{(2)} \right). \text{ Variance:}$$

$$\text{Var} \left[ \hat{\theta}_{AS} \right] = \frac{1}{4} \left( \text{Var} \left[ \hat{\theta}_{IS}^{(1)} \right] + \text{Var} \left[ \hat{\theta}_{IS}^{(2)} \right] + 2\text{Cov} \left[ \hat{\theta}_{IS}^{(1)}, \hat{\theta}_{IS}^{(2)} \right] \right).$$

## C.3: Confidence intervals part 1

$$\hat{\theta}_{AS} = \frac{1}{2} \left( \hat{\theta}_{IS}^{(1)} + \hat{\theta}_{IS}^{(2)} \right). \text{ Variance:}$$

$$\text{Var} \left[ \hat{\theta}_{AS} \right] = \frac{1}{4} \left( \text{Var} \left[ \hat{\theta}_{IS}^{(1)} \right] + \text{Var} \left[ \hat{\theta}_{IS}^{(2)} \right] + 2\text{Cov} \left[ \hat{\theta}_{IS}^{(1)}, \hat{\theta}_{IS}^{(2)} \right] \right).$$

How to make confidence interval? Rewrite  $\hat{\theta}_{AS}$

## C.3: Confidence intervals part 1

$$\hat{\theta}_{AS} = \frac{1}{2} \left( \hat{\theta}_{IS}^{(1)} + \hat{\theta}_{IS}^{(2)} \right). \text{ Variance:}$$

$$\text{Var} \left[ \hat{\theta}_{AS} \right] = \frac{1}{4} \left( \text{Var} \left[ \hat{\theta}_{IS}^{(1)} \right] + \text{Var} \left[ \hat{\theta}_{IS}^{(2)} \right] + 2\text{Cov} \left[ \hat{\theta}_{IS}^{(1)}, \hat{\theta}_{IS}^{(2)} \right] \right).$$

How to make confidence interval? Rewrite  $\hat{\theta}_{AS}$

$$\hat{\theta}_{AS} = \frac{1}{n} \sum_{i=1}^n \underbrace{\frac{1}{2} \left( h \left( X_i^{(1)} \right) w \left( X_i^{(1)} \right) + h \left( X_i^{(2)} \right) w \left( X_i^{(2)} \right) \right)}_{=Y_i} = \frac{1}{n} \sum_{i=1}^n Y_i.$$

## C.3: Confidence intervals part 1

$$\hat{\theta}_{AS} = \frac{1}{2} \left( \hat{\theta}_{IS}^{(1)} + \hat{\theta}_{IS}^{(2)} \right). \text{ Variance:}$$

$$\text{Var} \left[ \hat{\theta}_{AS} \right] = \frac{1}{4} \left( \text{Var} \left[ \hat{\theta}_{IS}^{(1)} \right] + \text{Var} \left[ \hat{\theta}_{IS}^{(2)} \right] + 2\text{Cov} \left[ \hat{\theta}_{IS}^{(1)}, \hat{\theta}_{IS}^{(2)} \right] \right).$$

How to make confidence interval? Rewrite  $\hat{\theta}_{AS}$

$$\hat{\theta}_{AS} = \frac{1}{n} \sum_{i=1}^n \underbrace{\frac{1}{2} \left( h \left( X_i^{(1)} \right) w \left( X_i^{(1)} \right) + h \left( X_i^{(2)} \right) w \left( X_i^{(2)} \right) \right)}_{=Y_i} = \frac{1}{n} \sum_{i=1}^n Y_i.$$

$Y_i$ 's independent  $\implies$  can use  $t$ -distribution to create C.I.

## C.3: Confidence intervals part 2

Use sample variance:

$$\hat{\sigma}_{AS}^2 = \frac{1}{n} \left( \frac{1}{n-1} \sum_{i=1}^n (Y_i - \hat{\theta}_{AS})^2 \right)$$

## C.3: Confidence intervals part 2

Use sample variance:

$$\hat{\sigma}_{AS}^2 = \frac{1}{n} \left( \frac{1}{n-1} \sum_{i=1}^n (Y_i - \hat{\theta}_{AS})^2 \right)$$

And we know:

$$\frac{\hat{\theta}_{AS} - \theta}{\hat{\sigma}_{AS}} \sim t_{n-1}$$

## C.3: Confidence intervals part 2

Use sample variance:

$$\hat{\sigma}_{AS}^2 = \frac{1}{n} \left( \frac{1}{n-1} \sum_{i=1}^n (Y_i - \hat{\theta}_{AS})^2 \right)$$

And we know:

$$\frac{\hat{\theta}_{AS} - \theta}{\hat{\sigma}_{AS}} \sim t_{n-1}$$

From which we construct a confidence interval.