

Compulsory exercise 1: Group 39

TMA4268 Statistical Learning V2021

Alexander J Ohrt, Jim Totland

09 februar, 2021

Problem 1

a)

We assume that \mathbf{Y} is a multivariate normal, which gives the distribution $\mathbf{Y} \sim N_n(\mathbf{X}\beta, \sigma^2\mathbf{I})$.

$$\begin{aligned} E(\tilde{\beta}) &= E((\mathbf{X}^T\mathbf{X} + \lambda\mathbf{I})^{-1}\mathbf{X}^T\mathbf{Y}) = (\mathbf{X}^T\mathbf{X} + \lambda\mathbf{I})^{-1}\mathbf{X}^T E(\mathbf{Y}) \\ &= (\mathbf{X}^T\mathbf{X} + \lambda\mathbf{I})^{-1}\mathbf{X}^T E(\mathbf{X}\beta + \varepsilon) = (\mathbf{X}^T\mathbf{X} + \lambda\mathbf{I})^{-1}\mathbf{X}^T\mathbf{X}\beta \end{aligned}$$

and

$$\begin{aligned} \text{Cov}(\tilde{\beta}) &= \text{Cov}((\mathbf{X}^T\mathbf{X} + \lambda\mathbf{I})^{-1}\mathbf{X}^T\mathbf{Y}) = ((\mathbf{X}^T\mathbf{X} + \lambda\mathbf{I})^{-1}\mathbf{X}^T) \text{Cov}(\mathbf{Y}) ((\mathbf{X}^T\mathbf{X} + \lambda\mathbf{I})^{-1}\mathbf{X}^T)^T \\ &= ((\mathbf{X}^T\mathbf{X} + \lambda\mathbf{I})^{-1}\mathbf{X}^T) \sigma^2 I(\mathbf{X}(\mathbf{X}^T\mathbf{X} + \lambda\mathbf{I})^{-T}) = \sigma^2 ((\mathbf{X}^T\mathbf{X} + \lambda\mathbf{I})^{-1}\mathbf{X}^T\mathbf{X}(\mathbf{X}^T\mathbf{X} + \lambda\mathbf{I})^{-T}) \\ &= \sigma^2 ((\mathbf{X}^T\mathbf{X} + \lambda\mathbf{I})^{-1}\mathbf{X}^T\mathbf{X}(\mathbf{X}^T\mathbf{X} + \lambda\mathbf{I})^{-1}), \end{aligned}$$

where we have used that $(\mathbf{X}^T\mathbf{X} + \lambda\mathbf{I})^{-1} = (\mathbf{X}^T\mathbf{X} + \lambda\mathbf{I})^{-T}$ in the last equality (USIKKER PÅ DENNE SISTE! DET BLIR I HVERT FALL HELT RIKTIG Å BEHOLDE -T). In both these equations it is apparent that the moments are equal to those of the OLS estimator when $\lambda = 0$.

b)

The requested moments of $\tilde{f}(\mathbf{x}_0)$ are

$$E(\tilde{f}(\mathbf{x}_0)) = E(\mathbf{x}_0^T \tilde{\beta}) = \mathbf{x}_0^T E(\tilde{\beta}) = \mathbf{x}_0^T (\mathbf{X}^T\mathbf{X} + \lambda\mathbf{I})^{-1}\mathbf{X}^T\mathbf{X}\beta$$

and

$$\begin{aligned} \text{Cov}(\tilde{f}(\mathbf{x}_0)) &= \text{Cov}(\mathbf{x}_0^T \tilde{\beta}) = \mathbf{x}_0^T \text{Cov}(\tilde{\beta}) \mathbf{x}_0 \\ &= \sigma^2 \mathbf{x}_0^T ((\mathbf{X}^T\mathbf{X} + \lambda\mathbf{I})^{-1}\mathbf{X}^T\mathbf{X}(\mathbf{X}^T\mathbf{X} + \lambda\mathbf{I})^{-1}) \mathbf{x}_0. \end{aligned}$$

c)

The expected MSE at \mathbf{x}_0 is

$$\begin{aligned}
E[(y_0 - \tilde{f}(x_0))^2] &= [E(\tilde{f}(x_0) - f(x_0))^2 + \text{Var}(\tilde{f}(x_0)) + \text{Var}(\varepsilon)] \\
&= [E(\tilde{f}(x_0)) - E(f(x_0))]^2 + \text{Cov}(\tilde{f}(x_0)) + \sigma^2 I \\
&= [x_0^T (\mathbf{X}^T \mathbf{X} + \lambda \mathbf{I})^{-1} \mathbf{X}^T \mathbf{X} \beta - x_0^T \beta]^2 + \sigma^2 x_0^T ((\mathbf{X}^T \mathbf{X} + \lambda \mathbf{I})^{-1} \mathbf{X}^T \mathbf{X} (\mathbf{X}^T \mathbf{X} + \lambda \mathbf{I})^{-1}) x_0 + \sigma^2 I
\end{aligned}$$

FOR NOE GRISERI

```
id <- "1X_80KcoYbng1XvYFDirxjEW7LtpNr1m" # google file ID
values <- dget(sprintf("https://docs.google.com/uc?id=%s&export=download", id))
X = values$X
dim(X)
```

```
## [1] 100 81
```

```
x0 = values$x0
dim(x0)
```

```
## [1] 81 1
```

```
beta = values$beta
dim(beta)
```

```
## [1] 81 1
```

```
sigma = values$sigma
sigma
```


```
## [1] 0.5
```

d)

The expression $\tilde{\beta}(\lambda) = ((1 + \lambda)I)^{-1} \hat{\beta}$ is inserted into value below.

```
library(ggplot2)
bias = function(lambda, X, x0, beta) {
  p = ncol(X)
  lambda * diag(p)
  # value = mean(t(x0) %*% solve((1+lambda)*diag(p)) %*% beta - t(x0) %*% beta )^2
  # value = mean(t(x0) %*% solve(diag(p) + lambda*t(X) %*% X) %*% beta - t(x0) %*%
  # beta )^2
  value = mean(t(x0) %*% solve(t(X) %*% X + lambda * diag(p)) %*% t(X) %*% X %*%
    beta - t(x0) %*% beta)^2
  # Tror dette skal være rett algebraisk, men synes resultatet er merkelig kanskje!
  # Kanskje oppgaven ovenfor burde fullføres først? value = mean(t(x0) %*% (diag(p)
  # + 1/lambda*t(X)%*%X)%*%beta - t(x0) %*% beta)^2


  # All of these give different plots!
  return(value)
}
lambdas = seq(0, 2, length.out = 500)
BIAS = rep(NA, length(lambdas))
for (i in 1:length(lambdas)) BIAS[i] = bias(lambdas[i], X, x0, beta)
dfBias = data.frame(lambdas = lambdas, bias = BIAS)
ggplot(dfBias, aes(x = lambdas, y = bias)) + geom_line(color = "red") + xlab(expression(lambda)) +
  ylab(expression(bias^2))
```



Comments: I think this is wrong. Perhaps a good idea to finish c) first, to get a better expression for the bias also. I think the expression is correct however, despite it not being “forkortet”, but this should be done later. However, I think the result is weird.

e)

```
variance = function(lambda, X, x0, sigma) {
  p = ncol(X)
  inv = solve(t(X) %*% X + lambda * diag(p))
  value = sigma * (inv %*% t(X) %*% X %*% inv)
  return(value)
}
lambdas = seq(0, 2, length.out = 500)
VAR = rep(NA, length(lambdas))
for (i in 1:length(lambdas)) VAR[i] = variance(lambdas[i], X, x0, sigma)
dfVar = data.frame(lambdas = lambdas, var = VAR)
ggplot(dfVar, aes(x = lambdas, y = var)) + geom_line(color = "green4") + xlab(expression(lambda)) +
  ylab("variance")
```




Comments: The variance decreases with lambda. It begins very high (which I do not know if holds with the regular estimator when $\lambda = 0$).

f)

```
exp_mse = BIAS + VAR + sigma
lambdas[which.min(exp_mse)]
```

```
## [1] 0.7815631
```

```
dfAll = data.frame(lambda = lambdas, bias = BIAS, var = VAR, exp_mse = exp_mse)
ggplot(dfAll) + geom_line(aes(x = lambda, y = exp_mse), color = "blue") + geom_line(aes(x = lambda,
  y = bias), color = "red") + geom_line(aes(x = lambda, y = var), color = "green4") +
  xlab(expression(lambda)) + ylab(expression(E(MSE)))
```



Comments: Again, unsure if this is correct.

Problem 2

```
# read file
id <- "1yYlEl5gYY3BEtJ4d7KWaFGIOEweJIn_" # google file ID
d.corona <- read.csv(sprintf("https://docs.google.com/uc?id=%s&export=download",
  id), header = T)
```

a)

Inspection of the data. Assuming that 0 = False and 1 = True (as usual), which means that a person has not deceased when `deceased = 0` and vice versa.

```
knitr::kable(table(Deceased = d.corona$deceased)) # Prøve å sette kolonne-navn, men fikk det ikke til
```

Deceased	Freq
0	1905
1	105

```
knitr::kable(table(d.corona$country, d.corona$sex))
```

	female	male
France	60	54
indonesia	30	39
japan	120	174
Korea	879	654

```
knitr::kable(table(d.corona$deceased, d.corona$sex))
```

	female	male
0	1046	859
1	43	62

```
knitr::kable(table(d.corona$country, d.corona$deceased)) # Må hente ut øverste raden herfra, men klart
```

	0	1
France	98	16
indonesia	64	5
japan	283	11
Korea	1460	73

b)

```
# Just in case they are not factors (this should perhaps be deleted later, since
# we could have checked if they were factors earlier).
d.corona$sex = factor(d.corona$sex)
d.corona$country = factor(d.corona$country)
lm.fit <- lm(deceased ~ ., data = d.corona) # perhaps a linear model is not the correct model to use?
```

```
summary(lm.fit)
```

```
##
## Call:
## lm(formula = deceased ~ ., data = d.corona)
##
## Residuals:
##      Min       1Q   Median       3Q      Max
## -0.20383 -0.07105 -0.04495 -0.02110  1.03018
##
## Coefficients:
##              Estimate Std. Error t value Pr(>|t|)
## (Intercept)    0.043862   0.025229   1.739 0.082263 .
## sexmale        0.030815   0.009902   3.112 0.001884 **
## age            0.001305   0.000218   5.984 2.57e-09 ***
## countryindonesia -0.053478   0.033584  -1.592 0.111455
## countryjapan    -0.097525   0.024269  -4.018 6.08e-05 ***
## countryKorea    -0.071966   0.021542  -3.341 0.000851 ***
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## Residual standard error: 0.2193 on 2004 degrees of freedom
## Multiple R-squared:  0.03144,    Adjusted R-squared:  0.02902
## F-statistic: 13.01 on 5 and 2004 DF,  p-value: 1.755e-12
```

- (i) The probability to die of covid for a male age 75 living in Korea can be predicted from the model. The prediction is ...
- (ii) The p-value for `sexmale` is relatively small, but we would not say that there is clear evidence that males have higher probability to die than females. The p-value could be low by chance also, and since it is not amazingly small, we do not think it is appropriate evidence of the question.
- (iii) The p-value of `countryjapan` is the smallest of the country-coefficients, which shows that it could potentially be important as a predictor. In the least, it does not exclude this possibility.
- (iv) Quantify the odds

c)

- (i)
- (ii)

I am guessing that “the appropriate model” would be to use linear regression, as used earlier, since this is the only regression-model we have learned thus far. Not quite sure if they want us to leave out some of the predictors however, or if they just want us to keep all of them in the model.

d)

I will make the models before answering the questions, which can/will be deleted later (since these are multiple choice).

```
# Make data sets.
trainID = sample(x = 1:nrow(d.corona), size = 1500, replace = F) # E.g
train = d.corona[trainID, ]
test = d.corona[!trainID, ]
nrow(d.corona)
```

```
## [1] 2010
```

```
# Ikke så greit å bruke indekser i d.corona tydeligvis. Kan se mer på dette  
# senere dersom det virker som en fornuftig måte å løse oppgaven på...
```

```
library(MASS)  
lda.fit <- lda(deceased ~ ., data = train)  
lda.fit.pred <- predict(lda.fit, newdata = test)$class  
table(predicted = lda.fit.pred, true = test$deceased)
```

```
## < table of extent 2 x 0 >
```

```
# This did not work.
```

EVENTUALLY THERE WILL ONLY BE TRUE OR FALSE IN EACH OF THESE, BUT I HAVE WRITTEN MY THOUGHTS HERE FIRST.

- (i) 5.511811 % seems to be the percentage of deaths in the dataset. This is not equal to the percentage given in the question (by accident). If they were equal I would say that this statement is TRUE. because, as I have understood it, the “null rate” for misclassification can be obtained in this case by always classifying the individual as healthy.
- (ii) Following the argumentation in the above point, LDA is useless if the misclassification rate is higher than approx 5 %. Need to make a LDA in code to test?
- (iii) Can we answer this from theory?
- (iv) The same concern goes for this statement as for the ones above.

Problem 3

```
# read file  
id <- "1i1cQPeoLLC_FyAH0nnqCnnrSBpn05_h0" # google file ID  
diab <- dget(sprintf("https://docs.google.com/uc?id=%s&export=download", id))  
t = MASS::Pima.tr2  
train = diab$ctrain  
test = diab$ctest
```

a)

```
logReg = glm(diabetes ~ ., data = train, family = "binomial")  
summary(logReg)
```

```
##  
## Call:  
## glm(formula = diabetes ~ ., family = "binomial", data = train)  
##  
## Deviance Residuals:  
##      Min       1Q   Median       3Q      Max   
## -2.8155  -0.6367  -0.3211   0.6147   2.2408   
##  
## Coefficients:  
##              Estimate Std. Error z value Pr(>|z|)      
## (Intercept) -10.583538   1.428276  -7.410 1.26e-13 ***  
## npreg        0.105109   0.062721   1.676 0.093775 .    
## glu          0.035586   0.005892   6.039 1.55e-09 ***  
## bp          -0.014654   0.013982  -1.048 0.294615
```

```
## skin          0.020379    0.020575    0.990 0.321962
## bmi           0.094683    0.031265    3.028 0.002458 **
## ped           1.931666    0.529573    3.648 0.000265 ***
## age          0.038291    0.020247    1.891 0.058594 .
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## (Dispersion parameter for binomial family taken to be 1)
##
## Null deviance: 381.91  on 299  degrees of freedom
## Residual deviance: 253.84  on 292  degrees of freedom
## AIC: 269.84
##
## Number of Fisher Scoring iterations: 5
```

(i) (assuming they want us to do it theoretically, and not in R, e.g. by plotting) We have that

$$p_i = \frac{e^{\beta_0 + \beta_1 x_{i1} + \beta_2 x_{i2} + \dots + \beta_7 x_{i7}}}{1 + e^{\beta_0 + \beta_1 x_{i1} + \beta_2 x_{i2} + \dots + \beta_7 x_{i7}}} = \frac{e^{\eta_i(x)}}{1 + e^{\eta_i(x)}} \implies \frac{p_i}{1 - p_i} = \frac{\frac{e^{\eta_i(x)}}{1 + e^{\eta_i(x)}}}{1 - \frac{e^{\eta_i(x)}}{1 + e^{\eta_i(x)}}} = e^{\eta_i(x)}$$

$$\implies \log\left(\frac{p_i}{1 - p_i}\right) = \eta_i(x) = \beta_0 + \beta_1 x_{i1} + \beta_2 x_{i2} + \dots + \beta_7 x_{i7}$$

(ii) Can be seen below

```
glm.probs <- predict(logReg, newdata = test, type = "response")
glm.preds <- ifelse(glm.probs > 0.5, 1, 0) #'Present', 'Non-present') # Could also use these names.
conf.table <- table(predicted = glm.preds, true = test$diabetes)
conf.table
```

```
##           true
## predicted  0   1
##           0 137  29
##           1  18  48
```

The sensitivity is 0.6233766 and the specificity is 0.883871.

b)