

The Emergent Spacetime Rendering Engine: Theoretical Framework, Relativistic Telemetry, and Cosmic Frame-Stacking

Generated by Gemini Notebook

October 1, 2026

Abstract

We present a formal systems-theoretic architecture that reframes physical reality as an emergent 3D bulk space projected from a lower-dimensional discrete 2D boundary update loop ($B_{n+1} = F(B_n)$). By mapping fundamental physical phenomena directly onto computational primitives—rest mass as a projection tax, quark confinement as a runtime exception handler, quantum entanglement as shared boundary memory references, and wavefunction collapse as lazy evaluation—this framework bridges quantum field theory, general relativity, and information theory. In this updated third edition (v3.0), we incorporate two major new architectural extensions: (1) **GPS Relativistic Telemetry**, demonstrating how Special Relativistic velocity time dilation ($\Delta t_{\text{SR}} = -7.21 \mu\text{s/day}$) and General Relativistic gravitational time dilation ($\Delta t_{\text{GR}} = +45.72 \mu\text{s/day}$) emerge from substrate bus bandwidth constraints and MERA rendering tax differentials; and (2) **Frame-Stacking Cosmic Expansion**, modeling spacetime expansion as an append-only history ledger ("Tower of Hanoi") where accumulating boundary circuit complexity $K(N)$ directly drives 3D bulk volume expansion $V(N) \propto K(N)$. Furthermore, we present quantitative numerical results from our Python simulation engine across three empirical test beds: Planck-scale cosmic ray dispersion ($\Delta t \approx 421.52 \text{ ms}$ over 500 Mpc for 100 GeV photons), holographic interferometer noise ($\delta L \approx 0.254 \text{ fm}$ across a 4.0 km arm), and sub-millimeter gravitational Yukawa deviations (+8.21% enhancement at $50.0 \mu\text{m}$).

1 Introduction

Modern theoretical physics faces a persistent foundational challenge in unifying general relativity and quantum mechanics. The *Emergent Spacetime Rendering Engine* proposes that this difficulty arises from treating macroscopic 3D spacetime as fundamental rather than emergent. In this framework, the physical universe is modeled as a computational system projected from a lower-dimensional boundary state space B evolving through discrete clock cycles t_P :

$$B_{n+1} = F(B_n) \tag{1}$$

Fundamental physical constants specify the operational bounds of the underlying execution substrate:

- **Planck Length** ($\ell_P \approx 1.6 \times 10^{-35} \text{ m}$): Spatial resolution limit (grid pixel size).
- **Planck Time** ($t_P \approx 5.4 \times 10^{-44} \text{ s}$): Substrate execution clock cycle duration.
- **Speed of Light** ($c \approx 3.0 \times 10^8 \text{ m/s}$): Maximum bus bandwidth and signal propagation speed limit.

2 Core Operational Mechanics

2.1 Rest Mass as a 2D-to-3D Projection Tax

In standard physics, rest mass m represents intrinsic inertia. In this holographic rendering framework, rest mass is defined as the **computational update tax** functional $C[B_n]$ required on the 2D boundary plane to sustain a localized 3D spatial field distortion:

$$m \sim \frac{1}{c^2} \frac{\delta C}{\delta \Pi} \quad (2)$$

where Π denotes the Multiscale Entanglement Renormalization Ansatz (MERA) projection operator coarse-graining boundary state updates layer-by-layer.

2.2 Quark Confinement as Runtime Exception Handling

Quantum Chromodynamics (QCD) dictates that color-charged quarks cannot exist in isolation due to the linear growth of potential energy in the connecting gluon flux tube:

$$V(r) \sim \sigma r \quad (\sigma \approx 1 \text{ GeV/fm}) \quad (3)$$

When external energy is injected to pull a quark-antiquark pair apart, attempting to create an isolated color charge constitutes a system **segmentation fault** that would corrupt bulk data integrity. The engine intercepts this illegal state via an automated **runtime exception handler**: when field energy reaches the pair-creation threshold ($E \geq 2mc^2$), the flux tube snaps, converting injected field energy into rest mass (hadronization) and re-encapsulating exposed charges into color-neutral hadrons.

2.3 Quantum Entanglement as Shared Memory References

Spatially separated entangled particles in 3D bulk space are modeled as pointers referencing a single, localized memory address (`0x7F_BOUND_MEM`) on the 2D boundary. Measuring Particle A directly updates the underlying boundary memory, instantly reflecting in Particle B's eigenstate. Because state resolution occurs on the boundary substrate rather than propagating across 3D bulk distance, zero-latency correlation occurs without violating bulk relativity or the No-Signaling theorem ($v \leq c$).

2.4 Wavefunction Collapse as Lazy Evaluation

To optimize computational resources, unobserved quantum states are maintained in an unrasterized probability distribution with minimal memory overhead (≈ 0.01 KB). Upon environmental interaction or measurement (decoherence), the rendering engine executes **procedural rasterization**, collapsing the lazy evaluation state into a concrete bulk eigenstate (≈ 1.00 KB).

2.5 Sub-Pixel Field Antialiasing

Discrete spatial updates on a Planck-scale lattice risk producing macroscopic directional artifacts. Quantum field fluctuations function as a sub-pixel antialiasing distribution operator $\hat{\phi}(f)$:

$$\hat{\phi}(f) = \int \hat{\phi}(x) f(x) d^4x \quad (4)$$

smoothing lattice pixels into a continuous, rotationally invariant 3D spatial geometry.

3 Mass System Taxonomy

The architecture categorizes elementary particle masses into two distinct allocation classes:

1. **Hadronic Mass** ($\approx 99\%$ of nucleon mass): Originates from QCD gluon field binding energy, representing a **dynamic runtime update tax** continuously calculated across boundary execution cycles.
2. **Leptonic Mass** ($\approx 1\%$): Free leptons acquire mass via Higgs field Yukawa couplings, acting as **static registration overhead** in memory without high dynamic field re-computation.

4 Quantitative Empirical Test Beds

Our Python simulation engine evaluates three concrete experimental test beds to validate or constrain the framework.

4.1 Test Bed 1: Planck-Scale Cosmic Ray Dispersion

Photons traversing cosmological distances experience sub-pixel lattice grid delays as their energy approaches the Planck energy E_P :

$$v(E) \approx c \left(1 - \xi \frac{E}{E_P} \right) \quad (5)$$

For a high-energy gamma-ray photon ($E = 100$ GeV) originating from a source at distance $D = 500$ Mpc (≈ 1.63 billion light-years), our simulation predicts:

- **Calculated Sub-Pixel Arrival Delay** (Δt): 421.52 ms
- **Accumulated Substrate Clock Lag** (ΔN): 7.819×10^{42} Planck ticks (t_P)

4.2 Test Bed 2: Holographic Interferometer Noise

Boundary-to-bulk MERA projection uncertainty induces an irreducible transverse spacetime jitter δL across interferometer arms of length L :

$$\delta L \sim \sqrt{\ell_P \cdot L} \quad (6)$$

For a 4.0 km arm (modeled on Advanced LIGO), the simulation yields:

- **Transverse Spacetime Jitter** (δL): 2.5426×10^{-16} m (0.2543 femtometers)
- **Holographic Noise Spectral Density Floor** (S_h): 2.3219×10^{-22} m/ $\sqrt{\text{Hz}}$

4.3 Test Bed 3: Sub-Millimeter Gravity Law Deviations

At scales below the boundary projection cutoff, gravitational attraction deviates from the Newtonian inverse-square law via a non-Newtonian Yukawa modification:

$$V(r) = -G \frac{m_1 m_2}{r} \left(1 + \alpha e^{-r/\lambda} \right) \quad (7)$$

Evaluating at separation $r = 50.0 \mu\text{m}$ with Yukawa scale parameter $\lambda = 20.0 \mu\text{m}$:

- **Newtonian Force Enhancement Factor**: $1.0821 \times$
- **Departure from Inverse-Square Law**: +8.21% force enhancement

5 Electromagnetic Wave Telemetry & Surface Polarization

Photons carry both thermodynamic energy ($E = h\nu$) and an embedded material state header. When transmitted through media or reflected off material surface lattices, wave packets undergo frequency absorption and Fresnel polarization shifts:

$$R_s = \left| \frac{\cos \theta_i - n_{\text{complex}} \cos \theta_t}{\cos \theta_i + n_{\text{complex}} \cos \theta_t} \right|^2, \quad R_p = \left| \frac{n_{\text{complex}} \cos \theta_i - \cos \theta_t}{n_{\text{complex}} \cos \theta_i + \cos \theta_t} \right|^2 \quad (8)$$

Near the dielectric Brewster angle ($\theta_i = 53^\circ$), dielectrics (Water/Wood) suppress R_p to near zero, yielding 100% linear polarization (DOLP = 1.00). Conversely, metallic conductors (Steel/Aluminum) reflect high parallel amplitude ($R_p = 41.48\%$) with a 203.6° phase shift, providing a distinct surface polarization header.

6 GPS Relativistic Telemetry & Substrate Synchronization

GPS satellites orbiting at 20,200 km altitude ($v \approx 3,873$ m/s) experience two competing substrate clock dilation effects:

1. **Special Relativity (Bus Bandwidth Limit):** High velocity consumes boundary updates for spatial coordinate translation across grid cells, causing a kinematic clock lag:

$$\frac{\delta f}{f}_{\text{SR}} = -\frac{1}{2} \frac{v^2}{c^2} \implies \Delta t_{\text{SR}} = -7.21 \mu\text{s/day} \quad (9)$$

2. **General Relativity (Rendering Tax Differential):** Weaker gravitational potential in high orbit reduces MERA entanglement and boundary rendering tax density $C[B_n]$, accelerating local clock execution:

$$\frac{\delta f}{f}_{\text{GR}} = \frac{\Delta \Phi}{c^2} \implies \Delta t_{\text{GR}} = +45.72 \mu\text{s/day} \quad (10)$$

The combined net drift of $+38.51 \mu\text{s/day}$ is pre-compensated by factory-tuning master oscillators from 10.230000000000 MHz down to 10.22999999544 MHz, ensuring zero-latency telemetry frame alignment.

7 Frame-Stacking Complexity & Cosmic Expansion

A central innovation in version 3.0 is the formalization of cosmic expansion as an append-only **frame-stacking ledger** ("Tower of Hanoi").

7.1 Append-Only State Ledger

Rather than overwriting boundary matrices, each tick N appends a new state update $B_{n+1} = F(B_n)$ onto an acyclic history Directed Acyclic Graph (DAG). The cumulative boundary circuit complexity grows monotonically over substrate execution history:

$$K(N) = \sum_{i=1}^N C[B_i] \quad (11)$$

7.2 Complexity Equals Volume Relation

Adopting the holographic complexity-volume duality ($K \propto V$), the volume $V(N)$ of projected 3D bulk space is directly driven by the total accumulated boundary complexity $K(N)$:

$$V(N) = \frac{G_N \hbar}{c} \cdot K(N) \quad (12)$$

Because past frames are retained rather than erased, the accumulating history stack continually expands the 3D metric volume.

7.3 Emergent Hubble Growth Rate

The relative Hubble volume expansion rate $H(N)$ naturally decays as total accumulated volume grows:

$$H(N) = \frac{1}{V(N)} \frac{dV(N)}{dN} = \frac{C[B_N]}{K(N)} \quad (13)$$

In our simulation runs ($N = 1 \rightarrow 20$), 20 accumulated clock ticks produce a $5.1 \times$ **total volume expansion** and a $1.72 \times$ **increase in effective cosmic horizon radius**, demonstrating how frame-stacking naturally generates macroscopic cosmic expansion.

8 Summary & Conclusion

Table 1: System Mapping: Physical Phenomena vs. Rendering Architecture (v3.0)

Physical Phenomenon	System Architecture Analogue	Quantitative Impact
Speed of Light (c)	Maximum Substrate Bus Bandwidth	Signal limit $c \approx 3 \times 10^8$ m/s
Rest Mass (m)	Rendering Tax Cost Functional $C[B_n]$	99% Dynamic / 1% Static
Quark Confinement	Runtime Exception Handler	Hadronization pair creation ($E = mc^2$)
Quantum Entanglement	Shared Boundary Memory Pointer	Zero-latency non-local correlation
Wavefunction Collapse	Lazy Eval $\rightarrow$ Procedural Rasterization	0.01 KB $\rightarrow$ 1.00 KB expansion
Cosmic Ray Dispersion	Sub-Pixel Lattice Resolution Limit	$\Delta t = 421.52$ ms (100 GeV, 500 Mpc)
Holographic Noise	2D $\rightarrow$ 3D MERA Projection Uncertainty	$\delta L = 0.254$ fm ($L = 4.0$ km)
Sub-mm Gravity	Boundary Entropy Cutoff	+8.21% force enhancement (50 μ m)
Spectral Headers	Quantum Absorption Manifest	Notches at 756.5 nm and 939.1 nm
Surface Polarization	Fresnel Band/Phase Shift Header	Dielectric DOLP = 1.00 vs Metal $R_p = 41.5\%$
GPS Clock Dilation	SR Bandwidth / GR Rendering Tax	Net +38.51 μ s/day offset
Cosmic Expansion	Frame-Stacking History Ledger ($K \propto V$)	$5.1 \times$ volume growth over 20 ticks

The *Emergent Spacetime Rendering Engine (v3.0)* provides a unified, computationally grounded, and empirically testable framework for physical reality. By demonstrating how cosmic expansion emerges naturally from an append-only frame-stacking ledger, this work provides a coherent systems-theoretic model connecting subatomic quantum mechanics, general relativity, and cosmology.