dynamics, and a small one to a few spirals that meander around the system. The number of clusters is seen to decrease with the strength of the activator diffusion and increase with noise intensity. A dynamical state with many clusters can be characterised by their size distribution, which is observed to scale with a power law, in agreement with recent results obtained for thermal waves in subexcitable media.

We thank M. BÄR for suggestions, and P. Jung for fruitful and enjoyable discussions on the dynamics of excitable media. We acknowledge the Deutsche Forschungsgemeinschaft (Sfb555) for financial support. JGO also thanks the A. von Humboldt-Stiftung and the Dirección General de Investigación Científica y Técnica (Spain, project PB96-0241).

REFERENCES

- Jung P. and Mayer-Kress G., Chaos, 5 (1995) 458; Jung P. and Mayer-Kress G., Phys. Rev. Lett., 74 (1995) 2134.
- Jung P., Phys. Rev. Lett., 78 (1997) 1723; Jung P., Cornell-Bell A., Madden K. and Moss F., J. Neurophysiol., 79 (1998) 1098.
- [3] SENDIÑA-NADAL I., MUÑUZURI A., VIVES D., PÉREZ-MUÑUZURI V., CASADEMUNT J., RAMÍREZ-PISCINA L., SANCHO J. M. and SAGUÉS F., Phys. Rev. Lett., 80 (1998) 5437.
- [4] KÁDÁR S., WANG J. and SHOWALTER K., Nature, 391 (1998) 770.
- [5] HOU Z., YANG L., XIAOBIN Z. and XIN H., Phys. Rev. Lett., 81 (1998) 2854.
- [6] Hempel H., Schimansky-Geier L. and García-Ojalvo J., Phys. Rev. Lett., 82 (1999) 3713.
- [7] WINFREE A. T., Chaos, 1 (1991) 303.
- [8] TABOADA J. J., MUÑUZURI A. P., PÉREZ-MUÑUZURI V., GÓMEZ-GESTEIRA M. and PÉREZ-VILLAR V., Chaos, 4 (1994) 519; MARKUS M., KLOSS G. and KUSCH I., Nature, 371 (1994) 402; NAGY-UNGVARAI Z. and MÜLLER S., Int. J. Bifurcation Chaos, 4 (1994) 1257; OUYANG Q. and FLESSELLES J.-M., Nature, 379 (1996) 143.
- [9] HILDEBRAND M., BÄR M. and EISWIRTH M., Phys. Rev. Lett., **75** (1995) 1503.
- [10] STRAIN M. C. and GREENSIDE H. S., Phys. Rev. Lett., 80 (1998) 2306.
- [11] KARMA A., Phys. Rev. Lett., 71 (1993) 1103; KARMA A., Chaos, 4 (1994) 461; see also Focus Issue: Fibrillation in Normal Ventricular Myocardium, Chaos, 8 (1998) pp. 1-306.
- [12] Ito H. and Glass L., Phys. Rev. Lett., 66 (1991) 671.
- [13] HEMPEL H., FRICKE T., MIETH M. and SCHIMANSKY-GEIER L., in *Nonlinear Physics of Complex Systems*, edited by J. Parisi, S. Müller and W. Zimmermann (Springer) 1996.
- [14] PANFILOV A. V. and HOLDEN A. V., Phys. Lett. A, 151 (1990) 23; COURTEMANCHE M. and WINFREE A. T., Int. J. Bifurcation Chaos, 1 (1991) 431.
- [15] BÄR M. and EISWIRTH M., Phys. Rev. E, 48 (1993) 1635; BÄR M. and OR-GUIL M., Phys. Rev. Lett., 82 (1999) 1160.
- [16] PANFILOV A. and HOGEWEG P., Phys. Lett. A, 176 (1993) 295; MARÉE A. F. M. and PANFILOV A. V., Phys. Rev. Lett., 78 (1997) 1819.
- [17] BIKTASHEV V. N., HOLDEN A. V., TSYGANOV M. A., BRINDLEY J. and HILL N. A., Phys. Rev. Lett., 81 (1998) 2815.
- [18] Barkley D., Kness M. and Tuckerman L. S., Phys. Rev. A, 42 (1990) 2489.
- [19] ABRAMOWITZ M. and STEGUN I. A., Handbook of Mathematical Functions (Dover) 1972.
- [20] Barkley D., *Physica D*, **49** (1991) 61.
- [21] GARCÍA-OJALVO J. and SANCHO J. M., Noise in Spatially Extended Systems (Springer) 1999.
- [22] Aranson I. S., Chaté H. and Tang L.-H., Phys. Rev. Lett., 80 (1998) 2646.
- [23] WANG J., KÁDÁR S., JUNG P. and SHOWALTER K., Phys. Rev. Lett., 82 (1999) 855.