

Level M Regression Models

Lecture 12

Colette Mair

School of Mathematics and Statistics



Inference

Introduction

- ▶ Now we will focus on the construction interval estimates for various parameters of our models

Confidence Interval

A 100c% confidence interval estimate for $\mathbf{b}^T \beta$ with confidence c is

$$\mathbf{b}^T \hat{\beta} \pm t \left(n - p; \frac{1 + c}{2} \right) \sqrt{\frac{RSS}{n - p} (\mathbf{b}^T (\mathbf{X}^T \mathbf{X})^{-1} \mathbf{b})}.$$

Prediction Interval

A 100c% prediction interval for a future observation y_f is

$$\mathbf{b}^T \hat{\beta} \pm t \left(n - p; \frac{1 + c}{2} \right) \sqrt{\frac{RSS}{n - p} (1 + \mathbf{b}^T (\mathbf{X}^T \mathbf{X})^{-1} \mathbf{b})}.$$

Confidence c

- ▶ In this course, we will assume $c = 0.95$ unless otherwise specified.

95% Confidence Interval

A 95% confidence interval estimate for $\mathbf{b}^T \beta$ is

$$\mathbf{b}^T \hat{\beta} \pm t(n - p; 0.975) \sqrt{\frac{RSS}{n - p} (\mathbf{b}^T (\mathbf{X}^T \mathbf{X})^{-1} \mathbf{b})}.$$

95% Prediction Interval

A 95% prediction interval for a future observation y_f is

$$\mathbf{b}^T \hat{\beta} \pm t(n-p; 0.975) \sqrt{\frac{RSS}{n-p} (1 + \mathbf{b}^T (\mathbf{X}^T \mathbf{X})^{-1} \mathbf{b})}.$$

Illustration

$$y_i = \alpha + \beta x_i \quad i = 1, \dots, n.$$

A 95% confidence interval estimate for β is

► $\beta^T = (\alpha \quad \beta)$

► $\mathbf{b}^T = (0 \quad 1)$

$\mathbf{b}^T \beta$ is

$$\mathbf{b}^T \hat{\beta} \pm t(n-p; 0.975) \sqrt{\frac{RSS}{n-p} (\mathbf{b}^T (\mathbf{X}^T \mathbf{X})^{-1} \mathbf{b})}.$$

Example


Autoanalyser

- ▶ Blood plasma concentrations are usually measured using a lengthy laboratory process.
- ▶ The autoanalyser is regularly tested to see if it is performing properly.
- ▶ A method using an autoanalyser is often used
- ▶ On this occasion, 12 measurements have been made on samples of known concentration (3 replicates at each of 4 concentrations).

Autoanalyser

- ▶ Construct a 95% C.I. for the population mean autoanalyser concentration when the true concentration is 6 units and interpret the interval.
- ▶ Construct a 95% prediction interval for the autoanalyser concentration when the true concentration is 6 units and interpret the interval.

Autoanalyser



Autoanalyser

$$\text{autoanalyser}_i = \beta_0 + \beta_1 \text{true}_i + \epsilon_i, \quad i = 1, \dots, 12$$


Autoanalyser

Simple linear model R output


term	estimate	std.error	statistic	p.value
(Intercept)	0.68	0.19	3.60	0.00487
true	0.85	0.04	20.75	0.00000

Autoanalyser


Residuals vs Fitted




Normal Q-Q



Cook's distance



Residuals vs Leverage



Autoanalyser

$$\begin{aligned}\mathbf{b}^T \hat{\beta} &= \hat{\alpha} + 6 \times \hat{\beta} \\ &= 0.683 + 0.85 \times 6 \\ &= 5.783\end{aligned}$$


$$RSS = 0.252$$

$$n = 12$$

$$t(10;0.975) = 2.228$$

$$(\mathbf{X}^T \mathbf{X})^{-1} = \frac{1}{180} \begin{pmatrix} 258 & -54 \\ -54 & 12 \end{pmatrix}$$

Autoanalyser



Autoanalyser

- ▶ Construct a 95% C.I. for the population mean autoanalyser concentration when the true concentration is 6 units and interpret the interval.
- ▶ Construct a 95% prediction interval for the autoanalyser concentration when the true concentration is 6 units and interpret the interval.

95% C.I.

95% prediction interval

Trees

- ▶ The dataset refers to the volume (cubic feet) and diameter (inches) (at 54 inches above the ground) and height (feet) for a sample of 31 black cherry trees in the Allegheny National Forest Pennsylvania.
- ▶ The data were collected in order to and an estimate for the volume of a tree (and therefore for the timber yield), given its height and diameter. A

Trees

Our full model for the trees data is

$$E(Y) = \alpha + \beta x_1 + \gamma x_2$$

where Y denotes $\log(\text{volume})$, x_1 denotes $\log(\text{diameter})$ and x_2 denotes $\log(\text{height})$ of 31 trees.

Trees

The fitted model produced:

$$\hat{\beta} = \begin{pmatrix} -6.632 \\ 1.983 \\ 1.117 \end{pmatrix}$$

$$RSS = 0.1855$$

$$(\mathbf{X}^T \mathbf{X})^{-1} = \begin{pmatrix} 96.5721 & 3.1393 & -24.1651 \\ 3.1393 & 0.8495 & -1.2275 \\ -24.1651 & -1.2275 & 6.3099 \end{pmatrix}$$

Trees

Construct 95% confidence intervals for β and γ to test their significance.

95% C.I. for β

95% C.I. for γ