

# Inner Minkowski Dimension of Products of Fractal Strings

Isaac Ashton, Joseph Ingenito, Peter Tonuzi

The College of New Jersey



# What We Will Cover

we're finishing this after our dry run and inevitable cuts



# What is a Fractal String

## Definition 1

A **fractal string**  $\Omega$  is a bounded open subset of the real line.

# What is a Fractal String

## Definition 1

A **fractal string**  $\Omega$  is a bounded open subset of the real line.

$$\Omega = \bigcup_{j=1}^{\infty} \ell_j, \quad \mathcal{L} = \text{Vol}_1(\Omega) = \sum_{j=1}^{\infty} \ell_j \implies \lim_{j \rightarrow \infty} \ell_j = 0.$$



# What is a Fractal String

## Definition 1

A **fractal string**  $\Omega$  is a bounded open subset of the real line.

$$\Omega = \bigcup_{j=1}^{\infty} \ell_j, \quad \mathcal{L} = \text{Vol}_1(\Omega) = \sum_{j=1}^{\infty} \ell_j \implies \lim_{j \rightarrow \infty} \ell_j = 0.$$

$$\ell_1 \geq \ell_2 \geq \ell_3 \geq \cdots \geq 0$$

# What is a Fractal String

## Definition 1

A **fractal string**  $\Omega$  is a bounded open subset of the real line.

$$\Omega = \bigcup_{j=1}^{\infty} \ell_j, \quad \mathcal{L} = \text{Vol}_1(\Omega) = \sum_{j=1}^{\infty} \ell_j \implies \lim_{j \rightarrow \infty} \ell_j = 0.$$

$$\ell_1 \geq \ell_2 \geq \ell_3 \geq \cdots \geq 0$$

The fractal string we studied the most in particular is the **Cantor String**



# Cantor String

# Inner-Tubular Neighborhood

## Definition 2

**Inner-tubular neighborhood:**  $V(\epsilon) = \text{vol}_1\{x \in \Omega \mid d(x, \partial\Omega) < \epsilon\}$

# Inner-Tubular Neighborhood

## Definition 2

**Inner-tubular neighborhood:**  $V(\epsilon) = \text{vol}_1\{x \in \Omega \mid d(x, \partial\Omega) < \epsilon\}$

**Counting function:**  $N_\Omega(x) = \sum_{\ell_j \geq x} 1$

**More Notation:**  $\mathcal{W}_\Omega(x) = \sum_{j: \ell_j < x} \ell_j$

**Now we can write:**  $V(\epsilon) = 2\epsilon \cdot N_\Omega(2\epsilon) + \mathcal{W}_\Omega(2\epsilon)$



# Cantor String Volume



# Inner Minkowski Dimension of a Fractal String

## Definition 3

**Dimension of  $\Omega$ :**  $D_\Omega = \inf \{ \alpha \geq 0 \mid V(\epsilon) = O(\epsilon^{1-\alpha}) \text{ as } \epsilon \rightarrow 0^+ \}$

# Inner Minkowski Dimension of a Fractal String

## Definition 3

**Dimension of  $\Omega$ :**  $D_\Omega = \inf \{ \alpha \geq 0 \mid V(\epsilon) = O(\epsilon^{1-\alpha}) \text{ as } \epsilon \rightarrow 0^+ \}$

**In other terms:**  $D_\Omega = \text{the smallest } \alpha \text{ st } \lim_{\epsilon \rightarrow 0^+} \frac{V(\epsilon)}{\epsilon^{1-\alpha}} = c, \ c \in \mathbb{R}$

# Inner Minkowski Dimension of a Fractal String

## Definition 3

**Dimension of  $\Omega$ :**  $D_\Omega = \inf \{ \alpha \geq 0 \mid V(\epsilon) = O(\epsilon^{1-\alpha}) \text{ as } \epsilon \rightarrow 0^+ \}$

**In other terms:**  $D_\Omega = \text{the smallest } \alpha \text{ st } \lim_{\epsilon \rightarrow 0^+} \frac{V(\epsilon)}{\epsilon^{1-\alpha}} = c, \ c \in \mathbb{R}$

**Cantor String Dimension:**  $D_{cs} = \log_3 2$

# Fractal Lawn

## Definition 4

A **fractal lawn**  $\Omega^2$  is the Cartesian product of two fractal strings  
 $\Omega^2 = \Omega_1 \times \Omega_2$ .

# Fractal Lawn

## Definition 4

A **fractal lawn**  $\Omega^2$  is the Cartesian product of two fractal strings  
 $\Omega^2 = \Omega_1 \times \Omega_2$ .

The Cartesian product of two fractal strings is a bounded open subset of  $\mathbb{R}^2$  since a fractal string is a bounded open subset of  $\mathbb{R}$ .



# Fractal Lawn

## Definition 4

A **fractal lawn**  $\Omega^2$  is the Cartesian product of two fractal strings  
 $\Omega^2 = \Omega_1 \times \Omega_2$ .

The Cartesian product of two fractal strings is a bounded open subset of  $\mathbb{R}^2$  since a fractal string is a bounded open subset of  $\mathbb{R}$ .

$$\Omega_1 = \bigcup_{j=1}^{\infty} \ell_j, \quad \Omega_2 = \bigcup_{i=1}^{\infty} p_i$$



# Fractal Lawn

## Definition 4

A **fractal lawn**  $\Omega^2$  is the Cartesian product of two fractal strings  
 $\Omega^2 = \Omega_1 \times \Omega_2$ .

The Cartesian product of two fractal strings is a bounded open subset of  $\mathbb{R}^2$  since a fractal string is a bounded open subset of  $\mathbb{R}$ .

$$\Omega_1 = \bigcup_{j=1}^{\infty} \ell_j, \quad \Omega_2 = \bigcup_{i=1}^{\infty} p_i$$

$$\ell_1 \geq \ell_2 \geq \ell_3 \geq \cdots \geq 0, \quad p_1 \geq p_2 \geq p_3 \geq \cdots \geq 0$$



# Cantor Lawn



# Fractal Lawn Volume I

**Inner-tubular neighborhood:**  $V(\epsilon) = \text{vol}_2\{x \in \Omega^2 \mid d(x, \partial\Omega^2) < \epsilon\}$



# Fractal Lawn Volume I

**Inner-tubular neighborhood:**  $V(\epsilon) = \text{vol}_2\{x \in \Omega^2 \mid d(x, \partial\Omega^2) < \epsilon\}$

- $\partial\Omega^2 = (\partial\Omega_1 \times \Omega_2) \cup (\Omega_1 \times \partial\Omega_2)$

# Fractal Lawn Volume I

**Inner-tubular neighborhood:**  $V(\epsilon) = \text{vol}_2\{x \in \Omega^2 \mid d(x, \partial\Omega^2) < \epsilon\}$

- $\partial\Omega^2 = (\partial\Omega_1 \times \Omega_2) \cup (\Omega_1 \times \partial\Omega_2)$

**Counting function:** Counts up to the critical length for any  $x > 0$

# Fractal Lawn Volume I

**Inner-tubular neighborhood:**  $V(\epsilon) = \text{vol}_2\{x \in \Omega^2 \mid d(x, \partial\Omega^2) < \epsilon\}$

- $\partial\Omega^2 = (\partial\Omega_1 \times \Omega_2) \cup (\Omega_1 \times \partial\Omega_2)$

**Counting function:** Counts up to the critical length for any  $x > 0$

- $N_{\Omega_i}(x) = \sum_{j: \ell_j \geq x} 1$

# Fractal Lawn Volume I

**Inner-tubular neighborhood:**  $V(\epsilon) = \text{vol}_2\{x \in \Omega^2 \mid d(x, \partial\Omega^2) < \epsilon\}$

- $\partial\Omega^2 = (\partial\Omega_1 \times \Omega_2) \cup (\Omega_1 \times \partial\Omega_2)$

**Counting function:** Counts up to the critical length for any  $x > 0$

- $N_{\Omega_i}(x) = \sum_{j: \ell_j \geq x} 1$

**Useful Definitions:**

# Fractal Lawn Volume I

**Inner-tubular neighborhood:**  $V(\epsilon) = \text{vol}_2\{x \in \Omega^2 \mid d(x, \partial\Omega^2) < \epsilon\}$

- $\partial\Omega^2 = (\partial\Omega_1 \times \Omega_2) \cup (\Omega_1 \times \partial\Omega_2)$

**Counting function:** Counts up to the critical length for any  $x > 0$

- $N_{\Omega_i}(x) = \sum_{j: \ell_j \geq x} 1$

**Useful Definitions:**

- $\mathcal{L}_i = \sum_{j \in \mathbb{N}} \ell_j$

# Fractal Lawn Volume I

**Inner-tubular neighborhood:**  $V(\epsilon) = \text{vol}_2\{x \in \Omega^2 \mid d(x, \partial\Omega^2) < \epsilon\}$

- $\partial\Omega^2 = (\partial\Omega_1 \times \Omega_2) \cup (\Omega_1 \times \partial\Omega_2)$

**Counting function:** Counts up to the critical length for any  $x > 0$

- $N_{\Omega_i}(x) = \sum_{j: \ell_j \geq x} 1$

**Useful Definitions:**

- $\mathcal{L}_i = \sum_{j \in \mathbb{N}} \ell_j$

- $\mathcal{W}_i(x) = \sum_{j: \ell_j < x} \ell_j$

# Fractal Lawn Volume I

**Inner-tubular neighborhood:**  $V(\epsilon) = \text{vol}_2\{x \in \Omega^2 \mid d(x, \partial\Omega^2) < \epsilon\}$

- $\partial\Omega^2 = (\partial\Omega_1 \times \Omega_2) \cup (\Omega_1 \times \partial\Omega_2)$

**Counting function:** Counts up to the critical length for any  $x > 0$

- $N_{\Omega_i}(x) = \sum_{j: \ell_j \geq x} 1$

**Useful Definitions:**

- $\mathcal{L}_i = \sum_{j \in \mathbb{N}} \ell_j$

- $\mathcal{W}_i(x) = \sum_{j: \ell_j < x} \ell_j$

- $\sum_{j=1}^{N_{\Omega_i}(x)} \ell_j = \mathcal{L}_i - \mathcal{W}_i(x)$



# Fractal Lawn Volume II

## Partially Covered Volume Box



# Fractal Lawn Volume II

## Partially Covered Volume Box



### ■ Partially covered volume

$$V_{partial}(\epsilon) = 2\epsilon N_{\Omega_1}(2\epsilon) \sum_{j=1}^{N_{\Omega_2}(2\epsilon)} \ell_j + 2\epsilon N_{\Omega_2}(2\epsilon) \sum_{j=1}^{N_{\Omega_1}(2\epsilon)} (p_j - 2\epsilon)$$

# Fractal Lawn Volume II

## Partially Covered Volume Box



- Partially covered volume

- $V_{partial}(\epsilon) = 2\epsilon N_{\Omega_1}(2\epsilon) \sum_{j=1}^{N_{\Omega_2}(2\epsilon)} \ell_j + 2\epsilon N_{\Omega_2}(2\epsilon) \sum_{j=1}^{N_{\Omega_1}(2\epsilon)} (p_j - 2\epsilon)$

- Fully covered volume

- $V_{full}(\epsilon) = \mathcal{L}_1 \mathcal{W}_2(2\epsilon) + \mathcal{L}_2 \mathcal{W}_1(2\epsilon)$

# Fractal Lawn Volume II

## Partially Covered Volume Box



### ■ Partially covered volume

$$V_{partial}(\epsilon) = 2\epsilon N_{\Omega_1}(2\epsilon) \sum_{j=1}^{N_{\Omega_2}(2\epsilon)} \ell_j + 2\epsilon N_{\Omega_2}(2\epsilon) \sum_{j=1}^{N_{\Omega_1}(2\epsilon)} (p_j - 2\epsilon)$$

### ■ Fully covered volume

$$V_{full}(\epsilon) = \mathcal{L}_1 \mathcal{W}_2(2\epsilon) + \mathcal{L}_2 \mathcal{W}_1(2\epsilon)$$

### ■ Double counted volume

$$V_{double}(\epsilon) = \mathcal{W}_1(2\epsilon) \mathcal{W}_2(2\epsilon)$$

# Fractal Lawn Volume III

$$V(\epsilon) = V_{partial}(\epsilon) + V_{full}(\epsilon) - V_{double}(\epsilon)$$



# Fractal Lawn Volume III

$$V(\epsilon) = V_{partial}(\epsilon) + V_{full}(\epsilon) - V_{double}(\epsilon)$$

$$V(\epsilon) = 2\epsilon \cdot N_{\Omega_1}(2\epsilon) \cdot \left( \sum_{j=1}^{N_{\Omega_2}(2\epsilon)} p_j \right) + 2\epsilon \cdot N_{\Omega_2}(2\epsilon) \cdot \left( \sum_{j=1}^{N_{\Omega_1}(2\epsilon)} (\ell_j - 2\epsilon) \right) +$$

$$\mathcal{L}_1 \mathcal{W}_2(2\epsilon) + \mathcal{L}_2 \mathcal{W}_1(2\epsilon) - \mathcal{W}_1(2\epsilon) \mathcal{W}_2(2\epsilon)$$

$$= 2\epsilon \cdot N_{\Omega_1} \cdot (\mathcal{L}_2 - \mathcal{W}_2) + 2\epsilon \cdot N_{\Omega_2} \cdot (\mathcal{L}_1 - \mathcal{W}_1 - 2\epsilon \cdot N_{\Omega_1}) + \\ \mathcal{L}_1 \mathcal{W}_2 + \mathcal{L}_2 \mathcal{W}_1 - \mathcal{W}_1 \mathcal{W}_2$$

# Fractal Lawn Volume IV

$$\begin{aligned} V(\epsilon) &= \mathcal{L}_1 \cdot (2\epsilon \cdot N_{\Omega_2} + \mathcal{W}_2) + \mathcal{L}_2 \cdot (2\epsilon \cdot N_{\Omega_1} + \mathcal{W}_1) - \\ &\quad (2\epsilon \cdot N_{\Omega_1} + \mathcal{W}_1) \cdot (2\epsilon \cdot N_{\Omega_2} + \mathcal{W}_2) \\ &= \mathcal{L}_1 \cdot (2\epsilon \cdot N_{\Omega_2} + \mathcal{W}_2) + \mathcal{L}_2 \cdot (2\epsilon \cdot N_{\Omega_1} + \mathcal{W}_1) - \\ &\quad (2\epsilon \cdot N_{\Omega_1} + \mathcal{W}_1) \cdot (2\epsilon \cdot N_{\Omega_2} + \mathcal{W}_2) \\ &= \mathcal{L}_1 V_2 + \mathcal{L}_2 V_1 - V_1 V_2 \end{aligned}$$

# Cantor Lawn Volume



# Dimension of Fractal Lawn I

## Definition 5

The **Dimension**  $D_{\Omega^2} = \inf\{\alpha \geq 0 \mid V(\epsilon) = O(\epsilon^{2-\alpha}) \text{ as } \epsilon \rightarrow 0^+\}$

# Dimension of Fractal Lawn I

## Definition 5

The **Dimension**  $D_{\Omega^2} = \inf\{\alpha \geq 0 \mid V(\epsilon) = O(\epsilon^{2-\alpha}) \text{ as } \epsilon \rightarrow 0^+\}$

- Since  $\Omega^2 \subset \mathbb{R}^2$ ,  $V$  is required to remain bounded by  $\epsilon^{2-\alpha}$  as  $\epsilon \rightarrow 0^+$  rather than the 1D case

# Dimension of Fractal Lawn I

## Definition 5

The **Dimension**  $D_{\Omega^2} = \inf\{\alpha \geq 0 \mid V(\epsilon) = O(\epsilon^{2-\alpha}) \text{ as } \epsilon \rightarrow 0^+\}$

- Since  $\Omega^2 \subset \mathbb{R}^2$ ,  $V$  is required to remain bounded by  $\epsilon^{2-\alpha}$  as  $\epsilon \rightarrow 0^+$  rather than the 1D case

$$V(\epsilon) = \mathcal{L}_1 V_2 + \mathcal{L}_2 V_1 - V_1 V_2$$

$$\lim_{\epsilon \rightarrow 0^+} V(\epsilon) = O(\epsilon^{1-D_{\Omega_2}}) + O(\epsilon^{1-D_{\Omega_1}}) + O(\epsilon^{2-(D_{\Omega_1}+D_{\Omega_2})})$$

$$\lim_{\epsilon \rightarrow 0^+} \frac{V(\epsilon)}{\epsilon^{2-\alpha}} = O(\epsilon^{1-D_{\Omega_2}} \epsilon^{2-\alpha}) + O(\epsilon^{1-D_{\Omega_1}} \epsilon^{2-\alpha}) + O(\epsilon^{2-(D_{\Omega_1}+D_{\Omega_2})} \epsilon^{2-\alpha})$$



# Dimension of Fractal Lawn II

$$\lim_{\epsilon \rightarrow 0^+} \frac{V(\epsilon)}{\epsilon^{2-\alpha}} = O(\epsilon^{1-\alpha(1+D_{\Omega_2})}) + O(\epsilon^{1-\alpha(1+D_{\Omega_1})}) + O(\epsilon^{\alpha-(D_{\Omega_1}+D_{\Omega_2})})$$

# Dimension of Fractal Lawn II

$$\lim_{\epsilon \rightarrow 0^+} \frac{V(\epsilon)}{\epsilon^{2-\alpha}} = O(\epsilon^{1-\alpha(1+D_{\Omega_2})}) + O(\epsilon^{1-\alpha(1+D_{\Omega_1})}) + O(\epsilon^{\alpha-(D_{\Omega_1}+D_{\Omega_2})})$$

$$\begin{cases} \alpha - (1 + D_{\Omega_1}) \geq 0 \\ \alpha - (1 + D_{\Omega_2}) \geq 0 \\ \alpha - (D_{\Omega_1} + D_{\Omega_2}) \geq 0 \end{cases}$$

# Dimension of Fractal Lawn II

$$\lim_{\epsilon \rightarrow 0^+} \frac{V(\epsilon)}{\epsilon^{2-\alpha}} = O(\epsilon^{1-\alpha(1+D_{\Omega_2})}) + O(\epsilon^{1-\alpha(1+D_{\Omega_1})}) + O(\epsilon^{\alpha-(D_{\Omega_1}+D_{\Omega_2})})$$

$$\begin{cases} \alpha - (1 + D_{\Omega_1}) \geq 0 \\ \alpha - (1 + D_{\Omega_2}) \geq 0 \\ \alpha - (D_{\Omega_1} + D_{\Omega_2}) \geq 0 \end{cases}$$

$$\begin{cases} \alpha \geq 1 + D_{\Omega_1} \\ \alpha \geq 1 + D_{\Omega_2} \end{cases}$$

# Dimension of Fractal Lawn II

$$\lim_{\epsilon \rightarrow 0^+} \frac{V(\epsilon)}{\epsilon^{2-\alpha}} = O(\epsilon^{1-\alpha(1+D_{\Omega_2})}) + O(\epsilon^{1-\alpha(1+D_{\Omega_1})}) + O(\epsilon^{\alpha-(D_{\Omega_1}+D_{\Omega_2})})$$

$$\begin{cases} \alpha - (1 + D_{\Omega_1}) \geq 0 \\ \alpha - (1 + D_{\Omega_2}) \geq 0 \\ \alpha - (D_{\Omega_1} + D_{\Omega_2}) \geq 0 \end{cases}$$

$$\begin{cases} \alpha \geq 1 + D_{\Omega_1} \\ \alpha \geq 1 + D_{\Omega_2} \end{cases}$$

$$D_{\Omega^2} = \max\{(1 + D_{\Omega_1}), (1 + D_{\Omega_2})\}$$



# Dimension of the Cantor Lawn I

What is the dimension of the Cantor Lawn?

# Dimension of the Cantor Lawn I

What is the dimension of the Cantor Lawn?

- Apply the volume formula for a **fractal lawn** to the Cartesian product of 2 Cantor Strings.

# Dimension of the Cantor Lawn I

What is the dimension of the Cantor Lawn?

- Apply the volume formula for a **fractal lawn** to the Cartesian product of 2 Cantor Strings.
  - $D_\Omega = \log_3 2$



# Dimension of the Cantor Lawn I

What is the dimension of the Cantor Lawn?

- Apply the volume formula for a **fractal lawn** to the Cartesian product of 2 Cantor Strings.
  - $D_\Omega = \log_3 2$
  - $\Omega^2 = \Omega \times \Omega$

# Dimension of the Cantor Lawn I

What is the dimension of the Cantor Lawn?

- Apply the volume formula for a **fractal lawn** to the Cartesian product of 2 Cantor Strings.
  - $D_\Omega = \log_3 2$
  - $\Omega^2 = \Omega \times \Omega$
  - $\Omega_1 = \Omega_2$



# Dimension of the Cantor Lawn I

What is the dimension of the Cantor Lawn?

- Apply the volume formula for a **fractal lawn** to the Cartesian product of 2 Cantor Strings.
  - $D_\Omega = \log_3 2$
  - $\Omega^2 = \Omega \times \Omega$ 
    - $\Omega_1 = \Omega_2$
  - $D_{\Omega^2} = \max\{(1 + D_\Omega), 2 \cdot D_\Omega\}$

# Dimension of the Cantor Lawn I

What is the dimension of the Cantor Lawn?

- Apply the volume formula for a **fractal lawn** to the Cartesian product of 2 Cantor Strings.
  - $D_\Omega = \log_3 2$
  - $\Omega^2 = \Omega \times \Omega$ 
    - $\Omega_1 = \Omega_2$
  - $D_{\Omega^2} = \max\{(1 + D_\Omega), 2 \cdot D_\Omega\}$ 
    - $D_{\Omega^2} = 1 + D_\Omega$

# Dimension of the Cantor Lawn I

What is the dimension of the Cantor Lawn?

- Apply the volume formula for a **fractal lawn** to the Cartesian product of 2 Cantor Strings.
  - $D_\Omega = \log_3 2$
  - $\Omega^2 = \Omega \times \Omega$
  - $\Omega_1 = \Omega_2$
- $D_{\Omega^2} = \max\{(1 + D_\Omega), 2 \cdot D_\Omega\}$
- $D_{\Omega^2} = 1 + D_\Omega$

$$\begin{aligned}V(\epsilon)_{\Omega^2} &= \mathcal{L} V(\epsilon)_\Omega + \mathcal{L} V(\epsilon)_\Omega - V(\epsilon)_\Omega \cdot V(\epsilon)_\Omega \\&= 2\mathcal{L} V(\epsilon)_\Omega - (V(\epsilon)_\Omega)^2\end{aligned}$$



# Dimension of the Cantor Lawn II

Let  $t = \{-\log_3(2\epsilon)\}$  be defined as the fractional portion of  $-\log_3(2\epsilon)$

$$\begin{aligned}V_{\Omega^2}(\epsilon) &= (2\epsilon)^{1-\log_3 2} \left( 2 \left(\frac{1}{2}\right)^t + 2 \left(\frac{3}{2}\right)^t \right) + (2\epsilon)^{2-\log_3 2} \left( 2 \left(\frac{3}{2}\right)^t + 2 \left(\frac{1}{2}\right)^t \right) \\&= -(2\epsilon)^{2-\log_3 4} \left( 2 \left(\frac{3}{4}\right)^t + \left(\frac{1}{4}\right)^t + \left(\frac{9}{4}\right)^t \right) - (4\epsilon + 4\epsilon^2)\end{aligned}$$



## Dimension of the Cantor Lawn II

Let  $t = \{-\log_3(2\epsilon)\}$  be defined as the fractional portion of  $-\log_3(2\epsilon)$

$$\begin{aligned}V_{\Omega^2}(\epsilon) &= (2\epsilon)^{1-\log_3 2} \left( 2 \left(\frac{1}{2}\right)^t + 2 \left(\frac{3}{2}\right)^t \right) + (2\epsilon)^{2-\log_3 2} \left( 2 \left(\frac{3}{2}\right)^t + 2 \left(\frac{1}{2}\right)^t \right) \\&= -(2\epsilon)^{2-\log_3 4} \left( 2 \left(\frac{3}{4}\right)^t + \left(\frac{1}{4}\right)^t + \left(\frac{9}{4}\right)^t \right) - (4\epsilon + 4\epsilon^2)\end{aligned}$$

$$\begin{cases} -\log_3 2 + 1 - (2 - \alpha) \geq 0 \\ -\log_3 2 + 2 - (2 - \alpha) \geq 0 \\ -\log_3 4 + 2 - (2 - \alpha) \geq 0 \end{cases}$$



# Dimension of the Cantor Lawn III

$$\begin{cases} \alpha \geq 1 + \log_3 2 \\ \alpha \geq 1 + \log_3 4 \end{cases}$$

# Dimension of the Cantor Lawn III

$$\begin{cases} \alpha \geq 1 + \log_3 2 \\ \alpha \geq 1 + \log_3 4 \end{cases}$$

$$D_{\Omega^2} = \max\{(1 + \log_3 2), \log_3 4\}$$



# Dimension of the Cantor Lawn III


$$\begin{cases} \alpha \geq 1 + \log_3 2 \\ \alpha \geq 1 + \log_3 4 \end{cases}$$

$$D_{\Omega^2} = \max\{(1 + \log_3 2), \log_3 4\}$$


$$D_{\Omega^2} = 1 + \log_3 2$$




# Dimension of the Cantor Lawn IV



# Dimension of the Cantor Lawn V



# 3 Dimensional Fractal Objects



$$\begin{aligned} V(\epsilon) = & 2\epsilon \cdot N_{\Omega_1}(2\epsilon) \cdot \left( \sum_{j=1}^{N_{\Omega_2}(2\epsilon)} p_j \right) \cdot \left( \sum_{j=1}^{N_{\Omega_3}(2\epsilon)} q_j \right) + \\ & 2\epsilon \cdot N_{\Omega_2}(2\epsilon) \cdot \left( \sum_{j=1}^{N_{\Omega_1}(2\epsilon)} \ell_j - 2\epsilon \right) \cdot \left( \sum_{j=1}^{N_{\Omega_3}(2\epsilon)} q_j \right) + \\ & 2\epsilon \cdot N_{\Omega_3}(2\epsilon) \cdot \left( \sum_{j=1}^{N_{\Omega_2}(2\epsilon)} p_j - 2\epsilon \right) \cdot \left( \sum_{j=1}^{N_{\Omega_1}(2\epsilon)} \ell_j - 2\epsilon \right) + \\ & \mathcal{W}_1 \mathcal{L}_2 \mathcal{L}_3 + \mathcal{L}_1 \mathcal{W}_2 \mathcal{L}_3 + \mathcal{L}_1 \mathcal{L}_2 \mathcal{W}_3 - \mathcal{L}_1 \mathcal{W}_2 \mathcal{W}_3 - \mathcal{W}_1 \mathcal{L}_2 \mathcal{W}_3 - \\ & \mathcal{W}_1 \mathcal{W}_2 \mathcal{L}_3 + \mathcal{W}_1 \mathcal{W}_2 \mathcal{W}_3 \end{aligned}$$

# 3 Dimensional Fractal Volume

$$V(\epsilon) = 2\epsilon \cdot N_{\Omega_1} \cdot (\mathcal{L}_2 - \mathcal{W}_3) \cdot (\mathcal{L}_3 - \mathcal{W}_3) + 2\epsilon \cdot N_{\Omega_2} \cdot (\mathcal{L}_1 - \mathcal{W}_1 - 2\epsilon \cdot N_{\Omega_1}) \cdot (\mathcal{L}_3 - \mathcal{W}_3) + 2\epsilon \cdot N_{\Omega_3} \cdot (\mathcal{L}_2 - \mathcal{W}_2 - 2\epsilon \cdot N_{\Omega_2}) \cdot (\mathcal{L}_1 - \mathcal{W}_1 - 2\epsilon \cdot N_{\Omega_1}) + \mathcal{W}_1 \mathcal{L}_2 \mathcal{L}_3 + \mathcal{L}_1 \mathcal{W}_2 \mathcal{L}_3 + \mathcal{L}_1 \mathcal{L}_2 \mathcal{W}_3 - \mathcal{L}_1 \mathcal{W}_2 \mathcal{W}_3 - \mathcal{W}_1 \mathcal{L}_2 \mathcal{W}_3 - \mathcal{W}_1 \mathcal{W}_2 \mathcal{L}_3 + \mathcal{W}_1 \mathcal{W}_2 \mathcal{W}_3$$

# 3 Dimensional Fractal Volume

$$V(\epsilon) = 2\epsilon \cdot N_{\Omega_1} \cdot (\mathcal{L}_2 - \mathcal{W}_3) \cdot (\mathcal{L}_3 - \mathcal{W}_3) + 2\epsilon \cdot N_{\Omega_2} \cdot (\mathcal{L}_1 - \mathcal{W}_1 - 2\epsilon \cdot N_{\Omega_1}) \cdot (\mathcal{L}_3 - \mathcal{W}_3) + 2\epsilon \cdot N_{\Omega_3} \cdot (\mathcal{L}_2 - \mathcal{W}_2 - 2\epsilon \cdot N_{\Omega_2}) \cdot (\mathcal{L}_1 - \mathcal{W}_1 - 2\epsilon \cdot N_{\Omega_1}) + \mathcal{W}_1 \mathcal{L}_2 \mathcal{L}_3 + \mathcal{L}_1 \mathcal{W}_2 \mathcal{L}_3 + \mathcal{L}_1 \mathcal{L}_2 \mathcal{W}_3 - \mathcal{L}_1 \mathcal{W}_2 \mathcal{W}_3 - \mathcal{W}_1 \mathcal{L}_2 \mathcal{W}_3 - \mathcal{W}_1 \mathcal{W}_2 \mathcal{L}_3 + \mathcal{W}_1 \mathcal{W}_2 \mathcal{W}_3$$

$$V(\epsilon) = \mathcal{L}_1 \mathcal{L}_2 V_3(\epsilon) + \mathcal{L}_1 V_2(\epsilon) \mathcal{L}_3 + V_1(\epsilon) \mathcal{L}_2 \mathcal{L}_3 - V_1(\epsilon) V_2(\epsilon) \mathcal{L}_3 - V_1(\epsilon) \mathcal{L}_2 V_3(\epsilon) - \mathcal{L}_1 V_2(\epsilon) V_3(\epsilon) + V_1(\epsilon) V_2(\epsilon) V_3(\epsilon)$$

# 4 Dimensional Fractal Objects

$$\begin{aligned} V(\epsilon) = & 2\epsilon \cdot N_{\Omega_1}(2\epsilon) \cdot \left( \sum_{j=1}^{N_{\Omega_2}(2\epsilon)} p_j \right) \cdot \left( \sum_{j=1}^{N_{\Omega_3}(2\epsilon)} q_j \right) \cdot \left( \sum_{j=1}^{N_{\Omega_4}(2\epsilon)} z_j \right) + 2\epsilon \cdot \\ & N_{\Omega_2}(2\epsilon) \cdot \left( \sum_{j=1}^{N_{\Omega_1}(2\epsilon)} \ell_j - 2\epsilon \right) \cdot \left( \sum_{j=1}^{N_{\Omega_3}(2\epsilon)} q_j \right) \cdot \left( \sum_{j=1}^{N_{\Omega_4}(2\epsilon)} z_j \right) + 2\epsilon \cdot N_{\Omega_3}(2\epsilon) \cdot \\ & \left( \sum_{j=1}^{N_{\Omega_2}(2\epsilon)} p_j - 2\epsilon \right) \cdot \left( \sum_{j=1}^{N_{\Omega_1}(2\epsilon)} \ell_j - 2\epsilon \right) \cdot \left( \sum_{j=1}^{N_{\Omega_4}(2\epsilon)} z_j \right) + 2\epsilon \cdot N_{\Omega_4}(2\epsilon) \cdot \\ & \left( \sum_{j=1}^{N_{\Omega_2}(2\epsilon)} p_j - 2\epsilon \right) \cdot \left( \sum_{j=1}^{N_{\Omega_1}(2\epsilon)} \ell_j - 2\epsilon \right) \cdot \left( \sum_{j=1}^{N_{\Omega_3}(2\epsilon)} q_j - 2\epsilon \right) + W_1 L_2 L_3 L_4 + \\ & L_1 W_2 L_3 L_4 + L_1 L_2 W_3 L_4 + L_1 L_2 L_3 W_4 - L_1 L_2 W_3 W_4 - L_1 W_2 L_3 W_4 - \\ & L_1 W_2 W_3 L_4 - W_1 L_2 L_3 W_4 - W_1 L_2 W_3 L_4 - W_1 W_2 L_3 L_4 + L_1 W_2 W_3 W_4 + \\ & W_1 L_2 W_3 W_4 + W_1 W_2 L_3 W_4 + W_1 W_2 W_3 L_4 - W_1 W_2 W_3 W_4 \end{aligned}$$

# 4 Dimensional Volume Formula

$$V^4(\epsilon) = \mathcal{L}_1 \mathcal{L}_2 \mathcal{L}_3 V_4(\epsilon) + \mathcal{L}_1 \mathcal{L}_2 V_3(\epsilon) \mathcal{L}_4 + \mathcal{L}_1 V_2(\epsilon) \mathcal{L}_3 \mathcal{L}_4 + V_1(\epsilon) \mathcal{L}_2 \mathcal{L}_3 \mathcal{L}_4 - V_1(\epsilon) V_2(\epsilon) \mathcal{L}_3 \mathcal{L}_4 - V_1(\epsilon) \mathcal{L}_2 V_3(\epsilon) \mathcal{L}_4 - V_1(\epsilon) \mathcal{L}_2 \mathcal{L}_3 V_4(\epsilon) - \mathcal{L}_1 V_2(\epsilon) V_3(\epsilon) \mathcal{L}_4 - \mathcal{L}_1 V_2(\epsilon) \mathcal{L}_3 V_4(\epsilon) - \mathcal{L}_1 \mathcal{L}_2 V_3(\epsilon) V_4(\epsilon) + V_1(\epsilon) V_2(\epsilon) V_3(\epsilon) \mathcal{L}_4 + V_1(\epsilon) V_2(\epsilon) \mathcal{L}_3 V_4(\epsilon) + V_1(\epsilon) \mathcal{L}_2 V_3(\epsilon) V_4(\epsilon) + \mathcal{L}_1 V_2(\epsilon) V_3(\epsilon) V_4(\epsilon) - V_1(\epsilon) V_2(\epsilon) V_3(\epsilon) V_4(\epsilon)$$



# 4 Dimensional Volume Formula

$$V^4(\epsilon) = \mathcal{L}_1 \mathcal{L}_2 \mathcal{L}_3 V_4(\epsilon) + \mathcal{L}_1 \mathcal{L}_2 V_3(\epsilon) \mathcal{L}_4 + \mathcal{L}_1 V_2(\epsilon) \mathcal{L}_3 \mathcal{L}_4 + V_1(\epsilon) \mathcal{L}_2 \mathcal{L}_3 \mathcal{L}_4 - V_1(\epsilon) V_2(\epsilon) \mathcal{L}_3 \mathcal{L}_4 - V_1(\epsilon) \mathcal{L}_2 V_3(\epsilon) \mathcal{L}_4 - V_1(\epsilon) \mathcal{L}_2 \mathcal{L}_3 V_4(\epsilon) - \mathcal{L}_1 V_2(\epsilon) V_3(\epsilon) \mathcal{L}_4 - \mathcal{L}_1 \mathcal{L}_2 V_3(\epsilon) V_4(\epsilon) + V_1(\epsilon) V_2(\epsilon) V_3(\epsilon) \mathcal{L}_4 + V_1(\epsilon) V_2(\epsilon) \mathcal{L}_3 V_4(\epsilon) + V_1(\epsilon) \mathcal{L}_2 V_3(\epsilon) V_4(\epsilon) + \mathcal{L}_1 V_2(\epsilon) V_3(\epsilon) V_4(\epsilon) - V_1(\epsilon) V_2(\epsilon) V_3(\epsilon) V_4(\epsilon)$$

$$V^3(\epsilon) = \mathcal{L}_1 \mathcal{L}_2 V_3(\epsilon) + \mathcal{L}_1 V_2(\epsilon) \mathcal{L}_3 + V_1(\epsilon) \mathcal{L}_2 \mathcal{L}_3 - V_1(\epsilon) V_2(\epsilon) \mathcal{L}_3 - V_1(\epsilon) \mathcal{L}_2 V_3(\epsilon) - \mathcal{L}_1 V_2(\epsilon) V_3(\epsilon) + V_1(\epsilon) V_2(\epsilon) V_3(\epsilon)$$

$$V^2(\epsilon) = \mathcal{L}_1 V_2(\epsilon) + V_1(\epsilon) \mathcal{L}_2 - V_1(\epsilon) V_2(\epsilon)$$

# Algebraic Manipulations

$$\begin{aligned}V^2(\epsilon) &= \mathcal{L}_1 V_2(\epsilon) + V_1(\epsilon) \mathcal{L}_2 - V_1(\epsilon) V_2(\epsilon) \\&= (-1) \cdot (V_1(\epsilon) - \mathcal{L}_1) \cdot (V_2(\epsilon) - \mathcal{L}_2) + \mathcal{L}_1 \mathcal{L}_2\end{aligned}$$

# Algebraic Manipulations

$$\begin{aligned}V^2(\epsilon) &= \mathcal{L}_1 V_2(\epsilon) + V_1(\epsilon) \mathcal{L}_2 - V_1(\epsilon) V_2(\epsilon) \\&= (-1) \cdot (V_1(\epsilon) - \mathcal{L}_1) \cdot (V_2(\epsilon) - \mathcal{L}_2) + \mathcal{L}_1 \mathcal{L}_2\end{aligned}$$

$$\begin{aligned}V^3(\epsilon) &= \mathcal{L}_1 \mathcal{L}_2 V_3(\epsilon) + \mathcal{L}_1 V_2(\epsilon) \mathcal{L}_3 + V_1(\epsilon) \mathcal{L}_2 \mathcal{L}_3 - V_1(\epsilon) V_2(\epsilon) \mathcal{L}_3 - V_1(\epsilon) \mathcal{L}_2 V_3(\epsilon) - \\&\quad \mathcal{L}_1 V_2(\epsilon) V_3(\epsilon) + V_1(\epsilon) V_2(\epsilon) V_3(\epsilon) \\&= (-1)^2 \cdot (V_1(\epsilon) - \mathcal{L}_1) \cdot (V_2(\epsilon) - \mathcal{L}_2) \cdot (V_3(\epsilon) - \mathcal{L}_3) + \mathcal{L}_1 \mathcal{L}_2 \mathcal{L}_3\end{aligned}$$

# Algebraic Manipulations

$$\begin{aligned} V^4(\epsilon) &= \mathcal{L}_1 \mathcal{L}_2 \mathcal{L}_3 V_4(\epsilon) + \mathcal{L}_1 \mathcal{L}_2 V_3(\epsilon) \mathcal{L}_4 + \mathcal{L}_1 V_2(\epsilon) \mathcal{L}_3 \mathcal{L}_4 + V_1(\epsilon) \mathcal{L}_2 \mathcal{L}_3 \mathcal{L}_4 - \\ &\quad V_1(\epsilon) V_2(\epsilon) \mathcal{L}_3 \mathcal{L}_4 - V_1(\epsilon) \mathcal{L}_2 V_3(\epsilon) \mathcal{L}_4 - V_1(\epsilon) \mathcal{L}_2 \mathcal{L}_3 V_4(\epsilon) - \mathcal{L}_1 V_2(\epsilon) V_3(\epsilon) \mathcal{L}_4 - \\ &\quad \mathcal{L}_1 V_2(\epsilon) \mathcal{L}_3 V_4(\epsilon) - \mathcal{L}_1 \mathcal{L}_2 V_3(\epsilon) V_4(\epsilon) + V_1(\epsilon) V_2(\epsilon) V_3(\epsilon) \mathcal{L}_4 + V_1(\epsilon) V_2(\epsilon) \mathcal{L}_3 V_4(\epsilon) + \\ &\quad V_1(\epsilon) \mathcal{L}_2 V_3(\epsilon) V_4(\epsilon) + \mathcal{L}_1 V_2(\epsilon) V_3(\epsilon) V_4(\epsilon) - V_1(\epsilon) V_2(\epsilon) V_3(\epsilon) V_4(\epsilon) \\ &= (-1)^3 \cdot (V_1(\epsilon) - \mathcal{L}_1) \cdot (V_2(\epsilon) - \mathcal{L}_2) \cdot (V_3(\epsilon) - \mathcal{L}_3) \cdot (V_4(\epsilon) - \mathcal{L}_4) + \mathcal{L}_1 \mathcal{L}_2 \mathcal{L}_3 \mathcal{L}_4 \end{aligned}$$

# General Volume Formula

$$V^2(\epsilon) = (-1) \cdot (V_1(\epsilon) - \mathcal{L}_1) \cdot (V_2(\epsilon) - \mathcal{L}_2) + \mathcal{L}_1 \mathcal{L}_2$$

$$V^3(\epsilon) = (-1)^2 \cdot (V_1(\epsilon) - \mathcal{L}_1) \cdot (V_2(\epsilon) - \mathcal{L}_2) \cdot (V_3(\epsilon) - \mathcal{L}_3) + \mathcal{L}_1 \mathcal{L}_2 \mathcal{L}_3$$

$$V^4(\epsilon) = (-1)^3 \cdot (V_1(\epsilon) - \mathcal{L}_1) \cdot (V_2(\epsilon) - \mathcal{L}_2) \cdot (V_3(\epsilon) - \mathcal{L}_3) \cdot (V_4(\epsilon) - \mathcal{L}_4) + \mathcal{L}_1 \mathcal{L}_2 \mathcal{L}_3 \mathcal{L}_4$$

$$V^n(\epsilon) = \left[ (-1)^{n-1} \cdot \prod_{k=1}^n (V_k(\epsilon) - \mathcal{L}_k) \right] + \prod_{k=1}^n \mathcal{L}_k$$



# Future Work

- Prove the general volume formula
- Find a general formula for dimension (based on general volume formula)

# References I

-  D. A. Bini and L. Robol. "Solving secular and polynomial equations: A multiprecision algorithm." *Journal of Computational and Applied Mathematics* 272 (2014), 276–292.
-  M. R. Bremner. *Lattice Basis Reduction: An Introduction to the LLL Algorithm and its Applications*, Taylor & Francis, Boca Raton, 2011.
-  M. L. Lapidus. "Fractal drum, inverse spectral problems for elliptic operators and a partial resolution of the Weyl-Berry conjecture." *Transactions of the American Mathematical Society*. (2) 325 (1991), 465–529.

## References II

-  M. L. Lapidus. "Vibrations of Fractal Drums, the Riemann Hypothesis, Waves in Fractal Media and the Weyl-Berry Conjecture." In *Ordinary and Partial Differential Equations*, edited by B. D. Sleeman and R. J. Jarvis, Vol. IV, Proc. Twelfth Intern. Conf. (Dundee, Scotland, UK, June 1992), Pitman Research Notes in Math. Series 289, pp. 126–209, London: Longman Scientific and Technical, 1993.
-  M. L. Lapidus. *In Search of the Riemann Zeros: Strings, fractal membranes and noncommutative spacetimes*, American Mathematical Society, Providence, Rhode Island, 2008.
-  M. L. Lapidus. "An overview of the complex fractal dimensions: From fractal strings to fractal drums, and back," *Contemporary Mathematics*, Vol. 731, Amer. Math. Soc., Providence, R.I., 2019, 143–269

## References III

-  M. L. Lapidus and C. Pomerance. "The Riemann Zeta-Function and the One-Dimensional Weyl-Berry Conjecture for Fractal Drums," *Proc. London Math. Soc.* (3) 66 (1993), 41–69.
-  M. L. Lapidus and H. Maier. "The Riemann Hypothesis and Inverse Spectral Problems for Fractal Strings." *J. London Math. Soc.* (2) 52 (1995), 15–34.
-  M. L. Lapidus and M. van Frankenhuysen. *Fractal Geometry and Number Theory (Complex dimensions of fractal strings and zeros of zeta functions)*, Birkhäuser, Boston, 2000.
-  M. L. Lapidus and M. van Frankenhuysen. "Complex Dimensions of Self-Similar Fractal Strings and Diophantine Approximation," *J. Experimental Math.* (1) 12 (2003), 41–69.



## References IV

-  M. L. Lapidus and M. van Frankenhuysen. *Fractal Geometry, Complex Dimensions and Zeta Functions: Geometry and Spectra of Fractal Strings*, second edition (of the 2006 edition), Springer Monographs in Mathematics, Springer, New York, 2012.
-  M. L. Lapidus and G. Radunović and D. Žubrinić. *Fractal Zeta Functions and Fractal Drums: Higher-Dimensional Theory of Complex Dimensions*, Springer Monographs in Mathematics, Springer, New York, 2017.
-  A. K. Lenstra and H. W. Lenstra and L. Lovász. “Factoring polynomials with rational coefficients,” *Mathematische Annalen* (4) 261 (1982), 515–534.
-  W. M. Schmidt. *Diophantine Approximation*, Springer, New York, 1980.

# References V

-  J.-P. Serre. *A Course in Arithmetic*, English translation, Springer-Verlag, Berlin, 1973.
-  E. K. Voskanian. “On the Quasiperiodic Structure of the Complex Dimensions of Self-Similar Fractal Strings,” Ph. D. Dissertation, University of California, Riverside, 2019.