

Badass Title

Isaac Ashton, Joseph Ingenito, Peter Tonuzi

The College of New Jersey



What is a Fractal String

Definition 1

A **fractal string** Ω is a bounded open subset of the real line.

What is a Fractal String

Definition 1

A **fractal string** Ω is a bounded open subset of the real line.

$$\Omega = \bigcup_{j=1}^{\infty} \ell_j, \quad \mathcal{L} = \text{Vol}_1(\Omega) = \sum_{j=1}^{\infty} \ell_j \implies \lim_{j \rightarrow \infty} \ell_j = 0.$$



What is a Fractal String

Definition 1

A **fractal string** Ω is a bounded open subset of the real line.

$$\Omega = \bigcup_{j=1}^{\infty} \ell_j, \quad \mathcal{L} = \text{Vol}_1(\Omega) = \sum_{j=1}^{\infty} \ell_j \implies \lim_{j \rightarrow \infty} \ell_j = 0.$$

$$\ell_1 \geq \ell_2 \geq \ell_3 \geq \cdots \geq 0$$

What is a Fractal String

Definition 1

A **fractal string** Ω is a bounded open subset of the real line.

$$\Omega = \bigcup_{j=1}^{\infty} \ell_j, \quad \mathcal{L} = \text{Vol}_1(\Omega) = \sum_{j=1}^{\infty} \ell_j \implies \lim_{j \rightarrow \infty} \ell_j = 0.$$

$$\ell_1 \geq \ell_2 \geq \ell_3 \geq \cdots \geq 0$$

The fractal string we studied the most in particular is the **Cantor String**



Cantor String

Inner-Tubular Neighborhood

Definition 2

Inner-tubular neighborhood: $V(\epsilon) = \text{vol}_1\{x \in \Omega \mid d(x, \partial\Omega) < \epsilon\}$

Inner-Tubular Neighborhood

Definition 2

Inner-tubular neighborhood: $V(\epsilon) = \text{vol}_1\{x \in \Omega \mid d(x, \partial\Omega) < \epsilon\}$

Counting function: $N_\Omega(x) = \sum_{\ell_j^{-1} \leq x} 1$

More Notation: $\mathcal{W}_\Omega(x) = \sum_{j: \ell_j^{-1} > x} \ell_j$

Now we can write: $V(\epsilon) = 2\epsilon \cdot N_\Omega(2\epsilon) + \mathcal{W}_\Omega(2\epsilon)$



Cantor String Volume



Inner Minkowski Dimension of a Fractal String

Definition 3

Dimension of Ω : $D_\Omega = \inf \{ \alpha \geq 0 \mid V(\epsilon) = O(\epsilon^{1-\alpha}) \text{ as } \epsilon \rightarrow 0^+ \}$

Inner Minkowski Dimension of a Fractal String

Definition 3

Dimension of Ω : $D_\Omega = \inf \{ \alpha \geq 0 \mid V(\epsilon) = O(\epsilon^{1-\alpha}) \text{ as } \epsilon \rightarrow 0^+ \}$

In other terms: $D_\Omega = \text{the smallest } \alpha \text{ st } \lim_{\epsilon \rightarrow 0^+} \frac{V(\epsilon)}{\epsilon^{1-\alpha}} = c, \ c \in \mathbb{R}$

Inner Minkowski Dimension of a Fractal String


Definition 3

Dimension of Ω : $D_\Omega = \inf \{ \alpha \geq 0 \mid V(\epsilon) = O(\epsilon^{1-\alpha}) \text{ as } \epsilon \rightarrow 0^+ \}$

In other terms: $D_\Omega = \text{the smallest } \alpha \text{ st } \lim_{\epsilon \rightarrow 0^+} \frac{V(\epsilon)}{\epsilon^{1-\alpha}} = c, \ c \in \mathbb{R}$

Cantor String Dimension: $D_{cs} = \log_3 2$

3 Dimensional Fractal Objects



$$\begin{aligned} V(\epsilon) = & 2\epsilon \cdot N_{\Omega_1}(2\epsilon) \cdot \left(\sum_{j=1}^{N_{\Omega_2}(2\epsilon)} p_j \right) \cdot \left(\sum_{j=1}^{N_{\Omega_3}(2\epsilon)} q_j \right) + \\ & 2\epsilon \cdot N_{\Omega_2}(2\epsilon) \cdot \left(\sum_{j=1}^{N_{\Omega_1}(2\epsilon)} \ell_j - 2\epsilon \right) \cdot \left(\sum_{j=1}^{N_{\Omega_3}(2\epsilon)} q_j \right) + \\ & 2\epsilon \cdot N_{\Omega_3}(2\epsilon) \cdot \left(\sum_{j=1}^{N_{\Omega_2}(2\epsilon)} p_j - 2\epsilon \right) \cdot \left(\sum_{j=1}^{N_{\Omega_1}(2\epsilon)} \ell_j - 2\epsilon \right) + \\ & \mathcal{W}_1 \mathcal{L}_2 \mathcal{L}_3 + \mathcal{L}_1 \mathcal{W}_2 \mathcal{L}_3 + \mathcal{L}_1 \mathcal{L}_2 \mathcal{W}_3 - \mathcal{L}_1 \mathcal{W}_2 \mathcal{W}_3 - \mathcal{W}_1 \mathcal{L}_2 \mathcal{W}_3 - \\ & \mathcal{W}_1 \mathcal{W}_2 \mathcal{L}_3 + \mathcal{W}_1 \mathcal{W}_2 \mathcal{W}_3 \end{aligned}$$

3 Dimensional Fractal Volume

$$V(\epsilon) = 2\epsilon \cdot N_{\Omega_1} \cdot (\mathcal{L}_2 - \mathcal{W}_3) \cdot (\mathcal{L}_3 - \mathcal{W}_3) + 2\epsilon \cdot N_{\Omega_2} \cdot (\mathcal{L}_1 - \mathcal{W}_1 - 2\epsilon \cdot N_{\Omega_1}) \cdot (\mathcal{L}_3 - \mathcal{W}_3) + 2\epsilon \cdot N_{\Omega_3} \cdot (\mathcal{L}_2 - \mathcal{W}_2 - 2\epsilon \cdot N_{\Omega_2}) \cdot (\mathcal{L}_1 - \mathcal{W}_1 - 2\epsilon \cdot N_{\Omega_1}) + \mathcal{W}_1 \mathcal{L}_2 \mathcal{L}_3 + \mathcal{L}_1 \mathcal{W}_2 \mathcal{L}_3 + \mathcal{L}_1 \mathcal{L}_2 \mathcal{W}_3 - \mathcal{L}_1 \mathcal{W}_2 \mathcal{W}_3 - \mathcal{W}_1 \mathcal{L}_2 \mathcal{W}_3 - \mathcal{W}_1 \mathcal{W}_2 \mathcal{L}_3 + \mathcal{W}_1 \mathcal{W}_2 \mathcal{W}_3$$

3 Dimensional Fractal Volume

$$V(\epsilon) = 2\epsilon \cdot N_{\Omega_1} \cdot (\mathcal{L}_2 - \mathcal{W}_3) \cdot (\mathcal{L}_3 - \mathcal{W}_3) + 2\epsilon \cdot N_{\Omega_2} \cdot (\mathcal{L}_1 - \mathcal{W}_1 - 2\epsilon \cdot N_{\Omega_1}) \cdot (\mathcal{L}_3 - \mathcal{W}_3) + 2\epsilon \cdot N_{\Omega_3} \cdot (\mathcal{L}_2 - \mathcal{W}_2 - 2\epsilon \cdot N_{\Omega_2}) \cdot (\mathcal{L}_1 - \mathcal{W}_1 - 2\epsilon \cdot N_{\Omega_1}) + \mathcal{W}_1 \mathcal{L}_2 \mathcal{L}_3 + \mathcal{L}_1 \mathcal{W}_2 \mathcal{L}_3 + \mathcal{L}_1 \mathcal{L}_2 \mathcal{W}_3 - \mathcal{L}_1 \mathcal{W}_2 \mathcal{W}_3 - \mathcal{W}_1 \mathcal{L}_2 \mathcal{W}_3 - \mathcal{W}_1 \mathcal{W}_2 \mathcal{L}_3 + \mathcal{W}_1 \mathcal{W}_2 \mathcal{W}_3$$

$$V(\epsilon) = \mathcal{L}_1 \mathcal{L}_2 V_3(\epsilon) + \mathcal{L}_1 V_2(\epsilon) \mathcal{L}_3 + V_1(\epsilon) \mathcal{L}_2 \mathcal{L}_3 - V_1(\epsilon) V_2(\epsilon) \mathcal{L}_3 - V_1(\epsilon) \mathcal{L}_2 V_3(\epsilon) - \mathcal{L}_1 V_2(\epsilon) V_3(\epsilon) + V_1(\epsilon) V_2(\epsilon) V_3(\epsilon)$$

4 Dimensional Fractal Objects

$$\begin{aligned} V(\epsilon) = & 2\epsilon \cdot N_{\Omega_1}(2\epsilon) \cdot \left(\sum_{j=1}^{N_{\Omega_2}(2\epsilon)} p_j \right) \cdot \left(\sum_{j=1}^{N_{\Omega_3}(2\epsilon)} q_j \right) \cdot \left(\sum_{j=1}^{N_{\Omega_4}(2\epsilon)} z_j \right) + 2\epsilon \cdot \\ & N_{\Omega_2}(2\epsilon) \cdot \left(\sum_{j=1}^{N_{\Omega_1}(2\epsilon)} \ell_j - 2\epsilon \right) \cdot \left(\sum_{j=1}^{N_{\Omega_3}(2\epsilon)} q_j \right) \cdot \left(\sum_{j=1}^{N_{\Omega_4}(2\epsilon)} z_j \right) + 2\epsilon \cdot N_{\Omega_3}(2\epsilon) \cdot \\ & \left(\sum_{j=1}^{N_{\Omega_2}(2\epsilon)} p_j - 2\epsilon \right) \cdot \left(\sum_{j=1}^{N_{\Omega_1}(2\epsilon)} \ell_j - 2\epsilon \right) \cdot \left(\sum_{j=1}^{N_{\Omega_4}(2\epsilon)} z_j \right) + 2\epsilon \cdot N_{\Omega_4}(2\epsilon) \cdot \\ & \left(\sum_{j=1}^{N_{\Omega_2}(2\epsilon)} p_j - 2\epsilon \right) \cdot \left(\sum_{j=1}^{N_{\Omega_1}(2\epsilon)} \ell_j - 2\epsilon \right) \cdot \left(\sum_{j=1}^{N_{\Omega_3}(2\epsilon)} q_j - 2\epsilon \right) + W_1 L_2 L_3 L_4 + \\ & L_1 W_2 L_3 L_4 + L_1 L_2 W_3 L_4 + L_1 L_2 L_3 W_4 - L_1 L_2 W_3 W_4 - L_1 W_2 L_3 W_4 - \\ & L_1 W_2 W_3 L_4 - W_1 L_2 L_3 W_4 - W_1 L_2 W_3 L_4 - W_1 W_2 L_3 L_4 + L_1 W_2 W_3 W_4 + \\ & W_1 L_2 W_3 W_4 + W_1 W_2 L_3 W_4 + W_1 W_2 W_3 L_4 - W_1 W_2 W_3 W_4 \end{aligned}$$

4 Dimensional Volume Formula

$$V^4(\epsilon) = \mathcal{L}_1 \mathcal{L}_2 \mathcal{L}_3 V_4(\epsilon) + \mathcal{L}_1 \mathcal{L}_2 V_3(\epsilon) \mathcal{L}_4 + \mathcal{L}_1 V_2(\epsilon) \mathcal{L}_3 \mathcal{L}_4 + V_1(\epsilon) \mathcal{L}_2 \mathcal{L}_3 \mathcal{L}_4 - V_1(\epsilon) V_2(\epsilon) \mathcal{L}_3 \mathcal{L}_4 - V_1(\epsilon) \mathcal{L}_2 V_3(\epsilon) \mathcal{L}_4 - V_1(\epsilon) \mathcal{L}_2 \mathcal{L}_3 V_4(\epsilon) - \mathcal{L}_1 V_2(\epsilon) V_3(\epsilon) \mathcal{L}_4 - \mathcal{L}_1 V_2(\epsilon) \mathcal{L}_3 V_4(\epsilon) - \mathcal{L}_1 \mathcal{L}_2 V_3(\epsilon) V_4(\epsilon) + V_1(\epsilon) V_2(\epsilon) V_3(\epsilon) \mathcal{L}_4 + V_1(\epsilon) V_2(\epsilon) \mathcal{L}_3 V_4(\epsilon) + V_1(\epsilon) \mathcal{L}_2 V_3(\epsilon) V_4(\epsilon) + \mathcal{L}_1 V_2(\epsilon) V_3(\epsilon) V_4(\epsilon) - V_1(\epsilon) V_2(\epsilon) V_3(\epsilon) V_4(\epsilon)$$



4 Dimensional Volume Formula

$$V^4(\epsilon) = \mathcal{L}_1 \mathcal{L}_2 \mathcal{L}_3 V_4(\epsilon) + \mathcal{L}_1 \mathcal{L}_2 V_3(\epsilon) \mathcal{L}_4 + \mathcal{L}_1 V_2(\epsilon) \mathcal{L}_3 \mathcal{L}_4 + V_1(\epsilon) \mathcal{L}_2 \mathcal{L}_3 \mathcal{L}_4 - V_1(\epsilon) V_2(\epsilon) \mathcal{L}_3 \mathcal{L}_4 - V_1(\epsilon) \mathcal{L}_2 V_3(\epsilon) \mathcal{L}_4 - V_1(\epsilon) \mathcal{L}_2 \mathcal{L}_3 V_4(\epsilon) - \mathcal{L}_1 V_2(\epsilon) V_3(\epsilon) \mathcal{L}_4 - \mathcal{L}_1 \mathcal{L}_2 V_3(\epsilon) V_4(\epsilon) + V_1(\epsilon) V_2(\epsilon) V_3(\epsilon) \mathcal{L}_4 + V_1(\epsilon) V_2(\epsilon) \mathcal{L}_3 V_4(\epsilon) + V_1(\epsilon) \mathcal{L}_2 V_3(\epsilon) V_4(\epsilon) + \mathcal{L}_1 V_2(\epsilon) V_3(\epsilon) V_4(\epsilon) - V_1(\epsilon) V_2(\epsilon) V_3(\epsilon) V_4(\epsilon)$$

$$V^3(\epsilon) = \mathcal{L}_1 \mathcal{L}_2 V_3(\epsilon) + \mathcal{L}_1 V_2(\epsilon) \mathcal{L}_3 + V_1(\epsilon) \mathcal{L}_2 \mathcal{L}_3 - V_1(\epsilon) V_2(\epsilon) \mathcal{L}_3 - V_1(\epsilon) \mathcal{L}_2 V_3(\epsilon) - \mathcal{L}_1 V_2(\epsilon) V_3(\epsilon) + V_1(\epsilon) V_2(\epsilon) V_3(\epsilon)$$

$$V^2(\epsilon) = \mathcal{L}_1 V_2(\epsilon) + V_1(\epsilon) \mathcal{L}_2 - V_1(\epsilon) V_2(\epsilon)$$

Algebraic Manipulations

$$\begin{aligned}V^2(\epsilon) &= \mathcal{L}_1 V_2(\epsilon) + V_1(\epsilon) \mathcal{L}_2 - V_1(\epsilon) V_2(\epsilon) \\&= (-1) \cdot (V_1(\epsilon) - \mathcal{L}_1) \cdot (V_2(\epsilon) - \mathcal{L}_2) + \mathcal{L}_1 \mathcal{L}_2\end{aligned}$$

Algebraic Manipulations

$$\begin{aligned}V^2(\epsilon) &= \mathcal{L}_1 V_2(\epsilon) + V_1(\epsilon) \mathcal{L}_2 - V_1(\epsilon) V_2(\epsilon) \\&= (-1) \cdot (V_1(\epsilon) - \mathcal{L}_1) \cdot (V_2(\epsilon) - \mathcal{L}_2) + \mathcal{L}_1 \mathcal{L}_2\end{aligned}$$

$$\begin{aligned}V^3(\epsilon) &= \mathcal{L}_1 \mathcal{L}_2 V_3(\epsilon) + \mathcal{L}_1 V_2(\epsilon) \mathcal{L}_3 + V_1(\epsilon) \mathcal{L}_2 \mathcal{L}_3 - V_1(\epsilon) V_2(\epsilon) \mathcal{L}_3 - V_1(\epsilon) \mathcal{L}_2 V_3(\epsilon) - \\&\quad \mathcal{L}_1 V_2(\epsilon) V_3(\epsilon) + V_1(\epsilon) V_2(\epsilon) V_3(\epsilon) \\&= (-1)^2 \cdot (V_1(\epsilon) - \mathcal{L}_1) \cdot (V_2(\epsilon) - \mathcal{L}_2) \cdot (V_3(\epsilon) - \mathcal{L}_3) + \mathcal{L}_1 \mathcal{L}_2 \mathcal{L}_3\end{aligned}$$

Algebraic Manipulations

$$\begin{aligned} V^4(\epsilon) &= \mathcal{L}_1 \mathcal{L}_2 \mathcal{L}_3 V_4(\epsilon) + \mathcal{L}_1 \mathcal{L}_2 V_3(\epsilon) \mathcal{L}_4 + \mathcal{L}_1 V_2(\epsilon) \mathcal{L}_3 \mathcal{L}_4 + V_1(\epsilon) \mathcal{L}_2 \mathcal{L}_3 \mathcal{L}_4 - \\ &\quad V_1(\epsilon) V_2(\epsilon) \mathcal{L}_3 \mathcal{L}_4 - V_1(\epsilon) \mathcal{L}_2 V_3(\epsilon) \mathcal{L}_4 - V_1(\epsilon) \mathcal{L}_2 \mathcal{L}_3 V_4(\epsilon) - \mathcal{L}_1 V_2(\epsilon) V_3(\epsilon) \mathcal{L}_4 - \\ &\quad \mathcal{L}_1 V_2(\epsilon) \mathcal{L}_3 V_4(\epsilon) - \mathcal{L}_1 \mathcal{L}_2 V_3(\epsilon) V_4(\epsilon) + V_1(\epsilon) V_2(\epsilon) V_3(\epsilon) \mathcal{L}_4 + V_1(\epsilon) V_2(\epsilon) \mathcal{L}_3 V_4(\epsilon) + \\ &\quad V_1(\epsilon) \mathcal{L}_2 V_3(\epsilon) V_4(\epsilon) + \mathcal{L}_1 V_2(\epsilon) V_3(\epsilon) V_4(\epsilon) - V_1(\epsilon) V_2(\epsilon) V_3(\epsilon) V_4(\epsilon) \\ &= (-1)^3 \cdot (V_1(\epsilon) - \mathcal{L}_1) \cdot (V_2(\epsilon) - \mathcal{L}_2) \cdot (V_3(\epsilon) - \mathcal{L}_3) \cdot (V_4(\epsilon) - \mathcal{L}_4) + \mathcal{L}_1 \mathcal{L}_2 \mathcal{L}_3 \mathcal{L}_4 \end{aligned}$$

General Volume Formula

$$V^2(\epsilon) = (-1) \cdot (V_1(\epsilon) - \mathcal{L}_1) \cdot (V_2(\epsilon) - \mathcal{L}_2) + \mathcal{L}_1 \mathcal{L}_2$$

$$V^3(\epsilon) = (-1)^2 \cdot (V_1(\epsilon) - \mathcal{L}_1) \cdot (V_2(\epsilon) - \mathcal{L}_2) \cdot (V_3(\epsilon) - \mathcal{L}_3) + \mathcal{L}_1 \mathcal{L}_2 \mathcal{L}_3$$

$$V^4(\epsilon) = (-1)^3 \cdot (V_1(\epsilon) - \mathcal{L}_1) \cdot (V_2(\epsilon) - \mathcal{L}_2) \cdot (V_3(\epsilon) - \mathcal{L}_3) \cdot (V_4(\epsilon) - \mathcal{L}_4) + \mathcal{L}_1 \mathcal{L}_2 \mathcal{L}_3 \mathcal{L}_4$$

$$V^n(\epsilon) = \left[(-1)^{n-1} \cdot \prod_{k=1}^n (V_k(\epsilon) - \mathcal{L}_k) \right] + \prod_{k=1}^n \mathcal{L}_k$$



Future Work

- Prove the general volume formula
- Find a general formula for dimension (based on general volume formula)

References I

-  D. A. Bini and L. Robol. "Solving secular and polynomial equations: A multiprecision algorithm." *Journal of Computational and Applied Mathematics* 272 (2014), 276–292.
-  M. R. Bremner. *Lattice Basis Reduction: An Introduction to the LLL Algorithm and its Applications*, Taylor & Francis, Boca Raton, 2011.
-  M. L. Lapidus. "Fractal drum, inverse spectral problems for elliptic operators and a partial resolution of the Weyl-Berry conjecture." *Transactions of the American Mathematical Society*. (2) 325 (1991), 465–529.

References II

-  M. L. Lapidus. "Vibrations of Fractal Drums, the Riemann Hypothesis, Waves in Fractal Media and the Weyl-Berry Conjecture." In *Ordinary and Partial Differential Equations*, edited by B. D. Sleeman and R. J. Jarvis, Vol. IV, Proc. Twelfth Intern. Conf. (Dundee, Scotland, UK, June 1992), Pitman Research Notes in Math. Series 289, pp. 126–209, London: Longman Scientific and Technical, 1993.
-  M. L. Lapidus. *In Search of the Riemann Zeros: Strings, fractal membranes and noncommutative spacetimes*, American Mathematical Society, Providence, Rhode Island, 2008.
-  M. L. Lapidus. "An overview of the complex fractal dimensions: From fractal strings to fractal drums, and back," *Contemporary Mathematics*, Vol. 731, Amer. Math. Soc., Providence, R.I., 2019, 143–269

References III

-  M. L. Lapidus and C. Pomerance. "The Riemann Zeta-Function and the One-Dimensional Weyl-Berry Conjecture for Fractal Drums," *Proc. London Math. Soc.* (3) 66 (1993), 41–69.
-  M. L. Lapidus and H. Maier. "The Riemann Hypothesis and Inverse Spectral Problems for Fractal Strings." *J. London Math. Soc.* (2) 52 (1995), 15–34.
-  M. L. Lapidus and M. van Frankenhuysen. *Fractal Geometry and Number Theory (Complex dimensions of fractal strings and zeros of zeta functions)*, Birkhäuser, Boston, 2000.
-  M. L. Lapidus and M. van Frankenhuysen. "Complex Dimensions of Self-Similar Fractal Strings and Diophantine Approximation," *J. Experimental Math.* (1) 12 (2003), 41–69.



References IV

-  M. L. Lapidus and M. van Frankenhuysen. *Fractal Geometry, Complex Dimensions and Zeta Functions: Geometry and Spectra of Fractal Strings*, second edition (of the 2006 edition), Springer Monographs in Mathematics, Springer, New York, 2012.
-  M. L. Lapidus and G. Radunović and D. Žubrinić. *Fractal Zeta Functions and Fractal Drums: Higher-Dimensional Theory of Complex Dimensions*, Springer Monographs in Mathematics, Springer, New York, 2017.
-  A. K. Lenstra and H. W. Lenstra and L. Lovász. “Factoring polynomials with rational coefficients,” *Mathematische Annalen* (4) 261 (1982), 515–534.
-  W. M. Schmidt. *Diophantine Approximation*, Springer, New York, 1980.

References V

-  J.-P. Serre. *A Course in Arithmetic*, English translation, Springer-Verlag, Berlin, 1973.
-  E. K. Voskanian. “On the Quasiperiodic Structure of the Complex Dimensions of Self-Similar Fractal Strings,” Ph. D. Dissertation, University of California, Riverside, 2019.