

Inner Minkowski Dimension of Products of Fractal Strings

Isaac Ashton, Joseph Ingenito, Peter Tonuzi

The College of New Jersey

What is a fractal string

Definition 1

A **fractal string** \mathcal{L} is a bounded open subset Ω of the real line.

What is a fractal string

Definition 1

A **fractal string** \mathcal{L} is a bounded open subset Ω of the real line.

$$\Omega = \bigcup_{j=1}^{\infty} \ell_j, \quad \text{Vol}_1(\Omega) = \sum_{j=1}^{\infty} \ell_j \implies \lim_{j \rightarrow \infty} \ell_j = 0.$$

What is a fractal string

Definition 1

A **fractal string** \mathcal{L} is a bounded open subset Ω of the real line.

$$\Omega = \bigcup_{j=1}^{\infty} \ell_j, \quad \text{Vol}_1(\Omega) = \sum_{j=1}^{\infty} \ell_j \implies \lim_{j \rightarrow \infty} \ell_j = 0.$$

$$\ell_1 \geq \ell_2 \geq \ell_3 \geq \cdots \geq 0$$

What is a fractal string

Definition 1

A **fractal string** \mathcal{L} is a bounded open subset Ω of the real line.

$$\Omega = \bigcup_{j=1}^{\infty} \ell_j, \quad \text{Vol}_1(\Omega) = \sum_{j=1}^{\infty} \ell_j \implies \lim_{j \rightarrow \infty} \ell_j = 0.$$

$$\ell_1 \geq \ell_2 \geq \ell_3 \geq \cdots \geq 0$$

Introduce the Cantor string so that the audience can see an example right away. A movie would be nice!

Cantor String

Dimension of fractal string

Inner-tubular neighborhood: $V(\epsilon) = \text{vol}_1 \{x \in \Omega \mid d(x, \partial\Omega) < \epsilon\}$

Dimension of fractal string

Inner-tubular neighborhood: $V(\epsilon) = \text{vol}_1 \{x \in \Omega \mid d(x, \partial\Omega) < \epsilon\}$

Use a movie of the Cantor String to illustrate the volume formula!

Dimension of fractal string

Inner-tubular neighborhood: $V(\epsilon) = \text{vol}_1\{x \in \Omega \mid d(x, \partial\Omega) < \epsilon\}$

Use a movie of the Cantor String to illustrate the volume formula!

Counting function: $N_\Omega(x) = \sum_{\ell_j^{-1} \leq x}$

$$D_{\mathcal{L}} = \inf\{\alpha \geq 0 \mid V(\epsilon) = O(\epsilon^{1-\alpha}) \text{ as } \epsilon \rightarrow 0^+\}$$

Dimension of fractal string

Inner-tubular neighborhood: $V(\epsilon) = \text{vol}_1\{x \in \Omega \mid d(x, \partial\Omega) < \epsilon\}$

Use a movie of the Cantor String to illustrate the volume formula!

Counting function: $N_\Omega(x) = \sum_{\ell_j^{-1} \leq x}$

$$D_{\mathcal{L}} = \inf\{\alpha \geq 0 \mid V(\epsilon) = O(\epsilon^{1-\alpha}) \text{ as } \epsilon \rightarrow 0^+\}$$

Lapidus, van Frankenhuijsen, [9, 10, 11]

$$D_{\mathcal{L}} = \inf\{\alpha \in \mathbb{R} \mid \sum_{j=1}^{\infty} \ell_j^\alpha < \infty\} \implies 0 < D_{\mathcal{L}} < 1$$

Cantor string volume

Fractal lawn

Definition 2

A **fractal lawn** \mathcal{L}^2 is the Cartesian product of two fractal strings
 $\Omega = \Omega_1 \times \Omega_2$.

Fractal lawn

Definition 2

A **fractal lawn** \mathcal{L}^2 is the Cartesian product of two fractal strings $\Omega = \Omega_1 \times \Omega_2$.

The Cartesian product of two fractal strings is a bounded open subset of \mathbb{R}^2 since a fractal string is a bounded open subset of \mathbb{R} .

$$\Omega_1 = \bigcup_{j=1}^{\infty} \ell_j, \Omega_2 = \bigcup_{i=1}^{\infty} l_i$$

$$\ell_1 \geq \ell_2 \geq \ell_3 \geq \cdots \geq 0, l_1 \geq l_2 \geq l_3 \geq \cdots \geq 0$$

Cantor lawn

Dimension of fractal lawn

Inner-tubular neighborhood: $V(\epsilon) = \text{vol}_2\{x \in \Omega \mid d(x, \partial\Omega) < \epsilon\}$

- $\partial\Omega = (\partial\Omega_1 \times \Omega_2) \cup (\Omega_1 \times \partial\Omega_2)$

Counting function: Counts the critical length for any $x > 0$

- $N_{\Omega_i}(x) = \sum_{j: \ell_j^{-1} \leq x} 1$

Useful Definitions:

- $\mathcal{W}_i(x) = \sum_{j: \ell_j^{-1} \leq x} \ell_j$

- $\mathcal{L}_i(x) = \sum_{j=1}^{N_{\Omega_i}(x)} \ell_j = \Omega_i - \mathcal{W}_i(x)$

Dimension of fractal lawn



- Partially covered volume

- $V_{\text{partial}}(\epsilon) = 2\epsilon N_{\Omega_1}(\frac{1}{2\epsilon}) \mathcal{L}_1(\frac{1}{2\epsilon}) + 2\epsilon N_{\Omega_2}(\frac{1}{2\epsilon}) \mathcal{L}_2(\frac{1}{2\epsilon})$

- Fully covered volume

- $V_{\text{full}}(\epsilon) = \Omega_1 \mathcal{W}_2(\frac{1}{2\epsilon}) + \Omega_2 \mathcal{W}_1(\frac{1}{2\epsilon})$

- Double counted volume




- $V_{\text{double}}(\epsilon) = \mathcal{W}_1(\frac{1}{2\epsilon}) \mathcal{W}_2(\frac{1}{2\epsilon})$

Dimension of fractal lawn




$$\begin{aligned} V(\epsilon) &= V_{\text{partial}}(\epsilon) + V_{\text{full}}(\epsilon) - V_{\text{double}}(\epsilon) \\ V(\epsilon) &= 2\epsilon N_{\Omega_1}(\tfrac{1}{2\epsilon}) \mathcal{L}_1(\tfrac{1}{2\epsilon}) + 2\epsilon N_{\Omega_2}(\tfrac{1}{2\epsilon}) \mathcal{L}_2(\tfrac{1}{2\epsilon}) + \Omega_1 \mathcal{W}_2(\tfrac{1}{2\epsilon}) + \Omega_2 \mathcal{W}_1(\tfrac{1}{2\epsilon}) - \\ &\quad \mathcal{W}_1(\tfrac{1}{2\epsilon}) \mathcal{W}_2(\tfrac{1}{2\epsilon}) \end{aligned}$$

Cantor Lawn Volume GIF





References I

-  D. A. Bini and L. Robol. “Solving secular and polynomial equations: A multiprecision algorithm.” *Journal of Computational and Applied Mathematics* 272 (2014), 276–292.
-  M. R. Bremner. *Lattice Basis Reduction: An Introduction to the LLL Algorithm and its Applications*, Taylor & Francis, Boca Raton, 2011.
-  M. L. Lapidus. “Fractal drum, inverse spectral problems for elliptic operators and a partial resolution of the Weyl-Berry conjecture.” *Transactions of the American Mathematical Society*. (2) 325 (1991), 465–529.





References II

-  M. L. Lapidus. “Vibrations of Fractal Drums, the Riemann Hypothesis, Waves in Fractal Media and the Weyl-Berry Conjecture.” In *Ordinary and Partial Differential Equations*, edited by B. D. Sleeman and R. J. Jarvis, Vol. IV, Proc. Twelfth Intern. Conf. (Dundee, Scotland, UK, June 1992), Pitman Research Notes in Math. Series 289, pp. 126–209, London: Longman Scientific and Technical, 1993.
-  M. L. Lapidus. *In Search of the Riemann Zeros: Strings, fractal membranes and noncommutative spacetimes*, American Mathematical Society, Providence, Rhode Island, 2008.
-  M. L. Lapidus. “An overview of the complex fractal dimensions: From fractal strings to fractal drums, and back,” *Contemporary Mathematics*, Vol. 731, Amer. Math. Soc., Providence, R.I., 2019, 143–269

References III

-  M. L. Lapidus and C. Pomerance. “The Riemann Zeta-Function and the One-Dimensional Weyl-Berry Conjecture for Fractal Drums,” *Proc. London Math. Soc.* (3) 66 (1993), 41–69.
-  M. L. Lapidus and H. Maier. “The Riemann Hypothesis and Inverse Spectral Problems for Fractal Strings.” *J. London Math. Soc.* (2) 52 (1995), 15–34.
-  M. L. Lapidus and M. van Frankenhuysen. *Fractal Geometry and Number Theory (Complex dimensions of fractal strings and zeros of zeta functions)*, Birkhäuser, Boston, 2000.
-  M. L. Lapidus and M. van Frankenhuysen. “Complex Dimensions of Self-Similar Fractal Strings and Diophantine Approximation,” *J. Experimental Math.* (1) 12 (2003), 41–69.

References IV

-  M. L. Lapidus and M. van Frankenhuijsen. *Fractal Geometry, Complex Dimensions and Zeta Functions: Geometry and Spectra of Fractal Strings*, second edition (of the 2006 edition), Springer Monographs in Mathematics, Springer, New York, 2012.
-  M. L. Lapidus and G. Radunović and D. Žubrinić. *Fractal Zeta Functions and Fractal Drums: Higher-Dimensional Theory of Complex Dimensions*, Springer Monographs in Mathematics, Springer, New York, 2017.
-  A. K. Lenstra and H. W. Lenstra and L. Lovász. “Factoring polynomials with rational coefficients,” *Mathematische Annalen* (4) 261 (1982), 515–534.
-  W. M. Schmidt. *Diophantine Approximation*, Springer, New York, 1980.

References V



J.-P. Serre. *A Course in Arithmetic*, English translation, Springer-Verlag, Berlin, 1973.



E. K. Voskanian. “On the Quasiperiodic Structure of the Complex Dimensions of Self-Similar Fractal Strings,” Ph. D. Dissertation, University of California, Riverside, 2019.