

The Ricci flow on closed 3 manifolds with strictly positive Ricci curvature

Jinghao Yu

Southern University of Science and Technology, China

2021.05.28

Background

The survey I do is related to the Poincare conjecture. In dimension 3, Poincare's conjecture states that:

If M is a simply-connected closed smooth manifold of dimension 3, then M and the 3-sphere \mathbb{S}^3 are diffeomorphic.

This conjecture has been solved in 2003 by Grigori Perelman. The Ricci flow plays an important role in Perelman proof. The thesis I do is to read R.S. Hamilton's paper in 1982, which is the origin of the Ricci flow.

Background

The survey I do is related to the Poincare conjecture. In dimension 3, Poincare's conjecture states that:

If M is a simply-connected closed smooth manifold of dimension 3, then M and the 3-sphere \mathbb{S}^3 are diffeomorphic.

This conjecture has been solved in 2003 by Grigori Perelman. The Ricci flow plays an important role in Perelman proof. The thesis I do is to read R.S. Hamilton's paper in 1982, which is the origin of the Ricci flow.

Introduction

In Hamilton's paper, he showed this conclusion:

Theorem (Main theorem)

Let (M, g) be a closed Riemannian manifold of dimension 3 which admits a strictly positive Ricci curvature. Then (M, g) also admits a metric of constant positive sectional curvature.

Here is a classical conclusion in Riemannian geometry to correlate the main theorem with Poincare conjecture.

In Hamilton's paper, he showed this conclusion:

Theorem (Main theorem)

Let (M, g) be a closed Riemannian manifold of dimension 3 which admits a strictly positive Ricci curvature. Then (M, g) also admits a metric of constant positive sectional curvature.

Here is a classical conclusion in Riemannian geometry to correlate the main theorem with Poincare conjecture.

Introduction

In Hamilton's paper, he showed this conclusion:

Theorem (Main theorem)

*Let (M, g) be a closed Riemannian manifold of dimension 3 which admits a strictly positive Ricci curvature. Then (M, g) also admits a metric of **constant positive sectional curvature**.*

Theorem (space form)

*Let M^n be a complete Riemannian manifold with **constant sectional curvature** K . Then the universal covering \tilde{M} of M , with the covering metric, is isometric to:*

- (a) *hyperbolic space \mathbb{H}^n , if $K = -1$,*
- (b) *Euclidean space \mathbb{R}^n , if $K = 0$,*
- (c) *sphere \mathbb{S}^n , if $K = 1$.*

As a special case, the main theorem states that

Corollary

If M is simply connected smooth manifold endowed with a strictly positive Ricci curvature, then M is diffeomorphic to a 3-sphere.

That is why R.S. Hamilton's result is a special case of Poincare conjecture.

As a special case, the main theorem states that

Corollary

If M is simply connected smooth manifold endowed with a strictly positive Ricci curvature, then M is diffeomorphic to a 3-sphere.

That is why R.S. Hamilton's result is a special case of Poincare conjecture.

The **Ricci flow** is a partial differential equation of g in a Riemannian manifold (M, g_0) satisfying

$$\frac{\partial g}{\partial t} = -2Rc,$$

$$g(0) = g_0.$$

A natural question: do we know the existence and uniqueness of Ricci flow?

The **Ricci flow** is a partial differential equation of g in a Riemannian manifold (M, g_0) satisfying

$$\frac{\partial g}{\partial t} = -2Rc,$$

$$g(0) = g_0.$$

A natural question: do we know the existence and uniqueness of Ricci flow?

Short time existence

Theorem (Hamilton, DeTurck)

If (M^n, g_0) is a closed Riemannian manifold, there exists a unique solution $g(t)$ to the Ricci flow defined on some positive time interval $[0, \epsilon)$ such that $g(0) = g_0$.

The solution time interval can be naturally extended to a maximal time $[0, T)$, $0 < T \leq \infty$.

Short time existence

Theorem (Hamilton, DeTurck)

If (M^n, g_0) is a closed Riemannian manifold, there exists a unique solution $g(t)$ to the Ricci flow defined on some positive time interval $[0, \epsilon)$ such that $g(0) = g_0$.

The solution time interval can be naturally extended to a maximal time $[0, T)$, $0 < T \leq \infty$.

Curvature estimate

The existence of short time solution allows us to discuss the variation of curvature under Ricci flow. In dimension 3, this estimate is relatively easy.

Curvature estimate

The existence of short time solution allows us to discuss the variation of curvature under Ricci flow. In dimension 3, this estimate is relatively easy.

First, the Riemannian curvature can be expressed as the combination of Ricci curvature:

Theorem

When M is of dimension 3,

$$R_{ijkl} = R_{il}g_{jk} + R_{jk}g_{il} - R_{ik}g_{jl} - R_{jl}g_{ik} - \frac{1}{2}R(g_{il}g_{jk} - g_{ik}g_{jl})$$

Curvature estimate

Second, at least one index of R_{ijkl} repeats twice. Many components vanish because of Bianchi identity (the symmetry of Riemannian curvature).

Curvature estimate

Second, at least one index of R_{ijkl} repeats twice. Many components vanish because of Bianchi identity (the symmetry of Riemannian curvature).

Let λ, μ, v be the eigenvalues of Ricci curvature, then

Corollary

The sectional curvature is the only possible nonzero component of Riemannian curvature. Its value can be represented by the eigenvalue of Ricci curvature. For example,

$$R_{1221} = \frac{1}{2}(\lambda + \mu - v).$$

Curvature estimate

Thus, if we want to flow g to be a metric with constant sectional curvature, it suffices to control Ricci curvature. Mainly, we need to do two things:

- (1) First, **at each point** we make λ, μ, ν approach each other as time flows.
- (2) Second, we control the **global** difference of R_{ij} . In some extent, $R_{max} - R_{min}$ could be test this because scalar curvature $R = \lambda + \mu + \nu$.

Curvature estimate

Thus, if we want to flow g to be a metric with constant sectional curvature, it suffices to control Ricci curvature. Mainly, we need to do two things:

- (1) First, **at each point** we make λ , μ , ν approach each other as time flows.
- (2) Second, we control the **global** difference of R_{ij} . In some extent, $R_{max} - R_{min}$ could be test this because scalar curvature $R = \lambda + \mu + \nu$.

Curvature estimate

Thus, if we want to flow g to be a metric with constant sectional curvature, it suffices to control Ricci curvature. Mainly, we need to do two things:

- (1) First, **at each point** we make λ , μ , ν approach each other as time flows.
- (2) Second, we control the **global** difference of R_{ij} . In some extent, $R_{max} - R_{min}$ could be test this because scalar curvature $R = \lambda + \mu + \nu$.

curvature estimate: first step

The L^2 -distance among λ, μ, v

$$(\lambda - \mu)^2 + (\mu - v)^2 + (\lambda - v)^2$$

is a good quantity to measure pointwise difference.

It could also be expressed as a **geometric** quantity. Let $S = |Rc|^2$ then

$$S - \frac{1}{3}R^2 = \frac{1}{3}[(\lambda - \mu)^2 + (\mu - v)^2 + (\lambda - v)^2]$$

curvature estimate: first step

The L^2 -distance among λ, μ, v

$$(\lambda - \mu)^2 + (\mu - v)^2 + (\lambda - v)^2$$

is a good quantity to measure pointwise difference.

It could also be expressed as a **geometric** quantity. Let $S = |Rc|^2$ then

$$S - \frac{1}{3}R^2 = \frac{1}{3}[(\lambda - \mu)^2 + (\mu - v)^2 + (\lambda - v)^2]$$

Curvature estimate: first step

It could also be expressed as a geometric quantity. Let $S = |Rc|^2$ then

$$S - \frac{1}{3}R^2 = \frac{1}{3}[(\lambda - \mu)^2 + (\mu - \nu)^2 + (\lambda - \nu)^2]$$

Theorem

Let M^3 be a closed 3-manifold, with strictly positive Ricci curvature. Under the variation of Ricci flow, \exists constant $\delta > 0$ and $C \in \mathbb{R}_+$ both depending only on the initial metric such that on $0 \leq t < T$ we have

$$S - \frac{1}{3}R^2 \leq CR^{2-\delta}.$$

This estimate turns the pointwise control of λ, μ, ν to the control of **scalar curvature R** .

Curvature estimate: first step

It could also be expressed as a geometric quantity. Let $S = |Rc|^2$ then

$$S - \frac{1}{3}R^2 = \frac{1}{3}[(\lambda - \mu)^2 + (\mu - \nu)^2 + (\lambda - \nu)^2]$$

Theorem

Let M^3 be a closed 3-manifold, with strictly positive Ricci curvature. Under the variation of Ricci flow, \exists constant $\delta > 0$ and $C \in \mathbb{R}_+$ both depending only on the initial metric such that on $0 \leq t < T$ we have

$$S - \frac{1}{3}R^2 \leq C R^{2-\delta}.$$

This estimate turns the pointwise control of λ, μ, ν to the control of **scalar curvature R** .

Curvature estimate: second step

In previous analysis, we know the estimate of scalar curvature, in particular, the estimate of $R_{max} - R_{min}$, is important. A natural idea is to estimate the gradient $\nabla_i R$ since it tells the local difference of R .

Curvature estimate: second step

In previous analysis, we know the estimate of scalar curvature, in particular, the estimate of $R_{max} - R_{min}$, is important. A natural idea is to estimate the gradient $\nabla_i R$ since it tells the local difference of R .

Curvature estimate: second step

In previous analysis, we know the estimate of scalar curvature, in particular, the estimate of $R_{max} - R_{min}$, is important. A natural idea is to estimate the gradient $\nabla_i R$ since it tells the local difference of R .

Curvature estimate: second step

In previous analysis, we know the estimate of scalar curvature, in particular, the estimate of $R_{max} - R_{min}$, is important. A natural idea is to estimate the gradient $\nabla_i R$ since it tells the local difference of R .

Theorem (gradient estimate)

Let M^3 be a closed Riemannian 3 manifold with positive Ricci curvature. For every $\eta > 0$, \exists constant $C = C(\eta, g(0))$ depending only on η and the initial value of the metric such that on $0 \leq t < T$ we have

$$|\nabla_i R|^2 \leq \eta R^3 + C$$

Curvature estimate: second step

In previous analysis, we know the estimate of scalar curvature, in particular, the estimate of $R_{max} - R_{min}$, is important. A natural idea is to estimate the gradient $\nabla_i R$ since it tells the local difference of R .

Theorem (gradient estimate)

Let M^3 be a closed Riemannian 3 manifold with positive Ricci curvature. For every $\eta > 0$, \exists constant $\mathcal{C} = \mathcal{C}(\eta, g(0))$ depending only on η and the initial value of the metric such that on $0 \leq t < T$ we have

$$|\nabla_i R|^2 \leq \eta R^3 + \mathcal{C}$$

Theorem

$R_{max}/R_{min} \rightarrow 1$ as $t \rightarrow T$.

Blow up

Up to now, everything goes right.

- difference of $\lambda, \mu, v \implies R_{max} - R_{min}$.
- R_{max}/R_{min} is also controlled.
- If we can control the value of R , then $R_{max} - R_{min}$ is well-controlled.
- However, R itself goes to infinity.

Blow up

Up to now, everything goes right.

- difference of $\lambda, \mu, v \implies R_{max} - R_{min}$.
- R_{max}/R_{min} is also controlled.
- If we can control the value of R , then $R_{max} - R_{min}$ is well-controlled.
- However, R itself goes to infinity.

Blow up

Up to now, everything goes right.

- difference of $\lambda, \mu, v \implies R_{max} - R_{min}$.
- R_{max}/R_{min} is also controlled.
- If we can control the value of R , then $R_{max} - R_{min}$ is well-controlled.
- However, R itself goes to infinity.

Up to now, everything goes right.

- difference of $\lambda, \mu, v \implies R_{max} - R_{min}$.
- R_{max}/R_{min} is also controlled.
- If we can control the value of R , then $R_{max} - R_{min}$ is well-controlled.
- However, R itself goes to infinity.

Blow up

Up to now, everything goes right.

- difference of $\lambda, \mu, v \implies R_{max} - R_{min}$.
- R_{max}/R_{min} is also controlled.
- If we can control the value of R , then $R_{max} - R_{min}$ is well-controlled.
- However, R itself goes to infinity.

Blow up

However, R itself goes to infinity.

Theorem

If M is a Riemannian 3-manifold and $R \geq \rho > 0$ at $t = 0$, then R goes to infinity before the time $3/2\rho$.

Proof: (Sketch)

$$\begin{aligned}\frac{\partial R}{\partial t} &\geq \Delta R + \frac{2}{3}R^2 \\ \frac{df}{dt} &= \frac{2}{3}f^2, \quad f(0) = \rho\end{aligned}$$

Apply comparison principle:

$$R \geq f = \frac{3\rho}{3 - 2\rho t} \rightarrow \infty \quad \text{as } t \rightarrow 3/2\rho.$$

Blow up

However, R itself goes to infinity.

Theorem

If M is a Riemannian 3-manifold and $R \geq \rho > 0$ at $t = 0$, then R goes to infinity before the time $3/2\rho$.

Proof: (Sketch)

$$\begin{aligned}\frac{\partial R}{\partial t} &\geq \Delta R + \frac{2}{3}R^2 \\ \frac{df}{dt} &= \frac{2}{3}f^2, \quad f(0) = \rho\end{aligned}$$

Apply comparison principle:

$$R \geq f = \frac{3\rho}{3 - 2\rho t} \rightarrow \infty \quad \text{as } t \rightarrow 3/2\rho.$$

Blow up

However, R itself goes to infinity.

Theorem

If M is a Riemannian 3-manifold and $R \geq \rho > 0$ at $t = 0$, then R goes to infinity before the time $3/2\rho$.

Proof: (Sketch)

$$\begin{aligned}\frac{\partial R}{\partial t} &\geq \Delta R + \frac{2}{3}R^2 \\ \frac{df}{dt} &= \frac{2}{3}f^2, \quad f(0) = \rho\end{aligned}$$

Apply comparison principle:

$$R \geq f = \frac{3\rho}{3 - 2\rho t} \rightarrow \infty \quad \text{as } t \rightarrow 3/2\rho.$$

Blow up

It could also be shown that the Riemannian curvature will blow up.

Theorem

Suppose (M, g_0) is a 3-manifold with strictly positive Ricci curvature. Then its Ricci flow has a unique solution on a maximal time interval $0 \leq t < T \leq 3/2\rho$, and $\max_M |R_{ijkl}| \rightarrow \infty$ as $t \rightarrow T$.

Normalized Ricci flow

To avoid the blow-up, we introduce the normalized Ricci flow.

Normalized Ricci flow

To avoid the blow-up, we introduce the normalized Ricci flow.

Definition (normalized Ricci flow)

The normalized equation of Ricci flow on (M^n, g_0) :

$$\frac{\partial}{\partial \tilde{t}} \tilde{g}_{ij} = \frac{2}{n} \tilde{r} \tilde{g}_{ij} - 2 \tilde{R}_{ij}$$

$$\tilde{g}(0) = g_0$$

where $\tilde{r} = \int \tilde{R} d\tilde{\mu} / \int d\tilde{\mu}$ is the average of scalar curvature.

Geometrically, the additional term keeps the volume of manifold M a constant.

Normalized Ricci flow

To avoid the blow-up, we introduce the normalized Ricci flow.

Definition (normalized Ricci flow)

The normalized equation of Ricci flow on (M^n, g_0) :

$$\frac{\partial}{\partial t} \tilde{g}_{ij} = \frac{2}{n} \tilde{r} \tilde{g}_{ij} - 2 \tilde{R}_{ij}$$

$$\tilde{g}(0) = g_0$$

where $\tilde{r} = \int \tilde{R} d\tilde{\mu} / \int d\tilde{\mu}$ is the average of scalar curvature.

Geometrically, the additional term keeps **the volume of manifold M a constant**.

Normalized Ricci flow

Geometrically, the additional term promises the volume of manifold M a constant.

Lemma

It promise $\tilde{T} = \infty$ and \tilde{R}_{\max} could be bounded by a constant C :

$$\tilde{R}_{\max}(\tilde{t}) \leq C < \infty \quad \text{on} \quad 0 \leq \tilde{t} < \tilde{T}.$$

Curvature estimate: normalized case

An interesting fact is that the normalized Ricci flow could be given by multiplying g with a appropriate coefficient $\varphi(t)$:

$$g \longrightarrow \tilde{g} = \varphi(t)g.$$

It helps us to transform many estimates from origin case to the normalized case.

Curvature estimate: normalized case

An interesting fact is that the normalized Ricci flow could be given by multiplying g with a appropriate coefficient $\varphi(t)$:

$$g \longrightarrow \tilde{g} = \varphi(t)g.$$

It helps us to transform many estimates from origin case to the normalized case.

Curvature estimate: normalized case

Lemma

\exists constants $C < \infty$ and $\delta > 0$ such that

$$\tilde{S} - \frac{1}{3}\tilde{R}^2 \leq Ce^{-\delta\tilde{t}}$$

Lemma

\exists constant $C < \infty$ and $\delta > 0$ such that

$$\tilde{R}_{\max} - \tilde{R}_{\min} \leq Ce^{-\delta\tilde{t}}.$$

Curvature estimate: normalized case

Lemma

\exists constants $C < \infty$ and $\delta > 0$ such that

$$\tilde{S} - \frac{1}{3}\tilde{R}^2 \leq Ce^{-\delta\tilde{t}}$$

Lemma

\exists constant $C < \infty$ and $\delta > 0$ such that

$$\tilde{R}_{\max} - \tilde{R}_{\min} \leq Ce^{-\delta\tilde{t}}.$$

Corollary

$|\tilde{R}_{ij} - \frac{1}{3}\tilde{r}\tilde{g}_{ij}| \leq Ce^{-\delta\tilde{t}}$ where $\tilde{r} = \int \tilde{R}d\tilde{\mu} / \int d\tilde{\mu}$.

Hence, if the metric \tilde{g} has good enough convergence such that we can exchange the order of taking limit and taking differential, then $\tilde{g}(\infty)$ has a constant sectional curvature.

Hence, if the metric \tilde{g} has **good enough convergence** such that we can exchange the order of taking limit and taking differential, then $\tilde{g}(\infty)$ has a constant sectional curvature.

Theorem

As $\tilde{t} \rightarrow \infty$ the metrics $\tilde{g}_{ij}(\tilde{t})$ converge to the smooth limit metric $\tilde{g}_{ij}(\infty)$ in C^∞ -topology. In special, the curvature $\tilde{R}_{ij}(\tilde{t})$ converge to the curvature $\tilde{R}_{ij}(\infty)$.

The limit Ricci curvature is

$$\tilde{R}_{ij}(\infty) = \frac{1}{3}\tilde{r}(\infty)\tilde{g}_{ij}(\infty)$$

Hence, M has a metric $\tilde{g}(\infty)$ with sectional curvature $\frac{1}{6}\tilde{r}(\infty)$.

Hence, if the metric \tilde{g} has good enough convergence such that we can exchange the order of taking limit and taking differential, then $\tilde{g}(\infty)$ has a constant sectional curvature.

Theorem

As $\tilde{t} \rightarrow \infty$ the metrics $\tilde{g}_{ij}(\tilde{t})$ converge to the smooth limit metric $\tilde{g}_{ij}(\infty)$ in C^∞ -topology. In special, the curvature $\tilde{R}_{ij}(\tilde{t})$ converge to the curvature $\tilde{R}_{ij}(\infty)$.

The limit Ricci curvature is

$$\tilde{R}_{ij}(\infty) = \frac{1}{3}\tilde{r}(\infty)\tilde{g}_{ij}(\infty)$$

Hence, M has a metric $\tilde{g}(\infty)$ with sectional curvature $\frac{1}{6}\tilde{r}(\infty)$.

Proof: (Sketch)

- By tensor interpolation inequality we have

$$\int |\tilde{\nabla}^n \tilde{R}c|^p d\tilde{\mu} \leq C e^{-\delta \tilde{t}} \quad \forall n, p \in \mathbb{N}_+$$

- By Sobolev's inequality we show: for every $n \in \mathbb{N}_+$ we have

$$\|\tilde{\nabla}^n \tilde{R}c(\tilde{t})\|_\infty \leq C e^{-\delta \tilde{t}}$$

for some constant $C < \infty$, $\delta > 0$ depending on n .

Proof: (Sketch)

- By tensor interpolation inequality we have

$$\int |\tilde{\nabla}^n \tilde{R}c|^p d\tilde{\mu} \leq C e^{-\delta \tilde{t}} \quad \forall n, p \in \mathbb{N}_+$$

- By Sobolev's inequality we show: for every $n \in \mathbb{N}_+$ we have

$$\|\tilde{\nabla}^n \tilde{R}c(\tilde{t})\|_\infty \leq C e^{-\delta \tilde{t}}$$

for some constant $C < \infty$, $\delta > 0$ depending on n .

Proof: (Sketch)

- By tensor interpolation inequality we have

$$\int |\tilde{\nabla}^n \tilde{R}c|^p d\tilde{\mu} \leq C e^{-\delta \tilde{t}} \quad \forall n, p \in \mathbb{N}_+$$

- By Sobolev's inequality we show: for every $n \in \mathbb{N}_+$ we have

$$\|\tilde{\nabla}^n \tilde{R}c(\tilde{t})\|_\infty \leq C e^{-\delta \tilde{t}}$$

for some constant $C < \infty$, $\delta > 0$ depending on n .

Take this control into the normalized Ricci flow, we can control $\|\partial^n \tilde{g}\|_\infty$:

Corollary

For all $n \in \mathbb{N}_+$,

$$\|\partial^n \tilde{g}\|_\infty \leq C e^{-\delta \tilde{t}}$$

for some constant $C < \infty$, $\delta > 0$ depending on n .

It implies that $\forall \alpha = (\alpha_1, \dots, \alpha_n) \in \mathbb{N}^n$ we have

$$\frac{\partial^{|\alpha|} g(t)}{\partial x^\alpha} \rightarrow \frac{\partial^{|\alpha|} g(\infty)}{\partial x^\alpha} \quad \text{uniformly w.r.t } M \text{ as } t \rightarrow \infty.$$

Take this control into the normalized Ricci flow, we can control $\|\partial^n \tilde{g}\|_\infty$:

Corollary

For all $n \in \mathbb{N}_+$,

$$\|\partial^n \tilde{g}\|_\infty \leq C e^{-\delta \tilde{t}}$$

for some constant $C < \infty$, $\delta > 0$ depending on n .

It implies that $\forall \alpha = (\alpha_1, \dots, \alpha_n) \in \mathbb{N}^n$ we have

$$\frac{\partial^{|\alpha|} g(t)}{\partial x^\alpha} \rightarrow \frac{\partial^{|\alpha|} g(\infty)}{\partial x^\alpha} \quad \text{uniformly w.r.t } M \text{ as } t \rightarrow \infty.$$

Theorem

As $\tilde{t} \rightarrow \infty$ the metrics $\tilde{g}_{ij}(\tilde{t})$ converge to the smooth limit metric $\tilde{g}_{ij}(\infty)$ in C^∞ -topology. In special, the curvature $\tilde{R}_{ij}(\tilde{t})$ converge to the curvature $\tilde{R}_{ij}(\infty)$.

Theorem

As $\tilde{t} \rightarrow \infty$ the metrics $\tilde{g}_{ij}(\tilde{t})$ converge to the smooth limit metric $\tilde{g}_{ij}(\infty)$ in C^∞ -topology. In special, the curvature $\tilde{R}_{ij}(\tilde{t})$ converge to the curvature $\tilde{R}_{ij}(\infty)$.

Corollary (main theorem)

The limit metric $\tilde{g}_{ij}(\infty)$ has constant positive sectional curvature.

Corollary

If M is simply connected smooth manifold endowed with a strictly positive Ricci curvature, then M is diffeomorphic to a 3-sphere.

Thank you

Thank you!