


Comparison of P values from linear and linear mixed models

This is provided by the `lmerTest` package as illustrated fully with the documentation example here.

```
require(lmerTest)
boxplot(Reaction~Days, data=sleepstudy, main="Reaction by Days",
        xlab="Days", ylab="Reaction", col="blue", border="black")
```



We see a trend of Reaction by Days, so it is reasonable to fit a simple linear regression,

```
l <- lm(Reaction ~ Days, sleepstudy)
s <- summary(l)
s
```

```
##
## Call:
## lm(formula = Reaction ~ Days, data = sleepstudy)
##
## Residuals:
##      Min       1Q   Median       3Q      Max
## -110.848  -27.483    1.546   26.142  139.953
##
## Coefficients:
##              Estimate Std. Error t value Pr(>|t|)
## (Intercept)   251.405      6.610  38.033 < 2e-16 ***
## Days           10.467      1.238   8.454 9.89e-15 ***
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```


```
##
## Residual standard error: 47.71 on 178 degrees of freedom
## Multiple R-squared:  0.2865, Adjusted R-squared:  0.2825
## F-statistic: 71.46 on 1 and 178 DF,  p-value: 9.894e-15
```

```
names(s)
```

```
## [1] "call"          "terms"          "residuals"      "coefficients"
## [5] "aliased"        "sigma"          "df"             "r.squared"
## [9] "adj.r.squared" "fstatistic"     "cov.unscaled"
```

Now the Subject effect is treated as a random; from

```
boxplot(Reaction~Subject, data=sleepstudy, main="Reaction by Subject",
        xlab="Subject", ylab="Reaction", col="orange", border="brown")
```



it is more appropriate to fit a random effect model

```
f <- lme4::lmer(Reaction ~ Days + (Days | Subject), sleepstudy)
s <- summary(f)
s
```

```
## Linear mixed model fit by REML ['lmerMod']
## Formula: Reaction ~ Days + (Days | Subject)
## Data: sleepstudy
##
## REML criterion at convergence: 1743.6
##
## Scaled residuals:
##      Min       1Q   Median       3Q      Max
```

```
## -3.9536 -0.4634 0.0231 0.4633 5.1793
##
## Random effects:
## Groups Name Variance Std.Dev. Corr
## Subject (Intercept) 611.90 24.737
## Days 35.08 5.923 0.07
## Residual 654.94 25.592
## Number of obs: 180, groups: Subject, 18
##
## Fixed effects:
## Estimate Std. Error t value
## (Intercept) 251.405 6.824 36.843
## Days 10.467 1.546 6.771
##
## Correlation of Fixed Effects:
## (Intr)
## Days -0.138
```

```
names(s)
```

```
## [1] "methTitle" "objClass" "devcomp" "isLmer"
## [5] "useScale" "logLik" "family" "link"
## [9] "ngrps" "coefficients" "sigma" "vcov"
## [13] "varcor" "AICtab" "call" "residuals"
## [17] "fitMsgs" "optinfo"
```

We saw the same estimate of effect but a larger standard error for Days in the linear mixed model compared to that in the linear regression model.

```
m <- lmerTest::lmer(Reaction ~ Days + (Days | Subject), sleepstudy)
class(m)
```

```
## [1] "lmerModLmerTest"
## attr(,"package")
## [1] "lmerTest"
```

```
s <-summary(m)
names(s)
```

```
## [1] "methTitle" "objClass" "devcomp" "isLmer"
## [5] "useScale" "logLik" "family" "link"
## [9] "ngrps" "coefficients" "sigma" "vcov"
## [13] "varcor" "AICtab" "call" "residuals"
## [17] "fitMsgs" "optinfo"
```

```
with(s,coefficients)[,5]
```

```
## (Intercept) Days
## 1.158238e-17 3.273014e-06
```