

Math 600 Fall 2015 Homework #5

Due Nov. 5, Thu. in class

Note: For the Euclidean space \mathbb{R}^n , consider the usual metric induced by the Euclidean norm $\|\cdot\|_2$ on \mathbb{R}^n , unless otherwise stated.

1. Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be continuous on \mathbb{R} . Which of the following sets are necessarily open, closed, compact, and connected (no proof is needed)?

$$(1) \{f(x) \in \mathbb{R} \mid x \geq 0\}; \quad (2) \{f(x) \in \mathbb{R} \mid 0 \leq x \leq 1\}.$$

2. Let $\|\cdot\|_2$ be the standard Euclidean norm on \mathbb{R}^n , and $\|\cdot\|_\alpha$ be an arbitrary norm on \mathbb{R}^n . Recall that both $\|\cdot\|_2$ and $\|\cdot\|_\alpha$ are continuous on $(\mathbb{R}^n, \|\cdot\|_2)$.

(1) Let $\mathbb{S}^{n-1} := \{x \in \mathbb{R}^n : \|x\|_2 = 1\}$. Show that \mathbb{S}^{n-1} is closed in $(\mathbb{R}^n, \|\cdot\|_2)$. (*Hint:* write \mathbb{S}^{n-1} as a suitable inverse image.)

(2) Show that \mathbb{S}^{n-1} is compact in $(\mathbb{R}^n, \|\cdot\|_2)$.

(3) Show that there exist $x^* \in \mathbb{S}^{n-1}$ and $y^* \in \mathbb{S}^{n-1}$ such that

$$\|x^*\|_\alpha = \max_{x \in \mathbb{S}^{n-1}} \|x\|_\alpha, \quad \|y^*\|_\alpha = \min_{x \in \mathbb{S}^{n-1}} \|x\|_\alpha.$$

Further show that $\|x^*\|_\alpha > 0$ and $\|y^*\|_\alpha > 0$.

(4) Let $c_1 := \|x^*\|_\alpha$ and $c_2 := \|y^*\|_\alpha$. Use (3) to show that

$$c_2 \|x\|_2 \leq \|x\|_\alpha \leq c_1 \|x\|_2, \quad \forall x \in \mathbb{R}^n.$$

(*Hint:* for any $0 \neq x \in \mathbb{R}^n$, $\frac{x}{\|x\|_2} \in \mathbb{S}^{n-1}$.)

(5) Let $\|\cdot\|_\beta$ be another norm on \mathbb{R}^n . Use (4) to show that there exist $\kappa_1, \kappa_2 > 0$ such that

$$\kappa_2 \|x\|_\beta \leq \|x\|_\alpha \leq \kappa_1 \|x\|_\beta, \quad \forall x \in \mathbb{R}^n.$$

★ *This result shows that all norms on \mathbb{R}^n are equivalent.*

3. Let $f : \mathbb{R}^n \rightarrow \mathbb{R}^m$ be continuous on \mathbb{R}^n . Define the function $g : \mathbb{R}^n \rightarrow \mathbb{R}^n \times \mathbb{R}^m$ as:

$$g(x) := (x, f(x)), \quad \forall x \in \mathbb{R}^n.$$

(1) Show g is continuous on \mathbb{R}^n . (*Hint:* consider the sequential criterion.)

(2) For the function f given above, define the following set (called the *graph* of f):

$$\mathcal{S} := \{(x, f(x)) \mid x \in \mathbb{R}^n\} \subseteq \mathbb{R}^n \times \mathbb{R}^m.$$

Show that \mathcal{S} is closed and connected.

4. Show that a metric space (M, d) is connected if and only if the only open and closed sets in M are M and the empty set.

Miscellaneous practice problems: Do not submit

1. Let $f : [a, b] \rightarrow [a, b]$ be continuous on $[a, b]$, where the real numbers $a < b$. Show that f has a fixed point on $[a, b]$, namely, there exists $c \in [a, b]$ such that $f(c) = c$.
2. Let $f, g : (M, d) \rightarrow \mathbb{R}$ be continuous on M . Then (i) the set $\{x \in M : f(x) = g(x)\}$ is closed in (M, d) ; and (ii) the set $\{x \in M : f(x) > 2g(x)\}$ is open in (M, d) .