

a
Stress-associated
Ciliated-like
Epithelial core
Proliferative
AT2-like
Other
UMAP 2
UMAP 1
b
Stress-associated | 3,780 cells | 14 patients
Ciliated-like | 437 cells | 6 patients
Epithelial core | 3,187 cells | 14 patients
Proliferative | 2,585 cells | 8 patients
AT2-like | 3,653 cells | 13 patients
Other | 52 cells | 10 patients
BACH1 detected
DoRothEA BACH1
Hypoxia
Glycolysis
NRF2/oxidative
Heme metabolism
ROS pathway
OXPHOS
EMT
Proliferation
Column z
c
Original scores
ρ=0.91 [0.75, 0.97]
n=16; P=1.1×10−6
Shared genes removed
ρ=0.87 [0.65, 0.95]
n=16; P=1.3×10−5
Dataset and QC residuals
ρ=0.51 [0.01, 0.80]
n=16; P=0.046
Dataset, QC and state-PC residuals
ρ=0.14 [-0.39, 0.59]
n=16; P=0.617
Full residuals, >=20-cell patients
ρ=0.01 [-0.52, 0.54]
n=14; P=0.970
Spearman ρ
d
Stress-associated
ρ=0.88 [0.46, 0.98]
n=8; Holm P=0.022
Epithelial core
ρ=0.89 [0.26, 0.99]
n=6; Holm P=0.067
AT2-like
ρ=0.10 [-0.65, 0.75]
n=8; Holm P=0.840
Within-state patient-level ρ