

152_project code

Lucy Wang, Michelle Seo

5/3/2020

```
##EDA
##Plots/EDA
library(dplyr)

##
## Attaching package: 'dplyr'
## The following objects are masked from 'package:plyr':
##
##   arrange, count, desc, failwith, id, mutate, rename, summarise,
##   summarize
## The following objects are masked from 'package:stats':
##
##   filter, lag
## The following objects are masked from 'package:base':
##
##   intersect, setdiff, setequal, union

##filling in total_staff male/female calculation
##invalid data = 0 staff members
data15_18$TOTALSTAFFFF <- data15_18$CORRSTAFFFF+data15_18$OTHERSTAFFFF
data15_18$TOTALSTAFFM <- data15_18$TOTALSTAFF - data15_18$TOTALSTAFFFF
data15_18$STRATUM <- as.factor(data15_18$STRATUM)

summary(data15_18$TOTALSTAFF)

##   Min. 1st Qu.  Median    Mean 3rd Qu.    Max.
##   -9.0   32.0   78.0  181.2  190.5 11449.0

summary(data15_18$CORRSTAFFFF)

##   Min. 1st Qu.  Median    Mean 3rd Qu.    Max.
##   -9.00    8.00   17.00  42.71  37.00 4370.00

summary(data15_18$OTHERSTAFFFF)

##   Min. 1st Qu.  Median    Mean 3rd Qu.    Max.
##   -9.00    2.00    7.00  21.73  22.00  513.00


indices <- which(data15_18$TOTALSTAFF==0)

data15_18$ratio <- data15_18$TOTALSTAFFM/data15_18$TOTALSTAFFFF
data15_18$ratio[data15_18$ratio<=0 | is.infinite(data15_18$ratio)] <- NA
```

```
ratio <- data15_18$ratio[is.na(data15_18$ratio)==F]
ratio <- ratio[is.infinite(ratio)==F]

ggplot()+
  geom_histogram(aes(ratio), bins = 30, fill="#4281a4")+
  ggtitle("2015-2018 Male to Female ratio of Jail Staff")+
  geom_vline(aes(xintercept=mean(ratio)))+
  scale_x_continuous(breaks = seq(0,25, 2.5))
```

2015–2018 Male to Female ratio of Jail Staff




```
data15_18 %>% group_by(STRATUM) %>% summarise(ct = n(), avg=mean(ratio, na.rm = T))
```

```
## # A tibble: 10 x 3
##   STRATUM    ct  avg
##   <fct>    <int> <dbl>
## 1 1      1386  2.33
## 2 2      133  2.76
## 3 3       70  1.99
## 4 4       32  2.07
## 5 5       44  2.19
## 6 7      828  2.15
## 7 8      306  1.98
## 8 9      239  2.43
## 9 10     219  1.89
## 10 12     262  1.81
```


```
mean(ratio)
```

```
## [1] 2.202839
```

```
sts <- boxplot.stats(data15_18$CORRSTAFF)$stats
## Correctional Staff
ggplot(data15_18, aes(x=STRATUM, y=CORRSTAFF)) +
  geom_boxplot(aes(group=STRATUM), fill="#9cafb7") +
  scale_x_discrete(breaks = seq(1,12,1)) +
  coord_cartesian(ylim = c(sts*1.05, sts/1.05)) +
  ggtitle("2015-2018 Correctional Staff Distributions by Stratum")
```



```
## Other Staff
sts <- boxplot.stats(data15_18$OTHERSTAFF)$stats
## Gender comparison per type
ggplot(data15_18, aes(x=STRATUM, y=OTHERSTAFF)) +
  geom_boxplot(aes(group=STRATUM), fill="#9cafb7") +
  scale_x_discrete(breaks = seq(1,12,1)) +
  coord_cartesian(ylim = c(sts*1.05, sts/1.05)) +
  ggtitle("2015-2018 Other Staff Distributions by Stratum")
```



##ADP/Staff Ratio


```
data13_18$STRATUM <- sapply(data13_18$STRATUM, round)
data13_18$year <- as.factor(data13_18$year)
data13_18$STRATUM <- as.factor(data13_18$STRATUM)

##inmate to staff ratio
data13_18$ratios2 <- data13_18$ADP/data13_18$TOTALSTAFF
data13_18$ratios2[is.infinite(data13_18$ratios2) | data13_18$ratios2 <=0] <- NA

grpRatio <- data13_18 %>% group_by(year, STRATUM) %>% summarise(avg = mean(ratios2, na.rm=T))
grpRatio <- grpRatio[is.na(grpRatio$avg)==F,]

ggplot(grpRatio, aes(year, avg))+
  geom_line(aes(group = STRATUM, color=STRATUM))+
  geom_point(size = 1)+
  ggtitle("Average Inmate to Staff ratio by year and Stratum")+
  xlab("Year")+
  ylab("Ratio")
```

Average Inmate to Staff ratio by year and Stratum



grpRatio

```
## # A tibble: 44 x 3
## # Groups:   year [6]
##   year STRATUM avg
##   <fct> <fct> <dbl>
## 1 2013 1      4.23
## 2 2013 12     4.21
## 3 2014 1      4.00
## 4 2014 12     4.19
## 5 2015 1      3.78
## 6 2015 2      3.92
## 7 2015 3      3.91
## 8 2015 4      4.93
## 9 2015 5      2.44
## 10 2015 7      4.37
## # ... with 34 more rows
```

```
data15_18$ratios3 <- data15_18$ADP/data15_18$CORRSTAFF
```

```
data15_18$ratios3[is.infinite(data15_18$ratios3) | data15_18$ratios3 <=0] <- NA
```

```
grpRatio <- data15_18 %>% group_by(year, STRATUM) %>% summarise(avg = mean(ratios3, na.rm=T))
```

```
grpRatio <- grpRatio[is.na(grpRatio$avg)==F,]
```

grpRatio

```
## # A tibble: 40 x 3
## # Groups:   year [4]
##   year STRATUM avg
```

```
##      <chr> <fct>    <dbl>
## 1 2015 1          4.98
## 2 2015 2          4.74
## 3 2015 3          5.01
## 4 2015 4          5.68
## 5 2015 5          3.24
## 6 2015 7          5.61
## 7 2015 8          5.13
## 8 2015 9          3.80
## 9 2015 10         2.56
## 10 2015 12        5.00
## # ... with 30 more rows

##Inmate staff Ratio
##Inmate to staff
ratio_fn <- function(dat) {
  ##Returns Ratio and SE
  strat_design<-svydesign(ids = ~1, strata = ~STRATUM, weights=~FINALWT, data = dat)
  r <- svyratio(~ADP, ~TOTALSTAFF, strat_design, na.rm=T)
  c(unname(r$ratio)[[1]], sqrt(unname(r$var)[[1]]))
}

yrs <- as.character(2013:2018)

estimates <- data.frame()

for (i in 1:6) {
  yr <- yrs[i]
  dat <- data13_18[data13_18$year==yr,]
  estimates <- rbind(estimates, ratio_fn(dat))
}


colnames(estimates) <- c("Ratio", "SE")
estimates$year <- yrs

estimates

##      Ratio      SE year
## 1 6.080715 0.5229804 2013
## 2 5.635782 0.4399577 2014
## 3 3.386003 0.1367137 2015
## 4 3.231654 0.1535714 2016
## 5 3.303769 0.1683169 2017
## 6 3.330533 0.1797114 2018

ggplot(estimates, aes(x = year, y = Ratio)) +
  geom_point(size = 2) +
  geom_errorbar(aes(ymin = Ratio-1.96*SE, ymax = Ratio+1.96*SE))+
  ggtitle("2013-18 Confidence intervals for Inmate to Staff Ratios")
```

2013–18 Confidence intervals for Inmate to Staff Ratios



```
estimates <- estimates[c("year", "Ratio", "SE")]
estimates
```

```
##   year   Ratio      SE
## 1 2013 6.080715 0.5229804
## 2 2014 5.635782 0.4399577
## 3 2015 3.386003 0.1367137
## 4 2016 3.231654 0.1535714
## 5 2017 3.303769 0.1683169
## 6 2018 3.330533 0.1797114
```

##Functions for estimation

```
yrs <- as.character(2015:2018)
##Female to Male
estimates <- data.frame()

ratio_fn2 <- function(dat) {
  ##Returns Ratio and SE
  strat_design<-svydesign(ids = ~1, strata = ~STRATUM, weights=~FINALWT, data = dat)
  r <- svyratio(~TOTALSTAFFM, ~TOTALSTAFFF, strat_design, na.rm=T)
  c(unnname(r$ratio)[[1]], sqrt(unnname(r$var)[[1]]))
}

for (i in 1:4) {
  yr <- yrs[i]
  dat <- data15_18[data15_18$year==yr,]
  estimates <- rbind(estimates, ratio_fn2(dat))
}
```

```


}

colnames(estimates) <- c("Ratio", "SE")
estimates$year <- yrs

ggplot(estimates, aes(x = year, y = Ratio)) +
  geom_point(size = 2) +
  geom_errorbar(aes(ymax = Ratio+1.96*SE, ymin = Ratio-1.96*SE))+
  ggtitle("2015-18 Confidence intervals for Male to Female Staff")

```

2015-18 Confidence intervals for Male to Female Staff



```

estimates <- estimates[c("year", "Ratio", "SE")]
data15_18$year <- as.factor(data15_18$year)
summary(aov(ratio~year, data15_18))

```

```

##              Df Sum Sq Mean Sq F value Pr(>F)
## year           3      9  2.846   1.145  0.33
## Residuals  3401  8456  2.486
## 114 observations deleted due to missingness

```

```
##t-test
```

```
library(tidyverse)
```

```

## -- Attaching packages ----- tidyverse 1.3.0 --
## v tibble  3.0.1    v stringr 1.4.0
## v tidyr   1.0.0    v forcats 0.4.0
## v purrr   0.3.3

```



```

## -- Conflicts ----- tidyverse_conflicts() --
## x dplyr::arrange() masks plyr::arrange()
## x purrr::compact() masks plyr::compact()
## x dplyr::count() masks plyr::count()
## x tidyr::expand() masks Matrix::expand()
## x dplyr::failwith() masks plyr::failwith()
## x dplyr::filter() masks stats::filter()
## x dplyr::id() masks plyr::id()
## x dplyr::lag() masks stats::lag()
## x dplyr::mutate() masks plyr::mutate()
## x tidyr::pack() masks Matrix::pack()
## x dplyr::rename() masks plyr::rename()
## x dplyr::summarise() masks plyr::summarise()
## x dplyr::summarize() masks plyr::summarize()
## x tidyr::unpack() masks Matrix::unpack()

library(ggpubr)

##
## Attaching package: 'ggpubr'

## The following object is masked from 'package:plyr':
##
## mutate

library(rstatix)

##
## Attaching package: 'rstatix'

## The following objects are masked from 'package:plyr':
##
## desc, mutate

## The following object is masked from 'package:stats':
##
## filter

#Inmate to staff
dat <- data15_18[data15_18$STRATUM != 1 & data15_18$STRATUM != 2,]
summary(aov(ratio~STRATUM, dat))

##
## Df Sum Sq Mean Sq F value Pr(>F)
## STRATUM 7 63 9.013 4.106 0.000173 ***
## Residuals 1933 4243 2.195
## ---
## Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
## 59 observations deleted due to missingness

inm_staff<-data13_18[data13_18$year != "2013" & data13_18$year != "2014",]

t_test <- inm_staff %>% pairwise_t_test(ratios2~STRATUM, pool.sd = F)
t_test <- t_test[t_test$p.adj < 0.05,]
t_test <- t_test[t_test$group1!=1 & t_test$group2 !=12,c("group1", "group2","p.adj")]
head(t_test)

## # A tibble: 6 x 3

```

```
##      group1 group2      p.adj
##      <chr>  <chr>      <dbl>
## 1 2        5        3.96e- 7
## 2 2        7        2.00e- 3
## 3 2       10        1.03e-11
## 4 3        5        1.55e- 5
## 5 3        7        2.00e- 3
## 6 3       10        6.26e- 9
```

Q1Q2

```
library(ggplot2)
library(dplyr)
library(gridExtra)
```

```
##
## Attaching package: 'gridExtra'

## The following object is masked from 'package:dplyr':
##
##      combine
library(survey)
```

#Loading data

```
fifteen_eighteen_data <- read.csv("data15_18.csv")
thirteen_fourteen_data <- read.csv("data13_14.csv")
total_data <- read.csv("data13_18.csv")
```

#Individual Years

```
thirteen_data <- total_data %>% filter(year==2013)
fourteen_data <- total_data %>% filter(year==2014)
fifteen_data <- total_data %>% filter(year==2015)
sixteen_data <- total_data %>% filter(year==2016)
seventeen_data <- total_data %>% filter(year==2017)
eighteen_data <- total_data %>% filter(year==2018)
```

#Total staff values using survey

```
survey_design13 <- svydesign(data=thirteen_data, ids=~1, weights=thirteen_data$FINALWT, strata=thirteen_data$TOTALSTAFF, design=survey_design13)
svytotal(thirteen_data$TOTALSTAFF, design=survey_design13)
```

```
##      total      SE
## [1,] 120274 13653
```

```
survey_design14 <- svydesign(data=fourteen_data, ids=~1, weights=fourteen_data$FINALWT, strata=fourteen_data$TOTALSTAFF, design=survey_design14)
svytotal(fourteen_data$TOTALSTAFF, design=survey_design14)
```

```
##      total      SE
## [1,] 131122 13688
```

```
survey_design15 <- svydesign(data=fifteen_data, ids=~1, weights=fifteen_data$FINALWT, strata=fifteen_data$TOTALSTAFF, design=survey_design15)
svytotal(fifteen_data$TOTALSTAFF, design=survey_design15)
```

```
##      total      SE
## [1,] 212489 12634
```

```

survey_design16 <- svydesign(data=sixteen_data, ids=~1, weights=sixteen_data$FINALWT, strata=sixteen_data$TOTALSTAFF, design=survey_design16)

##          total      SE
## [1,] 226294 14067

survey_design17 <- svydesign(data=seventeen_data, ids=~1, weights=seventeen_data$FINALWT, strata=seventeen_data$TOTALSTAFF, design=survey_design17)

##          total      SE
## [1,] 225690 14748

survey_design18 <- svydesign(data=eighteen_data, ids=~1, weights=eighteen_data$FINALWT, strata=eighteen_data$TOTALSTAFF, design=survey_design18)

##          total      SE
## [1,] 221564 14500

#Assault Analysis for years 13-14

thirteendata <- thirteen_fourteen_data %>% filter(year==2013)
fourteendata <- thirteen_fourteen_data %>% filter(year==2014)
thirteensurvey <- svydesign(data=thirteendata, ids=~1, weights=thirteendata$FINALWT, strata=thirteendata$TOTALSTAFF, design=thirteensurvey)

##          total      SE
## ASSAULTCORR 3000.9 443.15

svytotal(~ASSAULTCORR, design=thirteensurvey)

##          total      SE
## ASSAULTCORR 159.18 36.256

fourteensurvey <- svydesign(data=fourteendata, ids=~1, weights=fourteendata$FINALWT, strata=fourteendata$TOTALSTAFF, design=fourteensurvey)

##          total      SE
## ASSAULTCORR 2687.1 411.78

svytotal(~ASSAULTCORR, design=fourteensurvey)

##          total      SE
## ASSAULTCORR 93.392 26.478

#Chisq Test of Independence

sum(thirteendata$TOTALSTAFF-thirteendata$OTHERSTAFF)

## [1] 85808

sum(thirteendata$OTHERSTAFF)

## [1] 27551

sum(thirteendata$ASSAULTCORR)

## [1] 2854

sum(thirteendata$ASSAULTTOTRSTAFF)

## [1] 150

```

```
correctional_staff <- c(2854, 85808)
other_staff <- c(150, 27551)
chisquared_table <- data.frame(rbind(correctional_staff, other_staff))
names(chisquared_table) <- c("assault", "no_assault")
chisquared_table
```

```
##           assault no_assault
## correctional_staff    2854    85808
## other_staff           150    27551
```

```
chisq.test(chisquared_table)
```

```
##
## Pearson's Chi-squared test with Yates' continuity correction
##
## data:  chisquared_table
## X-squared = 600.58, df = 1, p-value < 2.2e-16
#Chisq Test of Independence
```

```
emptyvector <- c()
emptyvector[thirteendata$ADP < 2000] <- "one"
emptyvector[thirteendata$ADP > 2001 & thirteendata$ADP < 4001] <- "two"
emptyvector[thirteendata$ADP > 4001] <- "three"
thirteendata$category <- emptyvector
gettingvals1 <- thirteendata %>% select(category, ASSAULTCORR, ASSAULTOTRSTAFF) %>% group_by(category)
thirteendata$noassault <- thirteendata$TOTALSTAFF - thirteendata$ASSAULTCORR - thirteendata$ASSAULTOTRSTAFF
gettingvals2 <- thirteendata %>% select(noassault, category) %>% group_by(category) %>% summarise(sum=s)

assault <- c(1470, 878, 598)
no_assault <- c(60027, 29406, 20415)

chisquared_table <- data.frame(rbind(assault, no_assault))
names(chisquared_table) <- c("one", "two", "three")
chisquared_table
```

```
##           one  two three
## assault    1470  878  598
## no_assault 60027 29406 20415
```

```
chisq.test(chisquared_table)
```

```
##
## Pearson's Chi-squared test
##
## data:  chisquared_table
## X-squared = 26.216, df = 2, p-value = 2.029e-06
#Exploratory Analysis
```

```
staff_data <- total_data %>% select(year, TOTALSTAFF, OTHERSTAFF) %>% group_by(year) %>% summarise_all(
staff_data
```


```
## # A tibble: 6 x 3
##   year TOTALSTAFF OTHERSTAFF
##   <int>      <int>      <int>
## 1  2013    113359    27551
## 2  2014    122533    29766
```

```
## 3 2015      155897      32276
## 4 2016      165000      35260
## 5 2017      160366      33840
## 6 2018      156268      34786
```


#reference <https://stackoverflow.com/questions/1249548/side-by-side-plots-with-ggplot2>

```
a <- ggplot(data=staff_data)+geom_line(aes(x = year, y = TOTALSTAFF))+xlab("Year")+ylab("Number of People")
b <- ggplot(data=staff_data)+geom_line(aes(x = year, y = OTHERSTAFF))+xlab("Year")+ylab("Number of People")
grid.arrange(a, b)
```

Total Staff Members in Jail 2013–2018



Other Staff Members in Jail 2013–2018



```
assault_data <- thirteen_fourteen_data %>% select(ASSAULTCORR, ASSAULTOTRSTAFF, DEATHSTAFFYN)
assault_data$DEATHSTAFFYN[assault_data$DEATHSTAFFYN==9] <- 0
summary(assault_data)
```

```
##  ASSAULTCORR      ASSAULTOTRSTAFF  DEATHSTAFFYN
##  Min.   : 0.000    Min.   :0.000    Min.   :0.0000
##  1st Qu.: 0.000    1st Qu.:0.000    1st Qu.:0.0000
##  Median : 0.000    Median :0.000    Median :0.0000
##  Mean   : 3.097    Mean   :0.137    Mean   :0.4819
##  3rd Qu.: 0.000    3rd Qu.:0.000    3rd Qu.:0.0000
##  Max.   :217.000    Max.   :9.000    Max.   :2.0000
```

```
assault_data <- thirteen_fourteen_data %>% select(ASSAULTCORR, ASSAULTOTRSTAFF, DEATHSTAFFYN)
assault_data$DEATHSTAFFYN[assault_data$DEATHSTAFFYN==9] <- 0
assault_data <- assault_data %>% select(ASSAULTCORR, ASSAULTOTRSTAFF, DEATHSTAFFYN) %>% summarise_all(
assault_count <- data.frame("Type"=c("Assault on Corr Officers", "Assault on Other", "Deaths by Assault"))
assault_count
```

```
##
## 1 Assault on Corr Officers 2526
## 2 Assault on Other 88
## 3 Deaths by Assaults 836
```

```
ggplot(data=assault_count)+geom_col(aes(x=Type, y=Count))+xlab("Assault Type")+ylab("Number of Cases")+
```

