Causal Inference Methods and Case Studies

STAT24630

Jingshu Wang

Lecture 10

- Non-compliance in randomized experiment

  - Intention-to-treat effect

  - Principal stratification

  - The monotonicity and exclusion restriction assumptions

  - CATE estimand and the moment-based estimator

  - Connection with two-stage least square estimator

  - Weak instrument

The Sommer-Zeger vitamin A supplement data

- In principle, 8 different possible values of the triple $(Z_i, W_i^{\text{obs}}, Y_i^{\text{obs}})$

- Non-compliance: $Z_i \neq W_i^{\text{obs}}$

Assignment Z_i	Vitamin Supplements W_i^{obs}	Survival Y_i^{obs}	Number of Units ($N = 23{,}682$)
0	0	0	74
0	0	1	11,514
1	0	0	34
1	0	1	2385
1	1	0	12
1	1	1	9663

Three types of traditional analyses

Method	Estimate	Calculation	Row Comparison
ITT	0.0026	$= \dfrac{2385 + 9663}{12 + 9663 + 34 + 2385} - \dfrac{11514}{74 + 11514}$	3, 4, 5, & 6 vs. 1 & 2
As-treated	0.0065	$= \dfrac{9663}{12 + 9663} - \dfrac{11514 + 2385}{74 + 11514 + 34 + 2385}$	5 & 6 vs. 1, 2, 3, & 4
Per-protocol	0.0052	$= \dfrac{9663}{12 + 9663} - \dfrac{11514}{74 + 11514}$	5 & 6 vs. 1 & 2

Assignment Z_i	Vitamin Supplements W_i^{obs}	Survival Y_i^{obs}	Number of Units $(N = 23{,}682)$
0	0	0	74
0	0	1	11,514
1	0	0	34
1	0	1	2385
1	1	0	12
1	1	1	9663

Can we provide a better analysis?

Setup of the framework

- Treatment assignment (randomized encouragement): $Z_i \in \{0,1\}$

- Potential treatment variables: $(W_i(0), W_i(1))$
  - $W_i(z) = 1$: would receive the treatment if $Z_i = z$
  - $W_i(z) = 0$: would not receive the treatment if $Z_i = z$
- Observed treatment received: $W_i^{\text{obs}} = W_i(Z_i)$

- In the non-compliance setting, there are two "treatment": assignment to treatment and receipt of treatment
- Potential outcomes: $Y_i(z, w)$ potential outcome if unit is assigned to z and receive w
- Observed outcome: $Y_i^{\text{obs}} = Y_i(Z_i, W_i(Z_i))$
- We can also write the potential outcomes as $Y_i(z) = Y_i(z, W_i(z))$

Underlying assumptions

- No interference assumption for $W_i(z)$ and $Y_i(z, w)$

- Randomization of the treatment assignment
$$\left(Y_i(0,0), Y_i(0,1), Y_i(1,0), Y_i(1,1), W_i(0), W_i(1)\right) \perp Z_i$$

- We don't have
$$\left(Y_i(0,0), Y_i(0,1), Y_i(1,0), Y_i(1,1)\right) \perp W_i^{\text{obs}}$$
or
$$\left(Y_i(0,0), Y_i(0,1), Y_i(1,0), Y_i(1,1)\right) \perp W_i^{\text{obs}}|Z_i$$
We don't know why some units comply and some units don't

- Compliance can not be controlled by randomized experiment

Intention-to-treat (ITT) effects

- ITT effect on the receipt of treatment level

$$\text{ITT}_{W,i} = W_i(1) - W_i(0) \qquad \text{ITT}_W = \frac{1}{N}\sum_{i=1}^{N}\text{ITT}_{W,i} = \frac{1}{N}\sum_{i=1}^{N}\left(W_i(1) - W_i(0)\right)$$

- ITT effect on the outcome of primary interest

$$\text{ITT}_{Y,i} = Y_i(1, W_i(1)) - Y_i(0, W_i(0))$$

$$\text{ITT}_Y = \frac{1}{N}\sum_{i=1}^{N}\text{ITT}_{Y,i} = \frac{1}{N}\sum_{i=1}^{N}\left(Y_i(1, W_i(1)) - Y_i(0, W_i(0))\right)$$

Statistical analysis of ITT effects

- Statistical analyses of these effects follow exactly the same procedures as before

$$\widehat{\text{ITT}_W} = \overline{W}_1^{obs} - \overline{W}_0^{obs} \qquad \widehat{\mathbb{V}}(\widehat{\text{ITT}_W}) = \frac{s_{W,0}^2}{N_0} + \frac{s_{W,1}^2}{N_1}$$

$$s_{W,z}^2 = \sum_{i:W_i^{obs}=z} \frac{\left(W_i^{obs} - \overline{W}_z^{obs}\right)^2}{N_z - 1} = \frac{N_z}{N_z - 1} \overline{W}_z^{obs}(1 - \overline{W}_z^{obs})$$

$$\widehat{\text{ITT}_Y} = \overline{Y}_1^{obs} - \overline{Y}_0^{obs} \qquad \widehat{\mathbb{V}}(\widehat{\text{ITT}_Y}) = \frac{s_{Y,1}^2}{N_1} + \frac{s_{Y,0}^2}{N_0}$$

- We can also use regression analyses

- Drawback is that it estimates 'programmatic effectiveness' instead of 'biologic efficacy'

Principal stratification

- Stratify individuals based on their compliance status
- Four principal strata
  - Compliers (co) $\left(W_i(0), W_i(1)\right) = (0,1)$

  - Non-compliers (nc) $\begin{cases} \text{Always} - \text{takers (at)} \ \left(W_i(0), W_i(1)\right) = (1,1) \\ \text{never} - \text{takers (nt)} \ \left(W_i(0), W_i(1)\right) = (0,0) \\ \text{Defiers (df)} \qquad\qquad \left(W_i(0), W_i(1)\right) = (1,0) \end{cases}$

		$W_i(1)$	
		0	1
$W_i(0)$	0	nt	co
	1	df	at

Principal stratification

- Principal stratification depends on latent states of units!!

- Can not decide which principal strata each unit belong to simply based on the observed data
  - **one-sided compliance**: control group can never receive the treatment, but treatment group may not follow the assignment

		Assignment Z_i	
		0	1
Receipt of treatment W_i^{obs}	0	nt/co	nt
	1	–	co

- In general

		Z_i	
		0	1
W_i^{obs}	0	nt/co	nt/df
	1	at/df	at/co

ITT effect decomposition

- Denote the proportion of individuals that fall into each strata as $\pi_c, \pi_a, \pi_n, \pi_d$
  - For one-sided compliance data, $\pi_a = \pi_d = 0$
- Define the average ITT effect for each strata
  - For the treatment received $\text{ITT}_{W,c}, \text{ITT}_{W,a}, \text{ITT}_{W,n}, \text{ITT}_{W,d}$
  $$\text{ITT}_{W,c} = 1, \text{ITT}_{W,a} = 0, \text{ITT}_{W,n}=0, \text{ITT}_{W,d} = -1$$
  - For the primary outcome $\text{ITT}_c, \text{ITT}_a, \text{ITT}_n, \text{ITT}_d$

- For the ITT effect on treatment received

$$\text{ITT}_W = \sum_{i=1}^{N} \text{ITT}_{W,i} = \pi_c \text{ITT}_{W,c} + \pi_a \text{ITT}_{W,a} + \pi_n \text{ITT}_{W,n} + \pi_d \text{ITT}_{W,d} = \pi_c - \pi_d$$

- For the ITT effect on primary outcome

$$\text{ITT}_Y = \sum_{i=1}^{N} \text{ITT}_{Y,i} = \pi_c \text{ITT}_c + \pi_a \text{ITT}_a + \pi_n \text{ITT}_n + \pi_d \text{ITT}_d$$

Instrumental variables (IV)

Assumptions for Z_i being a valid IV:

- Randomization: $Z_i \in \{0,1\}$ are randomized
- Monotonicity: no defiers $\pi_d = 0$ or $W_i(0) \leq W_i(1)$ for all i
- Exclusion restriction: instrument affects the outcome only through treatment

$$Y_i(1,w) = Y_i(0,w)$$

- For always takers

$$\text{ITT}_{Y,i} = Y_i\big(1, W_i(1)\big) - Y_i\big(0, W_i(0)\big) = Y_i(1,1) - Y_i(0,1) = 0$$

so $\text{ITT}_a = 0$

- For never takers

$$\text{ITT}_{Y,i} = Y_i\big(1, W_i(1)\big) - Y_i\big(0, W_i(0)\big) = Y_i(1,0) - Y_i(0,0) = 0$$

so $\text{ITT}_n = 0$

- For compliers

$$\text{ITT}_{Y,i} = Y_i\big(1, W_i(1)\big) - Y_i\big(0, W_i(0)\big) = Y_i(1,1) - Y_i(0,0)$$

ITT_c is the average ``biological efficacy'' of the treatment on compliers

- Relevance: $\pi_c > 0$

Instrumental variables

Assumptions of Z_i being a valid IV :

- Randomization: $Z_i \in \{0,1\}$ are randomized
- Monotonicity: no defiers $\pi_d = 0$ or $W_i(0) \le W_i(1)$ for all i
- Exclusion restriction: instrument affects the outcome only through treatment
$$Y_i(1, w) = Y_i(0, w)$$

- Relevance: $\pi_c > 0$

- Then $\text{ITT}_W = \pi_c$ and $\text{ITT}_Y = \pi_c \text{ITT}_c + \pi_a \text{ITT}_a + \pi_n \text{ITT}_n + \pi_d \text{ITT}_d = \pi_c \text{ITT}_c$

- IV estimand: ITT_c Complier average treatment effect (CATE)
$$\text{CATE} = \text{ITT}_c = \frac{\text{ITT}_Y}{\text{ITT}_W}$$

- We can identify ITT_Y and ITT_W, so ITT_c is also identifiable
- CATE $\ne$ ATE unless ATE for noncompliers equals CATE

The monotonicity assumption

- Monotonicity: no defiers $\pi_d = 0$ or $W_i(0) \leq W_i(1)$ for all i

- Defiers are individuals who never follow treatment assignment no matter what treatment assignment is

- For one-sided compliance data, monotonicity is always satisfied

- Check the monotonicity assumption in general:
  - $\text{ITT}_W = \pi_c - \pi_d > 0$ if $\pi_d = 0$, so if we can reject the null that $\text{ITT}_W \geq 0$, then monotonicity assumption must fail
  - Otherwise, the monotonicity assumption is not testable

  - Need to decide whether the monotonicity assumption is reasonable or not based on domain knowledge

The exclusion restriction assumption

- Exclusion restriction: instrument affects the outcome only through treatment

$$Y_i(1, w) = Y_i(0, w)$$

- Double-blinding in experiments guarantees exclusion restriction

- The assumption in general is not testable, and need subject-matter knowledge to judge
- The subject-matter knowledge needed is often more subtle than that required to evaluate SUTVA

Moment-based IV estimator

- Causal estimand assuming a super population

$$\text{CATE} = \frac{\text{ITT}_Y}{\text{ITT}_W} = \frac{\mathbb{E}(Y_i(1) - Y_i(0))}{\mathbb{E}(W_i(1) - W_i(0))}$$

- Method-of-moment estimator:

$$\hat{\tau}^{iv} = \frac{\widehat{\text{ITT}}_Y}{\widehat{\text{ITT}}_W}$$

- How to estimate the variance of $\hat{\tau}^{iv}$?
  - Estimates $\widehat{\text{ITT}}_Y$ and $\widehat{\text{ITT}}_W$ are correlated because they use the same dataset
  - We can approximate the variance of $\hat{\tau}^{iv}$ when N is large (from delta method):

$$\mathbb{V}(\hat{\tau}^{iv}) \approx \frac{1}{\text{ITT}_W^4} \left\{ \text{ITT}_W^2 \mathbb{V}(\widehat{\text{ITT}}_Y) + \text{ITT}_Y^2 \mathbb{V}(\widehat{\text{ITT}}_W) - 2\text{ITT}_Y \text{ITT}_W \text{Cov}(\widehat{\text{ITT}}_W, \widehat{\text{ITT}}_Y) \right\}$$

  - Plug-in estimator of $\mathbb{V}(\hat{\tau}^{iv})$:

$$\widehat{\mathbb{V}}(\hat{\tau}^{iv}) \approx \frac{1}{\widehat{\text{ITT}}_W^4} \left\{ \widehat{\text{ITT}}_W^2 \widehat{\mathbb{V}}(\widehat{\text{ITT}}_Y) + \widehat{\text{ITT}}_Y^2 \widehat{\mathbb{V}}(\widehat{\text{ITT}}_W) - 2\widehat{\text{ITT}}_Y \widehat{\text{ITT}}_W \widehat{\text{Cov}}(\widehat{\text{ITT}}_W, \widehat{\text{ITT}}_Y) \right\}$$

Estimate the covariance

- The covariance between $\widehat{\text{ITT}}_Y$ and $\widehat{\text{ITT}}_W$:

$$\text{Cov}\left(\widehat{\text{ITT}}_W, \widehat{\text{ITT}}_Y\right) = \text{Cov}\left(\overline{W}_1^{\text{obs}} - \overline{W}_0^{\text{obs}}, \overline{Y}_1^{\text{obs}} - \overline{Y}_0^{\text{obs}}\right)$$

$$= \frac{\text{Cov}\left(Y_i(1), W_i(1)\right)}{N_1} + \frac{\text{Cov}\left(Y_i(0), W_i(0)\right)}{N_0}$$

- To estimate the covariance $\text{Cov}\left(Y_i(z), W_i(z)\right)$ for $z = 0,1$:

$$\widehat{\text{Cov}}\left(Y_i(z), W_i(z)\right) = \frac{1}{N_z - 1} \sum_{i:Z_i=z} \left(W_i^{\text{obs}} - \overline{W}_z^{\text{obs}}\right)\left(Y_i^{\text{obs}} - \overline{Y}_z^{\text{obs}}\right)$$

- So, the plug-in estimator is

$$\widehat{\text{Cov}}\left(\widehat{\text{ITT}}_W, \widehat{\text{ITT}}_Y\right) = \sum_{z=0}^{1} \frac{\sum_{i:Z_i=z}\left(W_i^{\text{obs}} - \overline{W}_z^{\text{obs}}\right)\left(Y_i^{\text{obs}} - \overline{Y}_z^{\text{obs}}\right)}{N_z(N_z - 1)}$$

- 95% confidence interval of CATE: $\left[\hat{\tau}^{iv} - 1.96\sqrt{\widehat{\mathbb{V}}(\hat{\tau}^{iv})}, \hat{\tau}^{iv} + 1.96\sqrt{\widehat{\mathbb{V}}(\hat{\tau}^{iv})}\right]$

Simplification for one-sided compliance data

As $W_i(0) \equiv 0$, we have

- $\widehat{\text{ITT}}_W = \bar{W}_1^{\text{obs}} - \bar{W}_0^{\text{obs}} = \bar{W}_1^{\text{obs}}$

- $\widehat{\mathbb{V}}\left(\widehat{\text{ITT}}_W\right) = \frac{s_{W,1}^2}{N_1} = \frac{\overline{W}_1^{\text{obs}}(1-\overline{W}_1^{\text{obs}})}{(N_1-1)}$ as $s_{W,0}^2 = 0$

- $\widehat{\text{Cov}}\left(\widehat{\text{ITT}}_W, \widehat{\text{ITT}}_Y\right) = \frac{\sum_{i:Z_i=1}\left(W_i^{\text{obs}}-\bar{W}_1^{\text{obs}}\right)\left(Y_i^{\text{obs}}-\bar{Y}_1^{\text{obs}}\right)}{N_1(N_1-1)}$

Result in Sommer-Zeger Vitamin Supplement data

ITT Estimates:

- $N_1 = 12 + 9663 + 34 + 2385 = 12094, N_0 = 74 + 11514 = 11588$

- $\widehat{\text{ITT}}_W = \bar{W}_1^{\text{obs}} = \frac{12+9663}{N_1} = 0.8, \widehat{\mathbb{V}}(\widehat{\text{ITT}}_W) = \frac{\bar{W}_1^{\text{obs}}(1-\bar{W}_1^{\text{obs}})}{N_1-1} = \frac{0.2*0.8}{12094*12093} = 0.0036^2$

- $\widehat{\text{ITT}}_Y = \frac{2385+9663}{N_1} - \frac{11514}{N_0} = 0.0026, \widehat{\mathbb{V}}(\widehat{\text{ITT}}_Y) = \sum_{z=0}^{1} \frac{\bar{Y}_z^{\text{obs}}(1-\bar{Y}_z^{\text{obs}})}{N_z-1} = 0.0009^2$

- 95% CI of ITT_Y: $(0.0008, 0.0044)$

CATE estimate:

- $\hat{\tau}^{iv} = \frac{0.0026}{0.8} = 0.0032$

- $\widehat{\text{Cov}}(\widehat{\text{ITT}}_W, \widehat{\text{ITT}}_Y) = -0.0000017$ (correlation -0.05)
- $\widehat{\mathbb{V}}(\hat{\tau}^{iv}) = 0.0012^2$

Assignment Z_i	Vitamin Supplements W_i^{obs}	Survival Y_i^{obs}	Number of Units $(N = 23,682)$
0	0	0	74
0	0	1	11,514
1	0	0	34
1	0	1	2385
1	1	0	12
1	1	1	9663

- 95% CI of CATE: $(0.0010, 0.0055)$
- The as-protocol or as-treated estimates are possibly biased up

Two-stage least square (2SLS) estimator

- Conventionally in econometrics, researchers use a two-stage least square approach for CATE

- The two-stage least square estimator is equivalent to $\hat{\tau}^{iv}$

- Two-stage least square
  - Stage 1: regress W_i^{obs} on Z_i : the coefficient of Z_i is ITT_W (regression with no covariate) the fitted coefficient on Z_i is $\widehat{\text{ITT}}_W$
  - Stage 1: regress Y_i^{obs} on Z_i : the coefficient of Z_i is ITT_Y (regression with no covariate) the fitted coefficient on Z_i is $\widehat{\text{ITT}}_Y$
  - Take the ratio of estimated coefficients, which is exactly $\hat{\tau}^{iv}$

- We can generalize 2SLS to incorporate covariates when estimating ITT_W and ITT_Y

The Angrist draft lottery data

Background

- Policy makers are interested in whether veterans are adequately compensated for their service.
- Angrist (1991) aims to measure the long-term labor market consequences of military service during the Vietnam era
- Question: estimate the causal effect of serving in the military during the Vietnam War on earnings
- We can not directly compare veterans and non-veterans, as they can be systematically different in unobserved ways, even after adjusting for differences in observed covariates

- Serving in the military or not during the Vietnam War could not randomized directly, but the military draft lottery of the Vietnam War was randomized
- This is called a natural experiment

The Angrist draft lottery data

Randomization

- For each birth year of birth cohort 1950-1952, a random ordering of the 365 days was constructed, a cutoff number was pre-determined, young men of that birth year who had a birth date with order before the cutoff "won" the lottery
- Randomization of birth date, instead of the individuals

- Theoretically, each date should be a unit, but in the book example, we treat each individual as a unit and consider the experiment as a completely randomized experiment (it's actually a stratified cluster randomized experiment).
Consequence is that we will tend to under-estimate the uncertainty of the causal estimator.

Relevance and two-sided non-compliance:

- Drafted individuals were required to prepare to serve in the military if fit for the service
- To serve the military, drafted individuals need to pass medical tests and have achieved minimum education level
- Individuals who were not draft eligible also can volunteer to serve in the military

The Angrist draft lottery data

	Non-Veterans ($N_c = 6,675$)				Veterans ($N_t = 2,030$)			
	Min	Max	Mean	(S.D.)	Min	Max	Mean	(S.D.)
Draft eligible	0	1	0.24	(0.43)	0	1	0.40	(0.49)
Yearly earnings (in $1,000's)	0	62.8	11.8	(11.5)	0	50.7	11.7	(11.8)
Earnings positive	0	1	0.88	(0.32)	0	1	0.91	(0.29)
Year of birth	50	52	51.1	(0.8)	50	52	50.9	(0.8)

Check assumptions
- **Monotonicity**: appears to be a reasonable assumption
  - The lottery numbers impose restrictions on individuals' behaviors.
  - Monotonicity means that no one responds to these restrictions by serving only if they are not required to do so
  - It is possible that there are some individuals who would be willing to volunteer if they are not drafted but would resist the draft if required, but it must be a very small fraction and are likely ignorable

The Angrist draft lottery data

	Non-Veterans ($N_c = 6{,}675$)				Veterans ($N_t = 2{,}030$)			
	Min	Max	Mean	(S.D.)	Min	Max	Mean	(S.D.)
Draft eligible	0	1	0.24	(0.43)	0	1	0.40	(0.49)
Yearly earnings (in $1,000's)	0	62.8	11.8	(11.5)	0	50.7	11.7	(11.8)
Earnings positive	0	1	0.88	(0.32)	0	1	0.91	(0.29)
Year of birth	50	52	51.1	(0.8)	50	52	50.9	(0.8)

Check assumptions
- **Exclusion restriction**: may be questionable
  - Consider the never-takers
  - Some never-takers are due to medical exemptions or exemptions due to their education or career choices. For them, the lottery numbers would likely not affect their future behaviors and the outcome
  - Some never-takers did have exemptions but changed their plan (enter graduate school or move to Canada) if they had a low draft number to avoid serving in the military. For them, exclusion restriction can be violated.

Analysis results

ITT Estimates:

- $\widehat{\text{ITT}}_W = 0.1460$, $\widehat{\mathbb{V}}\left(\widehat{\text{ITT}}_W\right) = 0.0108^2$

- $\widehat{\text{ITT}}_Y = -0.2129$, $\widehat{\mathbb{V}}\left(\widehat{\text{ITT}}_W\right) = \sum_{z=0}^{1} \frac{\bar{Y}_z^{\text{obs}}(1-\bar{Y}_z^{\text{obs}})}{N_z(N_z-1)} = 0.1980^2$

- 95% CI of ITT_Y: $(-0.6010, 0.1752)$

If we are willing to assume monotonicity and exclusion restriction
CATE estimate:

- $\hat{\tau}^{iv} = \frac{-0.2129}{0.1460} = -1.46$

- $\widehat{\mathbb{V}}\left(\hat{\tau}^{iv}\right) = 1.36^2$

- 95% CI of CATE: $(-4.13, 1.2)$

Weak instrument

- The instrumental variable is a weak instrument if the compliance probability (π_c or ITT_W) is small

- Problems using weak instrument
  - $\hat{\tau}^{iv} = \frac{\widehat{\text{ITT}}_Y}{\widehat{\text{ITT}}_W}$: the ratio is very unstable. If ITT_W is close to 0, then a small error (perturbation) in $\widehat{\text{ITT}}_W$ can lead to a large error in $\hat{\tau}^{iv}$
  - If the exclusion restriction assumption is violated, the bias in our estimator assuming exclusion restriction is inversely proportional to π_c

- How to identify weak instrument?
  - In the first stage linear regression model $W_i^{\text{obs}} = \alpha + \pi_c W_i + \varepsilon_i$, calculate the F-statistics to test whether $\pi_c = 0$
  - A rule of thumb is to check whether the F-statistics it larger to 10 or not.
  - F-statistics smaller than 10 indicates a weak instrument