


# Statistics How To

Statistics How To Videos



## Variance Inflation Factor



[Statistics Definitions](#) > Variance Inflation Factor

**CERTIFIED SERVICE**

**8-Quart Oil Change Package**  
**\$84.95\***  
ACDelco GM OE dexos1 Full Synthetic Oil Change and Tire Rotation

[Explore Offers](#)
[Schedule Service](#)

Chevrolet Cadillac of Turnersville

Expires 12/18/2022

[See Terms and Conditions](#)

You may want to read this article first: [What is Multicollinearity?](#)


## What is a Variance Inflation Factor?

A variance inflation factor(VIF) detects [multicollinearity](#) in [regression analysis](#). Multicollinearity is when there's [correlation](#) between predictors (i.e. [independent variables](#)) in a model; it's presence can adversely affect your regression results. The VIF estimates how much the variance of a regression coefficient is inflated due to multicollinearity in the model.

VIFs are usually calculated by software, as part of regression analysis. You'll see a VIF column as part of the output. VIFs are calculated by taking a predictor, and regressing it against every other predictor in the model. This gives you the [R-squared](#) values, which can then be plugged into the VIF formula. "i" is the predictor you're looking at (e.g.  $x_1$  or  $x_2$ ):

$$VIF = \frac{1}{1 - R_i^2}$$





## Interpreting the Variance Inflation Factor

Variance inflation factors range from 1 upwards. The numerical value for VIF tells you (in decimal form) what **percentage** the variance (i.e. the **standard error** squared) is inflated for each coefficient. For example, a VIF of 1.9 tells you that the variance of a particular coefficient is 90% bigger than what you would expect if there was no multicollinearity — if there was no correlation with other predictors.

A **rule of thumb** for interpreting the variance inflation factor:

- 1 = not correlated.
- Between 1 and 5 = moderately correlated.
- Greater than 5 = highly correlated.

Exactly how large a VIF has to be before it causes issues is a subject of debate. What is known is that the more your VIF increases, the less reliable your regression results are going to be. In general, a VIF above 10 indicates high correlation and is cause for concern. Some authors suggest a more conservative level of 2.5 or above.

Sometimes a high VIF is no cause for concern at all. For example, you can get a high VIF by including products or powers from other variables in your regression, like  $x$  and  $x^2$ . If you have high VIFs for **dummy variables** representing **nominal variables** with three or more categories, those are usually not a problem.


## References

Dodge, Y. (2008). [The Concise Encyclopedia of Statistics](#). Springer.


Everitt, B. S.; Skrondal, A. (2010), [The Cambridge Dictionary of Statistics](#), Cambridge University Press.

### CITE THIS AS:

**Stephanie Glen**. "Variance Inflation Factor" From [StatisticsHowTo.com](#): Elementary Statistics for the rest of us!  
<https://www.statisticshowto.com/variance-inflation-factor/>



**Need help with a homework or test question?** With [Chegg Study](#), you can get step-by-step solutions to your questions from an expert in the field. Your first 30 minutes with a Chegg tutor is free!



**Using a Frequency Distribution Table to Find Probabilities**

In a sample of 43 students:

- 15 had brown hair.
- 10 had black hair.
- 16 had blond hair.
- 2 had red hair.

Use a frequency distribution table to find the probability a person has neither red nor blond hair.

**Learn More**

Comments? Need to post a correction? Please [Contact Us](#).


- < Collinear Definition: What is Collinearity?
- > Multicollinearity: Definition, Causes, Examples

**NEED HELP with a homework problem? [CLICK HERE!](#)**

**Need help with a homework or test question?** With [Chegg Study](#), you can get step-by-step solutions to your questions from an expert in the field. Your first 30 minutes with a Chegg tutor is free!

**NEED HELP with a homework problem? [CLICK HERE!](#)**

Feel like “cheating” at Statistics? Check out our [Practically Cheating Statistics Handbook](#), which gives you hundreds of easy-to-follow answers in a convenient e-book.



**GET the Statistics & Calculus Bundle at a 40% discount!**

<https://www.statisticshowto.com/variance-inflation-factor/>

Using a Frequency Distribution Table to Find Probabilities

In a sample of 43 students:  
  
15 had brown hair.  
10 had black hair.  
16 had blond hair.  
2 had red hair.  
  
Use a frequency distribution table to find  
the probability a person has neither red nor blond hair.

Find the United card that's right for you

UNITED Club MileagePlus

D. BARRETT

VISA

UNITED Quest MileagePlus

D. BARRETT

VISA Signature

UNITED Explorer MileagePlus

D. BARRETT

VISA Signature

UNITED Gateway MileagePlus

D. BARRETT

VISA Signature

Learn more

NEED HELP with a homework problem? [CLICK HERE!](#)

The Practically Cheating

Calculus

Handbook

Feel like "cheating" at Calculus? Check out our [Practically Cheating Calculus Handbook](#), which gives you hundreds of easy-to-follow answers in a convenient e-book.



So easy, it's

Practically Cheating

GET the Statistics & Calculus Bundle at a 40% discount!



**Using a Frequency Distribution Table to Find Probabilities**

In a sample of 43 students:

- 15 had brown hair.
- 10 had black hair.
- 16 had blond hair.
- 2 had red hair.

Use a frequency distribution table to find the probability a person has neither red nor blond hair.

## Latest articles

[Census in Statistics: Simple Definition](#)

[Bimodal Distribution: What is it?](#)

[TI 83 for Statistics: Easy Steps for Common Problems](#)

[Blinding vs. Double Blind in Statistics: Definition, Examples](#)

[Hypothesis Test in Excel for the Population Mean \(Large Sample\)](#)


[Probability Distribution Table: What is it?](#)

[What is a Decile?](#)

[Normal Distribution Probability in Excel](#)

**NEED HELP with a homework problem? [CLICK HERE!](#)**





Using a Frequency Distribution Table to Find Probabilities

In a sample of 43 students:

15 had brown hair.  
10 had black hair.  
16 had blond hair.  
2 had red hair.

Use a frequency distribution table to find  
the probability a person has neither red nor blond hair.