

Intralobular Innervation and Lipocyte Contractility in the Liver

TAKATO UENO, MD* AND KYUICHI TANIKAWA, MD

From the Second Department of Medicine, Kurume University School of Medicine, Kurume, Japan

Date accepted: 6 June 1996

ABSTRACT

In the liver of humans, guinea pigs, cats, and tupaia, nerve endings are distributed all over the hepatic lobules from the portal spaces to the centrallobular spaces. Nerve endings in the intralobular spaces are located mainly in the space of Disse, and are closely related to lipocytes. In the human liver, various neurotransmitters such as substance P (SP) exist in the nerve endings. Lipocytes are believed to contract through these substances. In fact, the contraction of lipocytes is induced by SP. Moreover, lipocytes possess endothelin (ET) receptors (ET_A, ET_B), and the cells are contracted by ET-1 by way of ET receptors in the autocrine or paracrine mechanism. Contraction of lipocytes seems to be related to the enhancement of the intracellular Ca²⁺ and inositol phosphates. In addition, α -smooth muscle actin, which is a contractile protein, exists in the cytoplasm of lipocytes. Lipocyte contractility may be similar to that of vascular smooth muscle cells. On the other hand, prostaglandin E₂, iloprost, and adrenomedullin cause the elevation of c-AMP levels in lipocytes and relax the cells. In addition, lipocytes produce nitric oxide (NO) and inhibit contractility by an autocrine mechanism related to NO. In this way, lipocytes appear to be associated with the regulation of hepatic sinusoidal microcirculation by contraction and relaxation. In the cirrhotic liver, intralobular innervation is decreased or absent, but ET-1 and NO are overexpressed. These phenomena indicate that lipocytes may play an important role in the sinusoidal microcirculation through these agents rather than through intralobular innervation in liver cirrhosis. ©Elsevier Science Inc. 1997 Nutrition 1997;13:141–148

Key words: hepatic innervation, lipocyte (hepatic stellate cell), contractility

INTRODUCTION

Lipocytes located in the space of Disse are also called fat-storing cells, Ito cells, stellate cells, and perisinusoidal cells.^{1–4} These cells have many cytoplasmic processes and surround hepatic sinusoidal endothelial cells. The structure of the hepatic sinusoid resembles that of a capillary (Fig. 1).⁵ Hepatic sinusoidal endothelial cells are homologous to capillary endothelial cells, and lipocytes are homologous to pericytes located around capillaries. Pericytes regulate the microcirculation of the capillary by contraction or relaxation in response to neurotransmitters released from nerve endings or peptides such as endothelin (ET).⁶ Lipocytes, as well as pericytes, appear to be associated with nerves and function to regulate the hepatic sinusoidal microcirculation by contraction and relaxation. In this review, we discuss the relationship between intralobular innervation and lipocytes, and the contractility of lipocytes.

INTRALOBULAR INNERVATION AND LIPOCYTES IN THE LIVER

With regard to intrahepatic innervation, previous reports have indicated a wide variation in the distribution of nerves according to species studied and methods employed.^{7–14} In the human liver, nerve endings are distributed all over the hepatic lobules from the portal spaces to the centrolobular spaces.^{15–23} In the portal and centrolobular spaces, naked endings devoid of a Schwann sheath appose some mesenchymal cells such as fibroblasts, myofibroblasts, and smooth muscle cells, which form part of the adventitial wall of blood vessels and bile ducts (Figs. 2 and 3). These cell types are believed to control some aspects of hemodynamics in the portal and centrolobular spaces.

Nerve endings in the intralobular spaces are located mainly in the space of Disse. Nerve endings at this site are in close proximity to lipocytes and hepatocytes.^{15,19,21} Moreover, nerve endings are seen near hepatic sinusoidal endothelial cells (Fig. 4).¹⁹ In 1968, Ito and Shibasaki¹⁵ first reported the close relationship between

The original work presented in this review was supported in part by a Grant (No.06454269) for Scientific Research from the Japanese Ministry of Education.

Correspondence to: Takato Ueno, MD, Second Department of Medicine, Kurume University School of Medicine, 67 Asahi-machi, Kurume 830, Japan.




FIG. 1. Electron micrograph ($\times 11,000$) showing a sinusoid in the human liver. Hepatic sinusoidal endothelial cells with fenestrae (arrows) in the cytoplasm are enfolding by the cytoplasmic processes of lipocytes (L). A nerve bundle (N) closely apposes the process of a lipocyte. S, sinusoid.

nerve endings and lipocytes in the human liver. More recently, Bioulac-Sage et al.²⁰ also showed that nerve endings are located close to lipocytes in the human liver. Guinea pigs, cats, and tupaia have an intralobular innervation similar to that of humans.^{8,10} However, in the livers of mice and cats, intralobular innervation is not as abundant, and light and electron microscopic observation of such innervation is difficult.⁷⁻¹⁴

Generally, branches of the vagus nerve (a parasympathetic nerve), the splanchnic nerve (a sympathetic nerve), and sometimes the phrenic nerve enter the human liver. In the human liver, an adrenergic type of innervation has been suggested mainly at the lobular level,^{16,18} although some cholinergic innervation, based on cytochemical studies, has also been suggested.¹⁷ Moreover, innervation mediated by various neurotransmitters such as substance P (SP), vasoactive intestinal peptide (VIP), neuropeptide Y, somatostatin, cholecystokinin, and neurotensin has been reported to exist in the human liver.²¹⁻²³ The majority of studies have been conducted in attempts to correlate neurophysiologic data with specific metabolic functions of hepatocytes, glucose metabolism, biliary secretion, and so forth.^{24,25} Studies related to the autonomic control of hemodynamic, baroreceptive, and osmoreceptive functions have been based on macroscopic extrinsic innervation of the liver.^{26,27} Little attention has been paid to the intrinsic innervation and its physiologic implications for cells other than hepatocytes.

BACKGROUND OF LIPOCYTE CONTRACTILITY

In intralobular spaces, nerve bundles are observed along the space of Disse. Nerve endings with vesicles containing

neurotransmitters such as SP or VIP are often visible close to lipocytes.¹⁹ In certain respects, lipocytes around the sinusoids resemble pericytes around a capillary. Consequently, lipocytes have the potential to contract and may be involved in the regulation of hepatic sinusoidal microcirculation.

ET is a peptide composed of 21 amino acids that has been isolated from the supernatant after culture of porcine aortic endothelial cells.²⁸ Of the known vasoactive substances, ET has the most potent vasoconstrictive effects on smooth muscles.²⁸ The physiology of ET is currently the focus of much research. ET has three isoforms (ET-1, ET-2, and ET-3) with different sequences of amino acids.²⁹ There are at least two subtypes of ET receptors^{30,31}: ET_A and ET_B. According to the relative binding affinities and biological activities of the three isopeptides for their receptors, ET-1 is equipotent to ET-2 but more potent than ET-3 for ET_A receptor, whereas ET-1 is equipotent to ET-2 and ET-3 for ET_B receptor.³² Although a third receptor type (ET_C) has been pharmacologically and biochemically characterized, the cDNA encoding of this receptor type has not been identified.^{31,33} Receptors have been found not only in vascular smooth muscle but also in myocardium, brain, kidney, lung, adrenal gland, intestine, and liver.^{34,35} In the rat liver, ¹²⁵I-ET-1 binding sites, which localize ET receptors, are located mainly in the sinusoidal lining cells of hepatic lobules and are weakly expressed in the luminal space of the portal and central veins.^{36,37} In the intralobular spaces, the density is greatest in the periportal region and recedes progressively from the midzonal to the pericentral region (Fig. 5).³⁶ Ultrastructural autoradiographic observation mainly reveals grains in lipocytes, as




FIG. 2. Electron micrograph ($\times 6,500$) showing the portal area in the human liver. Nerve bundles (arrows) closely appose fibroblasts (F) localized around the portal veins (P), hepatic artery (A), and bile duct (B). S, Schwann cell.

well as in endothelial cells of the portal vein, central vein, and hepatic sinusoids, and Kupffer cells.^{36,37} Furuta et al.³⁷ reported that about 35% of all silver grains were located on the lipocytes. Moreover, rat lipocytes express both ET_A and ET_B receptors.³⁸ The mRNA for ET_B receptor, but not for ET_A receptor, is present in sinusoidal endothelial cells and Kupffer cells. The heterogeneity of ET-1-binding sites in hepatic lobules appears to parallel the blood flow pattern in hepatic sinusoids: the portal veins branch out into the hepatic sinusoids (6–7 cm H₂O blood pressure³⁹), and the hepatic arterial branches also drain into the sinusoids either directly (12–15 cm H₂O blood pressure⁴⁰) via the peribiliary plexus or through the portal venules. Thus, stable regulation of blood flow in the hepatic sinusoids is required predominantly in the periportal spaces compared with that in the mid-zonal and centrilobular spaces. According to McCuskey,⁴¹ sinusoidal endothelial cells and Kupffer cells are responsive to a wide variety of pharmacodynamic substances, and by contracting or relaxing, these cells regulate the rate and distribution of blood flow through the sinusoids. Although the participation of lipocytes remains to be clarified, these cells may play a leading role in the regulation of the hepatic sinusoidal microcirculation.

Lipocytes, as well as sinusoidal endothelial cells and Kupffer cells, contain a cytoskeleton that may be related to contractile activity.⁴² After contractile agents such as SP or ET-1 attaches to these receptors, the intracellular Ca^{2+} content increases, and the increased Ca^{2+} binds to calmodulin.⁴³ The Ca^{2+} -calmodulin complex enhances myosin-light chain kinase activity, and its activity affects contractile proteins such as actin and

myosin.⁴⁴ Through these reactions, lipocytes seem to contract. Recently, it was reported that lipocytes contain α -smooth muscle actin, which is closely involved in the contractility of smooth muscle cells.^{45–47}

CONTRACTION OF LIPOCYTES

Lipocytes contract or relax in response to various agents. Kawada et al.⁴⁸ recently reported that ET-1 is a lipocyte contraction agonist that concomitantly stimulates the formation of inositol phosphates and a transient increase in Ca^{2+} content. They also noted that contraction is induced but less markedly by prostaglandin (PG) $F_{2\alpha}$ and a thromboxane A_2 analogue. Since $PGF_{2\alpha}$ is released from lipocytes after stimulation with noradrenaline, lipocytes may contract in response to an autocrine mechanism under the influence of noradrenaline released from adrenergic nerve endings.⁴⁹ We reported the contractile response of cultured rat lipocytes to ET-1 and SP.⁵⁰ Even though lipocytes contract in response to treatment with ET-1 or SP, the extent and onset of the contraction differ in response to the two peptides (Fig. 6). That is, the effect of ET-1 has a stronger and more prolonged effect compared with that of SP. Yanagisawa et al.²⁸ reported that ET-1 induces contraction of vascular smooth muscle cells lasting for more than 1 h. Therefore, ET-1 may play a role in prolonging tension in the hepatic sinusoidal walls.

Pinzani et al.⁵¹ demonstrated that the response of cultured human lipocytes to ET-1 includes an increase in intracellular free calcium content. An increase in Ca^{2+} content of more than four-fold was associated with a simultaneous and transient reduction in cell area. In cultured vascular smooth muscle cells




FIG. 3. Electron micrograph ($\times 11,000$) showing the central vein in the human liver. Enlarged nerve endings (N) contact the surface of myofibroblasts (M). The distal end of the nerve ending is closely associated with the basal lamina (arrows) of the endothelial cell of the central vein (C).

and glomerular mesangial cells, ET-1 activates phospholipase C after binding to its receptors, leading in turn to the formation of inositol triphosphate (IP_3) and diacylglycerol.^{52,53} Thus, the rise in cytosolic Ca^{2+} induced by the peptide is derived mainly from intracellular stores due to the formation of IP_3 . The contraction of lipocytes may be similar to that of vascular smooth muscle cells and glomerular mesangial cells. Zhang et al.⁵⁴ studied sinusoidal and extrasinusoidal constrictor responses of the hepatic microcirculation to ET-1 and ET-3, and the possible roles of lipocytes, Kupffer cells, and sinusoidal endothelial cells in mediating this response using isolated rat livers under high-power intravital microscopy. ET-1 was found to induce significant sinusoid constriction at the sites containing lipocytes but not at those including Kupffer cells or endothelial cells, whereas ET-3 had no effect on sinusoid narrowing. Moreover, they noted that ET-1-induced sinusoid constriction was mediated by ET_A receptor on lipocytes.⁵⁴ Sinusoidal constriction by ET-1 in the perfused liver reaches a maximum in a short time compared with the time required for maximal contraction of cultured lipocytes by ET-1.^{48,50,54} A few factors such as the dose of ET-1 and differences in experimental circumstance between *in vivo* and *in vitro* appear to be related to the discrepancy. That is, since lipocytes *in vivo* are located in three-dimensional spaces such as the space of Disse through limited cell-to-cell contact or cell-to-matrix contact, the cells appear to be able to contract promptly in concert with hepatic sinusoidal endothelial cells in response to lower concentrations of ET-1.⁵⁵

In contrast, PGE_2 , Iloprost, which is a PGI_2 analogue, sodium nitroprusside, and adrenomedullin derived from the hu-

man adrenal medulla promote the relaxation of lipocytes.^{48,56} PGE_2 , Iloprost, and adrenomedullin cause the elevation of cAMP levels in lipocytes, whereas sodium nitroprusside triggers cGMP accumulation. In addition, lipocytes produce nitric oxide (NO) and inhibit contractility by an autocrine mechanism related to NO.^{57,58} In this way, lipocytes appear to be associated with the regulation of hepatic sinusoidal microcirculation by contraction and relaxation. Moreover, hepatic sinusoidal endothelial cells are closely related to lipocytes with respect to hepatic sinusoidal microcirculation. These cells possess a cell body with a nucleus and slender extended processes containing pores (fenestrae) that are arranged in so-called sieve plates and allow direct contact between the plasma and its solutes and cells, such as hepatocytes or lipocytes, which are located behind the endothelial barrier. That is, the fenestrated endothelial lining forms a barrier excluding direct contact of cells in the space of Disse with larger particles such as blood cells or large chylomicrons, and is involved in the transfer of substances such as nutrients between the sinusoidal and the space of Disse by way of the fenestrae.⁵⁹

Fenestrae can be altered by various factors. Enlargement of fenestrae diameter has been reported after alcohol abuse,⁶⁰ sinusoidal pressure,⁶¹ hypoxia, and endotoxins,⁶² while application of serotonin and noradrenalin decreases fenestrae diameter.⁶³ In addition, the relative sizes of the sinusoidal diameter and blood cells flowing into the sinusoids are closely related to the transfer between the sinusoids and the space of Disse. Commonly, a considerable number of blood cells are too large to fit into sinusoids. However, red blood cells pass in a single row and adapt their shapes to various



FIG. 4. Electron micrograph ($\times 15,000$) showing the sinusoidal area in the human liver. Nerve endings (arrows) with many vesicles are observed close to a lipocyte (L) and hepatocyte (H). The nerve ending is associated with the sinusoidal endothelial cell (E) for a short distance. S, sinusoid.




FIG. 5. Light microscopic autoradiographs showing ^{125}I -SP (a) and ^{125}I -ET-1 (b) binding sites in the normal human liver. (a) A concentrated accumulation of ^{125}I -SP grains is observed in lipocytes (arrows) of the perisinusoidal walls. $\times 800$. (b) Many more ^{125}I -ET-1 grains are observed in sinusoidal lining cells of the periportal area as compared with the midzonal or pericentral areas. C, central vein $\times 200$; ET, endothelin; P, portal vein; SP, substance P.

seems to contribute to vascular hyporeactivity in the presence of high doses but not low doses of ET-1.⁷¹

Autonomic hyperactivity with increased cardiac output and other manifestations is observed in patients with cirrhosis,⁷⁷ whereas intralobular innervation is decreased or absent in the cirrhotic liver. However, the nerve fiber density is increased around the portal vein branches and vessels in the fibrous septa.⁷⁹⁻⁸¹

In addition, liver fibrosis is characterized by the accumulation of extracellular matrix components in the space of Disse, forming a so-called "capillarization" of the sinusoids.⁸² At this stage, resistance in the sinusoidal walls is increased, and the hepatic sinusoidal microcirculation may be limited.⁸³ With the

progression of liver fibrosis, intralobular innervation becomes scant, whereas the role of lipocytes in the hepatic sinusoidal microcirculation may become more prominent. As shown in Figure 7, the regulation of blood flow, resistance, pressure, osmolarity and glucose metabolism, and so on in the sinusoids of normal liver is believed to be performed mainly by neuropeptides such as SP or VIP released from nerve endings. However, the regulation of these functions in sinusoids of the cirrhotic liver seems to be regulated by agents such as ET-1 or NO in a paracrine or autocrine manner. Regardless, the relationship between intralobular innervation and the contractility of lipocytes in liver disorders requires further investigation.

REFERENCES

- Ito T, Nemoto M. Über die Kupfferschen Sternzellen und die "Fett-speicherungszellen" ("fat storing cells") in der Blutkapillarenwand der menschlichen Leber. *Okajima Folia Anat Jpn* 1952;24:243
- Wake K. Perisinusoidal stellate cells (fat-storing cells, interstitial cells, lipocytes), their related structure in and around the liver sinusoids, and vitamin A-storing cells in extrahepatic organs. *Int Rev Cytol* 1980;66:303
- Blomhoff R, Wake K. Perisinusoidal stellate cells of the liver: important roles in retinol metabolism and fibrosis. *FASEB J* 1991;5:271
- Ramadori G. The stellate cell (Ito-cell, fat-storing cells, lipocyte, perisinusoidal cell) of the liver. *Virchows Arch B* 1991;61:147
- Fawcett DW. Blood and lymphovascular system. In: Fawcett DW, ed. *A textbook of histology*, 11th ed. Philadelphia: WB Saunders, 1986:367
- Joyce NC, Haire MF, Palade GE. Contractile proteins in pericytes. I. immunoperoxidase localization of tropomyosin. *J Cell Biol* 1985;100:1379
- Yamada E. Some observations on the nerve terminal on the liver parenchymal cell of the mouse as revealed by electron microscopy. *Okajima Folia Anat Jpn* 1965;40:663
- Forssman WG, Ito S. Hepatocyte innervation in primates. *J Cell Biol* 1977;74:299
- Barja F, Mathison R. Adrenergic and peptidergic (substance P and vasoactive intestinal polypeptide) innervation of the rat portal vein. *Blood Vessels* 1982;19:263
- Ohata M. Electron microscope study on the innervation of guinea pig liver. *Arch Histol Jpn* 1984;47:149
- Sasaki Y, Kamada T, Hayashi N, et al. Immunohistochemical distribution of glucagon, substance P and vasoactive intestinal polypeptide in hepatic vasculature of the rat. *Hepatology* 1984;4:1184
- Barja F, Mathison R. Sensory innervation of the rat portal vein and the hepatic artery. *J Auton Nerv Syst* 1984;10:117
- Ito Y, Magari S, Sakanaka M. Immunoelectron-microscopic localization of peptidergic nerve fibers around lymphatic capillaries in the rat liver. *Arch Histol Cytol* 1990;53(suppl):199
- Fehér E, Fodor M, Fehér J. Ultrastructural localization of somatostatin- and substance P-immunoreactive nerve fibers in the feline liver. *Gastroenterology* 1992;102:287
- Ito T, Shibasaki S. Electron microscopic study on the hepatic sinusoidal wall and the fat-storing cells in the normal human liver. *Arch Histol Jap* 1968;29:137
- Nobin A, Falck B, Ingemansson S, Järhult J, Rosengren E. Organization and function of the sympathetic innervation of human liver. *Acta Physiol Scand* 1977;452(suppl):103
- Amenta F, Cavallotti C, Ferrante F, Tonelli F. Cholinergic nerves in the human liver. *Histochem J* 1981;13:419
- Kyösola K, Penttilä O, Ihäniemi T, Varis K, Salaspuro M. Adrenergic innervation of the human liver. A fluorescence histochemical analysis of clinical liver biopsy specimens. *Scand J Gastroenterol* 1985;20:254
- Ueno T, Inuzuka S, Torimura T, et al. Intrinsic innervation of the human liver. *J Clin Electron Microscopy* 1988;21:481
- Bioulac-Sage P, Lafon ME, Saric J, Balabaud C. Nerves and perisinusoidal cells in human liver. *J Hepatol* 1990;10:105
- Ueno T, Inuzuka S, Torimura T, et al. Distribution of substance P and vasoactive intestinal peptide in the human liver: light and electron immunoperoxidase methods of observation. *Am J Gastroenterol* 1991;86:1633
- El-Salhy M, Stenling R, Grimelius L. Peptidergic innervation and endocrine cells in the human liver. *Scand J Gastroenterol* 1993;28:809
- Ding WG, Tooyama I, Kitasato H, Fujimura M, Kimura H. Phylogenetic and ontogenetic study of neuropeptide Y-containing nerves in the liver. *Histochemical J* 1994;26:453
- Shimazu T. Reciprocal innervation of the liver: its significance in metabolic control. *Adv Metab Disord* 1983;10:355
- Lautt WW. Afferent and efferent neural roles in liver function. *Prog Neurobiol* 1983;21:323
- Beverly JL, Yang ZJ, Meguid MM. Hepatic vagotomy effects on metabolic challenges during parenteral nutrition in rats. *Am J Physiol* 1994;266:R646
- Nijima A, Meguid MM. Parenteral nutrients in rat suppresses hepatic vagal afferent signals from portal vein to hypothalamus. *Surgery* 1994;116:294
- Yanagisawa M, Kurihara H, Kimura S, et al. A novel potent vasoconstrictor peptide produced by vascular endothelial cells. *Nature* 1988;332:411
- Inoue A, Yanagisawa M, Kimura S, et al. The human endothelin family: three structurally and pharmacologically distinct isopeptides predicted by three separate genes. *Proc Natl Acad Sci USA* 1989;86:2863
- Arai H, Hori S, Aramori I, Ohkubo H, Nakanishi S. Cloning and expression of a cDNA encoding an endothelin receptor. *Nature* 1990;348:730
- Sakurai T, Yanagisawa M, Takuwa Y, et al. Cloning of a cDNA encoding non-isopeptide-selective subtype of the endothelin receptor. *Nature* 1990;348:732
- Sakurai T, Yanagisawa M, Masaki T. Molecular characterization of endothelin receptors. *Trends Pharmacol Sci* 1992;13:103
- Eguchi S, Kozuka M, Hirose S, et al. Identification of G protein-coupled endothelin receptors in cultured bovine endothelial cells. *Biochem Biophys Res Commun* 1991;174:1343
- Battistini B, D'Orléans-Juste P, Sirois P. Endothelins: circulating plasma levels and presence in other biologic fluids. *Lab Invest* 1993;68:600
- Levin ER. Endothelins. *New Engl J Med* 1995;333:356
- Gondo K, Ueno T, Sakamoto M, Sakisaka S, Sata M, Tanikawa K. The endothelin-1 binding site in rat liver tissue: light- and electron-microscopic autoradiographic studies. *Gastroenterology* 1993;104:1745
- Furuya S, Naruse S, Nakayama T, Nokihiro K. Binding of ¹²⁵I-endothelin-1 to fat-storing cells in rat liver revealed by electron microscopic radioautography. *Anat Embryol* 1992;185:97
- Housset C, Rockey DC, Bissell DM. Endothelin receptors in rat liver: lipocytes as a contractile target for endothelin 1. *Proc Natl Acad Sci USA* 1993;90:9266
- Nakata K. Microcirculation and hemodynamical analysis of the blood circulation in the liver. *Acta Path Jap* 1967;17:361
- Rappaport AM. Microcirculatory units in the mammalian liver. *Bibl Anat* 1977;16:116
- McCuskey RS. A dynamic and static study of hepatic arterioles and hepatic sphincters. *Am J Anat* 1966;119:455
- Ballardini G, Fallani M, Biagini G, Bianchi FB, Pisi E. Desmin

- and actin in the identification of Ito cells and in monitoring their evolution to myofibroblasts in experimental liver fibrosis. *Virchows Arch [B]* 1988;56:45
43. Kakiuchi S, Yamazaki R. Calcium dependent phosphodiesterase activity and its activating factor (PAF) from brain. *Biochem Biophys Res Commun* 1970;41:1104
 44. Kamm KE, Stull JT. The function of myosin and myosin light chain kinase phosphorylation in smooth muscle. *Ann Rev Pharmacol Toxicol* 1985;25:593
 45. Ramadori G, Veit T, Schwöglar S, et al. Expression of the gene of the α -smooth muscle-actin isoform in rat liver and in rat fat-storing (ITO) cells. *Virchows Arch [B]* 1990;59:349
 46. Schmitt-Gräff A, Krüger S, Bochar F, Gabbiani G, Denk H. Modulation of alpha smooth muscle actin and desmin expression in perisinusoidal cells of normal and diseased human livers. *Am J Pathol* 1991;138:1233
 47. Rockey DC, Boyles JK, Gabbiani G, Friedman SL. Rat hepatic lipocytes express smooth muscle actin upon activation in vivo and in culture. *J Submicrosc Cytol Pathol* 1992;24:193
 48. Kawada N, Tran-thi TA, Klein H, Decker K. The contraction of hepatic stellate (Ito) cells stimulated with vasoactive substances. *Eur J Biochem* 1993;213:815
 49. Athari A, Hänecke K, Jungermann K. Prostaglandin $F_{2\alpha}$ and D_2 release from primary Ito cell cultures after stimulation with nor-adrenaline and ATP but not adenosine. *Hepatology* 1994;20:142
 50. Sakamoto M, Ueno T, Kin M, et al. Ito cell contraction in response to endothelin-1 and substance P. *Hepatology* 1993;18:978
 51. Pinzani M, Failli P, Ruocco C, et al. Fat-storing cells as liver-specific pericytes. *J Clin Invest* 1992;90:642
 52. Marsden PA, Danthuluri NR, Brenner BM, Ballermann BJ, Brock TA. Endothelin action on vascular smooth muscle involves inositol triphosphate and calcium mobilization. *Biochem Biophys Res Commun* 1989;158:86
 53. Simonson MS, Wann S, Mené P, et al. Endothelin stimulates phospholipase C, Na^+/H^+ exchange, c-fos expression, and mitogenesis in rat mesangial cells. *J Clin Invest* 1989;83:708
 54. Zhang JX, Bauer M, Clemens MG. Vessel- and target cell-specific actions of endothelin-1 and endothelin-3 in rat liver. *Am J Physiol* 1995;269:G269
 55. Bauer M, Zhang JX, Bauer I, Clemens MG. ET-1 induced alterations of hepatic microcirculation: sinusoidal and extrasinusoidal sites of action. *Am J Physiol* 1994;267:G143
 56. Kawada N, Inoue M. Effect of adrenomedullin on hepatic pericytes (stellate cells) of the rat. *FEBS Letters* 1994;356:109
 57. Helyar L, Bundschuh DS, Laskin JD, Laskin DL. Induction of hepatic Ito cell nitric oxide production after acute endotoxemia. *Hepatology* 1994;20:1509
 58. Rockey DC, Chung JJ. Inducible nitric oxide synthase in rat hepatic lipocytes and the effect of nitric oxide on lipocyte contractility. *J Clin Invest* 1995;95:1199
 59. Wisse E, Zanker RB, Charels K, Van Der Smissen P, McCuskey RS. The liver sieve: considerations concerning the structure and function of endothelial fenestrae, the sinusoidal wall and the space of Disse. *Hepatology* 1985;5:683
 60. Mak KM, Lieber CS. Alterations in endothelial fenestrations in liver sinusoids of baboons fed alcohol: a scanning electron microscopic study. *Hepatology* 1984;4:386
 61. Fraser R, Bowler LM, Day WA, et al. High perfusion pressure damages the sieving ability of sinusoidal endothelium in rat livers. *Br J Exp Pathol* 1980;61:222
 62. Frenzel H, Kremer B, Hücker H. The liver sinusoids under various pathological conditions. A TEM and SEM study of rat liver after respiratory hypoxia, telecobalt-irradiation and endotoxin application. In: Wisse E, Knook DL, eds. *Kupffer cells and other liver sinusoidal cells*. Amsterdam: Elsevier, 1977:213
 63. Oda M, Nakamura M, Watanabe N, et al. Some dynamic aspects of the hepatic microcirculation—demonstration of sinusoidal endothelial fenestrae as a possible regulatory factor. In: Tsuchiya M, et al., eds. *Intravital observation of organ microcirculation*. The Hague: Excerpta Medica Foundation, 1983:105
 64. Kent G, Gay S, Inoue T, et al. Vitamin A-containing lipocytes and formation of type III collagen in liver injury. *Proc Natl Acad Sci USA* 1976;73:3719
 65. Inuzuka S, Ueno T, Torimura T, et al. Immunohistochemistry of the hepatic extracellular matrix in acute viral hepatitis. *Hepatology* 1990;12:249
 66. Yamaoka K, Nouchi T, Marumo F, Sato C. α -smooth-muscle actin expression in normal and fibrotic human livers. *Dig Dis Sci* 1993;38:1473
 67. Davis BH, Vucic A. The effect of retinol on Ito cell proliferation in vitro. *Hepatology* 1988;8:788
 68. Matsumoto H, Uemasu J, Kitano M, Kawasaki H. Clinical significance of plasma endothelin-1 in patients with chronic liver disease. *Dig Dis Sci* 1994;39:2665
 69. Kawamura E, Yamanaka N, Okamoto E, Tomoda F, Furukawa K. Response of plasma and tissue endothelin-1 to liver ischemia and its implication in ischemia-reperfusion injury. *Hepatology* 1995;21:1138
 70. Guarner C, Soriano G, Tomas A, et al. Increased serum nitrite and nitrate levels in patients with cirrhosis: relationship to endotoxemia. *Hepatology* 1993;18:1139
 71. Clària J, Jiménez W, Ros J, et al. Increased nitric oxide-dependent vasorelaxation in aortic rings of cirrhotic rats with ascites. *Hepatology* 1994;20:1615
 72. Laskin DL, Del Valle MR, Heck DE, et al. Hepatic nitric oxide production following acute endotoxemia in rats is mediated by increased inducible nitric oxide synthase gene expression. *Hepatology* 1995;22:223
 73. Bird G, Severn A, Danesh BJ, Liew FY. Nitric oxide in alcohol-induced liver failure. *J Hepatol* 1993;19:316
 74. Harbrecht BG, Di Silvio M, Demetris AJ, Simmons RL, Billiar TR. Tumor necrosis factor- α regulates in vivo nitric oxide synthesis and induces liver injury during endotoxemia. *Hepatology* 1994;20:1055
 75. Lopez-Talavera JC, Merrill WW, Groszmann RJ. Tumor necrosis factor α : a major contributor to the hyperdynamic circulation in prehepatic portal-hypertensive rat. *Gastroenterology* 1995;108:761
 76. Oshita M, Takei Y, Kawano S, et al. Roles of endothelin-1 and nitric oxide in the mechanism for ethanol-induced vasoconstriction in rat liver. *J Clin Invest* 1993;91:1337
 77. Bomzon A, Blendis LM. The nitric oxide hypothesis and the hyperdynamic circulation in cirrhosis. *Hepatology* 1994;20:1343
 78. Hartleb M, Moreau R, Cailmail S, Gaudin C, Lebre C. Vascular hyporesponsiveness to endothelin I in rats with cirrhosis. *Gastroenterology* 1994;107:1085
 79. Miyazawa Y, Fukuda Y, Imoto M, Koyama Y, Nagura H. Immunohistochemical studies on the distribution of nerve fibers in chronic liver diseases. *Am J Gastroenterol* 1988;83:1108
 80. Lee JA, Ahmed Q, Hines JE, Burt AD. Disappearance of hepatic parenchymal nerves in human liver cirrhosis. *Gut* 1992;33:87
 81. Jaskiewicz K, Voigt MD, Robson SC. Distribution of hepatic nerve fibers in liver diseases. *Digestion* 1994;55:247
 82. Ueno T, Inuzuka S, Torimura T, et al. Serum hyaluronate reflects hepatic sinusoidal capillarization. *Gastroenterology* 1993;105:475
 83. Iwao T, Toyonaga A, Sumino M, et al. Portal hypertensive gastropathy in patients with cirrhosis. *Gastroenterology* 1992;102:2060