

tion on

se is known to  
ease process, as  
within the sys-

fusion, operates  
orce to bile for-  
bile acids and  
ed water flow

Reichen & Si-  
ation or extra-  
e flow, but it is  
tion *per se* in-  
ore, whether in-  
definition of a  
kely that biliary  
ar change with  
arance.

and

many drugs is  
7; Tillement et  
relates with the  
ations of albu-  
e 2 main drug  
er et al. 1988;  
vidence suggests  
rotein for drugs  
ng a qualitative  
al. 1988). This  
studies which  
titative changes  
of the glycopro-  
Pedersen et al.  
ation of hepatic  
have been ex-

7).

pharmacokinetic

renal impair-  
ment, i.e. when  
vidence of other

known causes of renal failure are absent (Papper 1983).

This is normally thought of as a problem only in severe and rapidly advancing liver disease. However, phenazone and aminophenazone (aminopyrine) clearance, used to indicate the severity of liver disease, has been shown to correlate with measures of both proximal and distal tubular Na<sup>+</sup> handling in patients with chronic stable cirrhosis (Wood et al. 1988) and cirrhotic rats (Wensing et al. 1990b). While these changes may reflect an alteration in the handling mechanisms peculiar to Na<sup>+</sup> balance, there remains a possibility of generalised change in tubular function. If this were so, xenobiotic clearance by the kidney which is dependent on tubular mechanisms could be altered in patients with even moderate degrees of liver impairment.

#### 4. Methods and Observations with Drugs and Markers of Drug Disposition

The majority of published clinical pharmacokinetic studies in liver disease have used a relatively simple mechanistic framework to interpret their experimental findings. Thus, changes in the systemic clearance or systemic availability of orally administered drugs have been ascribed to a reduction in hepatic intrinsic clearance and/or effective hepatic blood flow. This approach was the basis for the intact hepatocyte and sick cell hypotheses (Branch 1982; Branch & Shand 1976). With our increasing understanding of the pathological changes and of the mechanisms by which these changes may alter drug disposition in liver disease, it is now possible to interpret experimental findings using a broader mechanistic framework. This section examines the results of some selected studies of drugs and other markers of drug disposition in terms of the mechanisms of altered drug disposition described in section 3. Blaschke (1977) classified the hepatic elimination of drugs according to whether hepatic blood flow, intrinsic clearance or unbound fraction was rate limiting; the same classification is used here, and the review concentrates on more recently published studies.

#### 4.1 Capacity-Limited Binding-Insensitive Hepatic Elimination

Drugs in this category have a low hepatic intrinsic clearance relative to hepatic blood flow and are less than 30% bound to plasma proteins (Blaschke 1977). The hepatic clearance of such drugs is low and will be unaffected by changes in plasma protein binding or hepatic blood flow. Therefore, the interpretation of findings from studies of the disposition of these drugs in liver disease should be more straightforward than that for drugs with flow- or protein binding-limited hepatic clearance. The hepatic clearance of drugs with capacity-limited binding-insensitive hepatic elimination may still be affected in liver disease by loss of cell mass, reduction in enzyme mass per cell, alteration in the intrahepatic distribution of enzymes or sinusoidal capillarisation. Findings with typical drugs in this category are discussed below.

##### 4.1.1 Phenazone

Phenazone is one of the standard model compounds for studying hepatic oxidative drug metabolism (Vesell 1979; Vesell & Page 1968) and has found widespread use in the study of the effect of liver disease on hepatic drug disposition. The clearance of phenazone following oral administration (and assuming complete bioavailability) has been measured in patients with a wide variety of hepatic disorders, but a reduction in clearance is usually observed only in severe disease. For example, there is little impairment in chronic persistent hepatitis, chronic active hepatitis, mild cirrhosis or hepatosplenic schistosomiasis (Daneshmend et al. 1982; Horvath et al. 1986; Villeneuve et al. 1987). However, phenazone clearance is reduced by 50 to 70% in advanced or decompensated cirrhosis (Daneshmend et al. 1982; Kawasaki et al. 1988; Kirch et al. 1989; McQuinn et al. 1988; Mehta et al. 1986; Pentikainen et al. 1986, 1989; Villeneuve et al. 1987) and in certain other severe disorders such as acute episodes of hereditary hepatic porphyria (Birnie et al. 1987).

In many of the studies cited above, phenazone was administered concurrently with another drug

metabolism in liver  
23-432, 1979  
interactions in cirrho-  
8, 1986  
function. Hepatology  
E. Intravenous admin-  
chronic liver disease.  
position in chronic liver  
ical Pharmacokinetics  
PV, Peters M, et al. A  
e hepatic function in  
DS. Influence of liver  
hepatic monooxygenase  
Clinical Pharmacology  
iver function. Medical  
1979  
c drug oxidation. Im-  
s. Clinical Pharmacol-  
von Mollendorff E.  
rmacokinetics of toral  
dose in patients with  
of the liver. Arzneim-  
nier-Layrargues G. Af-  
H] muscimol (GABA-  
iazepine receptors are  
cirrhotic patients with  
1084-1088, 1988  
Lean AJ. Variation in  
rug fraction: discrimi-  
limination. Journal of  
985  
ura L, Sherlock S. The  
ment of hepatic blood  
Clinical Science 21: 43-  
ulation. In Arias et al.  
y, pp. 627-646, Raven  
C. Downregulation of  
teroid 16 $\alpha$ -hydroxylose,  
rtal bypass. Relevance  
d drug metabolism in  
Investigation 83: 1211-  
ships among beta-adra-  
tion, and liver func-  
ndinavian Journal of  
patients with Laennec's  
y. European Journal of  
33  
dson CS. Indocyanine  
erties, plasma decay,  
ical Investigation 39:

- Daneshmend TK, Homeida M, Kaye CM, Elamin AA, Roberts CJC. Disposition of oral metronidazole in hepatic cirrhosis and in hepatosplenic schistosomiasis in hepatic cirrhosis and in hepatosplenic schistosomiasis. *Gut* 23: 807-813, 1982
- Dao MT, Villeneuve J-P. Kinetics and dynamics of triamterene at steady-state in patients with cirrhosis. *Clinical and Investigative Medicine* 11: 6-9, 1988
- Davey PG. Pharmacokinetics in liver disease. *Journal of Antimicrobial Chemotherapy* 21: 1-8, 1988
- Desmond PV, Patwardhan RV, Johnson RF, Schenker S. Impaired elimination of caffeine in cirrhosis. *Digestive Diseases and Science* 25: 193-197, 1980
- Dunn MA, Kamal R. Hepatic schistosomiasis. *Hepatology* 1: 653-661, 1981
- Du Souich P, Erill S. Metabolism of procainamide and p-aminobenzoic acid in patients with chronic liver disease. *Clinical Pharmacology and Therapeutics* 22: 588-595, 1977
- Echizen H, Ishizaki T. Superiority of disease-specific over conventional formula in predicting creatinine clearance from serum creatinine in patients with liver cirrhosis. *Therapeutic Drug Monitoring* 10: 369-375, 1988
- Erlinger S. What is cholestasis in 1985? *Journal of Hepatology* 1: 687-693, 1985
- Farrell GC, Cooksley WGE, Powell LW. Drug metabolism in liver disease: activity of hepatic microsomal metabolising enzymes. *Clinical Pharmacology and Therapeutics* 26: 483-492, 1979
- Farrell GC, Koltai A, Murray M. Source of raised serum estrogens in male rats with portal bypass. *Journal of Clinical Investigation* 81: 221-228, 1988
- Farrell GC, Zaluzny L. Portal vein ligation selectively lowers hepatic cytochrome P450 levels in rats. *Gastroenterology* 85: 275-282, 1983
- Forrest JAH, Andriaenssens P, Finlayson NDC, Prescott LF. Paracetamol metabolism in chronic liver disease. *European Journal of Clinical Pharmacology* 15: 427-431, 1979
- Forrest JAH, Finlayson NDC, Adjepon-Yamoah KK, Prescott LF. Antipyrine, paracetamol and lignocaine elimination in chronic liver disease. *British Medical Journal* 1: 1384-1387, 1977
- Frost RW, Lettieri JT, Krol G, Shambles EC, Lessner KL. The effect of cirrhosis on the steady-state pharmacokinetics of oral ciprofloxacin. *Clinical Pharmacology and Therapeutics* 45: 608-616, 1989
- Fuller R, Hoppel C, Ingalls ST. Furosemide kinetics in patients with hepatic cirrhosis with ascites. *Clinical Pharmacology and Therapeutics* 30: 461-467, 1981
- Gengo FM, Fagan SC, Krol G, Bernhard H. Nimodipine disposition and haemodynamic effects in patients with cirrhosis and age-matched controls. *British Journal of Clinical Pharmacology* 23: 46-53, 1987
- Gerbes AL, Remien J, Jungst D, Sauerbruch Y, Paumgartner G. Evidence for down-regulation of beta-2-adrenoceptors in cirrhotic patients with severe ascites. *Lancet* 1: 1409-1411, 1986
- Ghabrial H, Desmond PV, Watson KJR, Gijssbers AJ, Horman PJ, et al. The effects of age and chronic liver disease on the elimination of temazepam. *European Journal of Clinical Pharmacology* 30: 93-97, 1986
- Giacomini KM, Giacomini JC, Gibson TP, Levy G. Propoxyphene and norpropoxyphene plasma concentrations after oral propoxyphene in cirrhotic patients with and without surgically constructed portacaval shunt. *Clinical Pharmacology and Therapeutics* 28: 417-424, 1980
- Gonzalez FJ. Molecular genetics of the P-450 superfamily. *Pharmacology and Therapeutics* 45: 1-38, 1990
- Gonzalez G, Aransibia A, Rivas MI, Caro P, Antezana C. Pharmacokinetics of frusemide in patients with hepatic cirrhosis. *European Journal of Clinical Pharmacology* 22: 315-320, 1982
- Goresky CA. A linear method for determining liver sinusoidal and extracellular volume. *American Journal of Physiology* 204: 626-640, 1963
- Goresky CA, Bach GG, Nadeau BE. On the uptake of material by the intact liver: the transport and net removal of galactose. *Journal of Clinical Investigation* 52: 991-1009, 1973
- Goresky CA, Ziegler WH, Bach GG. Capillary exchange modelling. Barrier-limited and flow-limited distribution. *Circulation Research* 27: 739-764, 1970
- Grahnen A, Jameson S, Loof L, Tyllstrom J, Lindstrom B. Pharmacokinetics of cimetidine in advanced cirrhosis. *European Journal of Clinical Pharmacology* 26: 347-355, 1984
- Greenway CV, Stark RD. Hepatic vascular bed. *Physiological Reviews* 51: 23-65, 1971
- Gross JB, Reichen J, Zeltner TB, Zimmermann A. The evolution of changes in quantitative liver function tests in a rat model of biliary cirrhosis: correlation with morphometric measurement of hepatocyte mass. *Hepatology* 7: 457-463, 1987
- Groszmann R, Kotelanski B, Cohn JN, Khatri IM. Quantitation of portasystemic shunting from the splenic and mesenteric beds in alcoholic liver disease. *American Journal of Medicine* 53: 715-722, 1972
- Groszmann RJ, Kravetz O, Paryzow O. Arteriovenous (AV) shunting in the liver. *Gastroenterology* 70: 983, 1976
- Groszmann RJ, Kravetz O, Paryzow O. Intrahepatic arteriovenous shunting in cirrhosis of the liver. *Gastroenterology* 73: 201-204, 1977
- Guechot J, Loric S, Vaubourdolle M, Chretien Y, Giboudeau J, et al. Effect of protein binding on testosterone extraction by human cirrhotic liver: evidence for a dissociation-limited uptake. *Journal of Clinical Endocrinology and Metabolism* 69: 200-203, 1989
- Guengerich FP. Characterization of human microsomal cytochrome P-450 enzymes. *Annual Review of Pharmacology and Toxicology* 29: 241-264, 1989
- Gugler R, Muller-Liebenau B, Somogyi A. Altered disposition of cimetidine in liver cirrhotic patients. *British Journal of Clinical Pharmacology* 14: 421-429, 1982
- Hasselstrom J, Eriksson S, Persson A, Rane A, Svensson JO, et al. The metabolism and bioavailability of morphine in patients with severe liver cirrhosis. *British Journal of Clinical Pharmacology* 29: 289-297, 1990
- Heinzow B, Corbett H, Constantinides S, Bourne R, McLean AJ. Interaction between oral hydralazine and propranolol I. Changes in absorption, presystemic clearance and splanchnic blood flow. *Journal of Pharmacology and Experimental Therapeutics* 229: 509-514, 1984
- Henderson JM, Kutner MH, Bain RP. First-order clearance of plasma galactose: the effect of liver disease. *Gastroenterology* 83: 1090-1096, 1982
- Hepner GW, Vesell ES. Assessment of aminopyrine metabolism in man after administration of 14C-aminopyrine. Effects of phenobarbital, disulfiram and portal cirrhosis. *New England Journal of Medicine* 291: 1384-1388, 1974
- Hepner GW, Vesell ES, Lipton A, Harvey HA, Wilkinson GR, et al. Disposition of aminopyrine, antipyrine, diazepam and indocyanine green in patients with liver disease or on anti-convulsant drug therapy: diazepam breath test and correlations in drug elimination. *Journal of Laboratory and Clinical Medicine* 90: 440-456, 1977
- Hoefs J, Sakimura I, Reynolds T. Direct measurement of intrahepatic shunting by the portal vein injection of microspheres. *Gastroenterology* 75: 968, 1978
- Holazo AO, Chen SS, McMahon FG, Ryan JR, Konikoff JJ, et al. The influence of liver dysfunction on the pharmacokinetics of carprofen. *Journal of Clinical Pharmacology* 25: 109-114, 1985
- Holstege A, Staiger M, Haag K, Gerok W. Correlation of caffeine elimination and Child's classification in liver cirrhosis. *Klinische Wochenschrift* 67: 6-15, 1989
- Homeida MMA, Ali HM, Arbab BMO, Harron DWG. Propran-