

We thank the Special Trustees of Guy's Hospital and Roussel UCLAF for financial support and Mrs Ann Hollington for help in preparing the typescript.

Correspondence should be addressed to R. H. D., Gastroenterology Unit, 18th floor, Guy's Tower, Guy's Hospital, London SE1 9RT.

REFERENCES

- Ruppin DC, Dowling RH. Is recurrence inevitable after gallstone dissolution by bile acid treatment? *Lancet* 1982; i: 181-85.
- Villanova N, Bazzoli F, Frabboni R, et al. Gallstone recurrence after successful oral bile acid treatment: a follow-up study and evaluation of long term post-dissolution treatment. *Gastroenterology*, 1987; 92: 1789 (abstr).
- Lanzini A, Jazrawi RP, Kupfer RM, Maudgal DP, Joseph AEA, Northfield TC. Gallstone recurrence after medical dissolution. An overestimated threat? *J Hepatol* 1986; 3: 241-46.
- O'Donnell LDJ, Heaton KW. Recurrence and re-recurrence of gallstones after medical dissolution: longterm follow up. *Gut* 1988; 29: 655-58.
- Lee SP, Carey MC, LaMont JT. Aspirin prevention of cholesterol gallstone formation in prairie dogs. *Science* 1981; 211: 1429-30.
- LaMont JT, Turner BS, Di Benedetto D, Handin R, Schafer AI. Arachidonic acid stimulates mucin secretion in prairie dog gallbladder. *Am J Physiol* 1983; 245: G92-G98.
- Hood K, Gleeson D, Ruppin DC, Dowling RH, and the BBGSG: The British/Belgian Gallstone Study Group's post-dissolution trial. *Gut* 1987; 28: A1359 (abstr).
- Hood K, Gleeson D, Ruppin DC, Dowling RH, and the British/Belgian Gallstone Study Group. Can gallstone recurrence be prevented? The British-Belgian postdissolution trial. *Gastroenterology* 1988; 94: A548 (abstr).
- Cantafora A, DiBiase A, Alvaro D, Angelico M, Marin M, Attili AF. High performance liquid chromatographic analysis of molecular species of phosphatidylcholine-development of quantitative assay and its application to human bile. *Clin Chim Acta* 1983; 134: 281-95.
- van Berge Henegouwen GP, van der Werf SDJ, Ruben AT. Fatty acid composition of phospholipids in bile in man: promoting effect of deoxycholate on arachidonate. *Clin Chim Acta* 1987; 165: 27-37.
- van der Werf SDJ, van Berge Henegouwen GP, Ruben A. Th, Palsma DMH. Biliary lipids, bile acid metabolism, gallbladder motor function and small intestinal transit during ingestion of a sub-fifty oral contraceptive. *J Hepatol* 1987; 4: 318-26.
- Lee SP, et al. In: Abstracts of International meeting on Pathochemistry, Pathophysiology and Pathomechanics of the Biliary System, Bologna, March, 1988.
- Carey MC. In: Paumgartner G, Stiehl A, Gerok W, eds. Trends in bile acid research. Lancaster: Kluwer (in press).
- Fisher RS, Stelzer F, Rock E, Malmud LS. Abnormal gallbladder emptying in patients with gallstones. *Dig Dis Sci* 1982; 27: 1019-24.
- Forgacs IC, Maisey MN, Murphy GM, Dowling RH. Influence of gallstones and ursodeoxycholic acid therapy on gallbladder emptying. *Gastroenterology*, 1984; 87: 299-307.
- Pomerantz IS, Shaffer EA. Abnormal gallbladder emptying in a subgroup of patients with gallstones. *Gastroenterology* 1985; 88: 787-91.
- Dowling RH, Gleeson DC, Hood KA, Ruppin DC, and the British/Belgian Gallstone Study Group. Gallstone recurrence and postdissolution management. In: Paumgartner G, Suchl A, Gerok W, eds. Bile acids and the liver. Lancaster: MTP, 1987 355-67.
- Brotschi EA, Lamorte WW, Williams LF, Jr Effect of dietary cholesterol and indomethacin on cholelithiasis and gallbladder motility in guinea pigs. *Dig Dis Sci*, 1984; 29: 1050-56.
- Mok HYI, Bell GD, Dowling RH. Effect of different doses of chenodeoxycholic acid on bile lipid composition and on frequency of side-effects in patients with gallstones. *Lancet* 1974; ii: 253-57.
- Thistle JL, Yu PYS, Hofmann AF, Ott BJ. Prompt return of bile to supersaturated state followed by gallstone recurrence after discontinuance of chenodeoxycholic acid therapy. *Gastroenterology* 1974; 66: 789 (abstr).
- Iser JH, Murphy GM, Dowling RH. Speed of change in biliary lipids and bile acids with chenodeoxycholic acid—is intermittent therapy feasible? *Gut* 1977; 18: 7-15.
- Pomare EW, Heaton KW, Low-Beer TS, Espiner HJ. The effect of wheat bran upon bile salt metabolism and upon the lipid composition of bile in gallstone patients. *Am J Dig Dis* 1976; 21: 521-26.
- Thomton JR, Emmett PM, Heaton KW. Diet and gallstones: effects of refined and unrefined carbohydrate diets on bile cholesterol saturation and bile acid metabolism. *Gut* 1983; 24: 2-6.
- Mok HYI, Grundy SM. Cholesterol and bile acid absorption during bile acid therapy in obese subjects undergoing weight reduction. *Gastroenterology* 1980; 78: 62-67.
- Bennion LJ, Grundy SM. Effects of obesity and caloric intake on biliary lipid metabolism in man. *J Clin Invest* 1975; 56: 996-1011.
- Broomfield P, Chopra R, Sheinbaum R, et al. Formation and prevention of lithogenic bile and gallstones during weight loss. *Gastroenterology* 1987; 92: 1721 (abstr).
- Sampliner RE, Bennett PH, Commess LJ, Rose FA, Burch TA. Gallbladder disease in Pima Indians. Demonstration of high prevalence and early onset by cholecystography. *N Engl J Med* 1970; 283: 1358-64.
- Heaton KW, Read AE. Gallstones in patients with disorders of the terminal ileum and disturbed bile salt metabolism. *Br Med J* 1969; iii: 494-96.
- Bouchier IAD. Postmortem study of the frequency of gallstones in patients with cirrhosis of the liver. *Gut* 1969; 10: 705-10.
- Nicholas P, Rinaudo PA, Conn HO. Increased incidence of cholelithiasis in Laennec's cirrhosis: a postmortem evaluation of pathogenesis. *Gastroenterology* 1972; 63: 112-21.
- Second International Study of Infarct Survival Collaborative Group. Randomised trial of intravenous streptokinase, oral aspirin, both, or neither among 17 187 cases of suspected acute myocardial infarction: ISIS-2. *Lancet* 1988; ii: 349-60.

Hypothesis

DEFENESTRATION OF HEPATIC SINUSOIDS AS A CAUSE OF HYPERLIPOPROTEINAEMIA IN ALCOHOLICS

SONYA A. CLARK
H. BRAMWELL COOK
ROBERT B. G. OXNER

HELEN B. ANGUS
PETER M. GEORGE
ROBIN FRASER

Departments of Pathology and Gastroenterology, Christchurch Hospital; and School of Medicine, Christchurch, New Zealand

Summary The hepatic sinusoidal endothelium separates sinusoidal blood from hepatocytes; changes in the porosity of this endothelium may affect the passage of chylomicrons into hepatocytes and influence lipid metabolism. Chronic exposure to ethanol reduces the porosity of the endothelium; this mechanism may underlie the hyperlipoproteinaemia observed in some people who drink heavily.

THE LIVER SIEVE

A POROUS sinusoidal endothelium with multiple open fenestrae and no basement membrane was found in animal livers by electron microscopy after perfusion-fixation,^{1,2} and shown in rats to separate large circulating chylomicrons from smaller chylomicron remnants.^{3,4} This porous barrier, or "liver sieve", separates the sinusoidal blood from the space of Disse and the hepatocytes—a strategic position for regulation of chylomicron catabolism and lipoprotein metabolism.^{5,6} Ultrastructural evidence for its function as a sieve^{3,4} has been supported by animal studies of the perfusion of radiolabelled chylomicrons.³ The selectivity of the sieve is decreased when the fenestrae are dilated, as in short-term ethanol ingestion,⁷ and abolished when the endothelium has been destroyed by excessive perfusion pressure.⁸ These changes may allow chylomicrons with an excess of dietary fat to reach the hepatocytes, and thereby affect hepatic steatosis.^{7,8} Various animal species have been shown to have open fenestrae lining the sinusoids.⁶ Large and small polystyrene spheres are also sieved in sheep: only spheres of less than 50 nm diameter are allowed to enter the space of Disse and to come into contact with hepatocytes.⁹ Rabbits and chickens have a sinusoidal endothelium only about half as porous as that of rats.^{10,11} This low porosity may accentuate the susceptibility of these species to dietary cholesterol by hindering the passage of cholesterol-rich chylomicron remnants from the blood to the hepatocytes for recognition and removal from circulation.¹²

Nicotine renders the rat susceptible to dietary cholesterol and hypercholesterolaemia by constriction of the endothelial fenestrae, causing a reduction of sinusoidal porosity.¹³ Adrenaline and noradrenaline have similar effect—a mechanism by which stress could influence atherogenesis.¹⁴ Pantethine lowers cholesterol levels in the rabbit, and increases hepatic sinusoidal porosity by dilation of the fenestrae.¹⁵ In tissue culture, the fenestrae of sinusoidal endothelial cells have also been shown to be sensitive to drugs which act by altering the cytoskeleton.¹⁶⁻¹⁸ In rats, ethanol at first dilates the fenestrae,⁷ but prolonged ingestion diminishes their number.¹⁹ In baboons, chronic ethanol

ingestion also leads to a reduction in the density of sinusoidal fenestrae.²⁰ Human beings who drink heavily, even those without cirrhosis, have hepatic sinusoids that resemble capillaries, with a defenestration of the endothelium and the appearance of a basement membrane.²¹

NEW FINDINGS IN MAN

We have examined human liver sinusoids by perfusion-fixation of small portions of diagnostic needle biopsy specimens wedged in a tapered capillary tube.^{22,23} Several specimens were taken from patients with no history of ethanol abuse, and transmission and scanning electron microscopy showed these patients to have the usual fenestrated endothelium with a structure similar to that found in animals (fig 1). A specimen from a 33-year-old publican, with a 10-year history of heavy alcohol intake, was also examined. On light microscopy, his liver exhibited only slight and patchy steatosis and a moderate increase in perisinusoidal fibrosis, but no cirrhosis. Transmission electron microscopy showed collagen and some basement membrane in the space of Disse, which separated the sinusoidal cells from the microvilli of the hepatocytes (fig 2). Scanning electron microscopy showed the endothelium to have a low density of fenestrae (fig 3). This patient also had a severe type V hyperlipoproteinaemia, with chylomicron and very low density lipoprotein (VLDL) triglycerides of up to 55 mmol/l and cholesterol of 16 mmol/l in the $d < 1.006$ kg/l ultracentrifugation fraction. The cholesterol in the high density lipoprotein fraction was also raised (2.2 mmol/l).

BLOOD LIPIDS IN ALCOHOLICS

Increased blood lipids are found in alcoholics,²⁴ approximately 25% of whom have hyperlipidaemia.²⁵ Some alcoholics have hyperlipoproteinaemia, mostly type IV or V.^{26,27} Type V hyperlipoproteinaemia is characterised by an accumulation of chylomicrons and VLDL.²⁵ The combination of alcohol and dietary fat can lead to hypertriglyceridaemia because of a decrease in the activity of lipoprotein lipase and/or apo C-II, a co-factor for lipoprotein lipase.²⁸⁻³⁰ In our patient the levels of both post-heparin lipolytic activity and apo C-II were within normal limits. Mendelson and Mello suggested that the triglyceride response in alcoholics is related to the presence of a pre-existing lipoprotein disorder.³¹ Alternatively, ethanol may alter the level of triglyceride-rich lipoproteins by modifying the absorption of ingested lipids, the hepatic production of lipids, or their rate of removal from circulation. Early workers suggested that the hypertriglyceridaemia of excess alcohol intake was mainly due to increased hepatic secretion of lipoproteins, but they also noted raised circulatory chylomicrons.²⁷ Many reports of alcoholic hypertriglyceridaemia describe chylomicron-like bands,³² as well as pre- β bands, on electrophoresis.^{31,32} One group reported that the excess triglyceride induced by alcohol in the absence of dietary fat was transported as a pre- β -lipoprotein but that, when fat was ingested, alimentary particles accumulated as well. In the pre- β -triglycerides, fatty acids provided by diet were nine times more common than those from de novo hepatic lipogenesis;²⁸ clearance of triglyceride-rich chylomicrons and VLDL may be slowed by a competition for common disposal mechanisms.^{28,33} The presence of a high level of chylomicron-like, intestinally-derived lipoproteins in type V alcoholic hypertriglyceridaemia has also been shown by the




Fig 1—Sinusoidal endothelium of normal human liver tissue.
($\times 11\,200$).

presence of two forms of apolipoprotein B (apo B). Apo B₄₈ is synthesised by the intestine and is a marker of chylomicrons, whereas apo B₁₀₀ is derived from the liver:^{34,35} apo B₄₈ can form up to 10% of the total VLDL fraction in type V hyperlipoproteinaemia.³⁴ In such individuals, apo B₄₈ is removed more slowly from the plasma than is apo B₁₀₀ and patients with type V hyperlipoproteinaemia may therefore have defective catabolism of chylomicron remnants.³⁵ Schneider et al³⁶ studied the levels of lipoprotein lipase and hepatic triglyceride lipase and suggested that ethanol-induced hyperlipoproteinaemia results from a combination of increased synthesis and decreased catabolism.




Fig 2—Sinusoid from a patient with alcoholic type V hyperlipoproteinaemia.

Arrow indicates a fenestra; s = sinusoid; e = endothelium; h = hepatocyte; fsc = fat-storing cells; ex = fat-storing cell extension; c = collagen.
($\times 11\,200$).




Fig 3—Sinusoidal endothelium of a patient with alcoholic type V hyperlipoproteinaemia.

rbc = red blood cell.
($\times 7500$).

HYPOTHESIS

We postulate that, in our patient, a decreased porosity of the liver sieve—caused by capillarisation of the sinusoidal endothelium, a reduction in the number of fenestrae, and the presence of collagen within the space of Disse—is a barrier to the usual movement of chylomicron remnants to the hepatocytes for catabolism. This hypothesis would explain the delayed catabolism of apo B₄₈-containing intestinal lipoproteins in alcoholic type V hypertriglyceridaemia. Low porosity of the liver sieve would decrease the passage of chylomicron remnants, while increased liver synthesis of triglyceride-rich lipoproteins would result from a lack of down-regulation of 3-hydroxy-3-methylglutaryl CoA reductase.^{6,10} Changes in the porosity of the liver sieve after ethanol ingestion have been reported in animals^{7,17,20} and in man.²¹ In alcohol-induced type V hyperlipoproteinaemia, abstinence from alcohol substantially reduces the triglyceride levels, which would suggest that the liver sieve regains its normal porosity. We predict that, as in animals, the porosity of the liver sieve in humans will be shown to have a profound influence on lipoprotein metabolism, liver pathology, and atherogenesis.

This work was supported by the Medical Research Council and the National Heart Foundation of New Zealand. We thank Miss S. Townshend for editorial advice, Miss J. E. A. Patterson for secretarial assistance, and Mrs L. M. Bowler-Fraser for technical assistance.

Correspondence should be addressed to R. F., Department of Pathology, Christchurch School of Medicine, Christchurch, New Zealand.

REFERENCES

- 1 Yamagishi M. Electron microscope studies on the fine structure of the sinusoidal wall and fat-storing cells of rabbit livers. *Arch Histol* 1959; **18**: 223–61.
- 2 Wisse E. An electron microscopic study of the fenestrated endothelial lining of rat liver sinusoids. *J Ultrastruct Res* 1970; **31**: 125–50.

- 3 Fraser R, Bosanquet AG, Day WA. Filtration of chylomicrons by the liver may influence cholesterol metabolism and atherosclerosis. *Atherosclerosis* 1978; **29**: 113–23.
- 4 Naito M, Wisse E. Filtration effect of endothelial fenestrations on chylomicron transport in neonatal rat liver sinusoids. *Cell Tissue Res* 1978; **190**: 371–82.
- 5 Wisse E, De Zanger RB, Charels K, Van der Smussen P, McCuskey RS. The liver sieve: considerations concerning the structure and function of endothelial fenestrae, the sinusoidal wall and the space of Disse. *Hepatology* 1985; **5**: 683–92.
- 6 Fraser R, Day WA, Fernando NS. Review: the liver sinusoidal cells. Their role in disorders of the liver, lipoprotein metabolism and atherogenesis. *Pathology* 1986; **18**: 5–11.
- 7 Fraser R, Bowler LM, Day WA. Damage of rat liver sinusoidal endothelium by ethanol. *Pathology* 1980; **12**: 371–76.
- 8 Fraser R, Bowler LM, Day WA, Dobbs B, Johnson HD, Lee D. High perfusion pressure damages the sieving ability of sinusoidal endothelium in rat livers. *Br J Exp Pathol* 1980; **61**: 222–28.
- 9 Wright PL, Smith KF, Day WA, Fraser R. Hepatic sinusoidal endothelium in sheep. An ultrastructural reinvestigation. *Anat Rec* 1983; **206**: 385–90.
- 10 Wright PL, Smith KF, Day WA, Fraser R. Small liver fenestrae may explain the susceptibility of rabbits to atherosclerosis. *Arteriosclerosis* 1983; **3**: 344–48.
- 11 Fraser R, Heslop VR, Murray FEM, Day WA. Ultrastructural studies of the portal transport of fat in chickens. *Br J Exp Pathol* 1986; **67**: 783–91.
- 12 Fraser R, Day WA, Fernando NS. Atherosclerosis and the liver sieve. In: Kim A, Knook DC, Wisse E, eds. Cells of the hepatic sinusoid, vol 1. Rijswijk, Netherlands. Kupffer Cell Foundation, 1986: 217–22.
- 13 Fraser R, Clark SA, Day WA, Murray FEM. Nicotine decreases the porosity of the rat liver sieve—a possible mechanism for hypercholesterolaemia. *Br J Exp Pathol* 1988; **69**: 345–50.
- 14 Wisse E, Van Dierendonck JH, De Zanger RB, Fraser R, McCuskey RS. On the role of liver endothelial filter in the transport of particulate fat (chylomicrons and their remnants) to parenchymal cells and the influence of certain hormones on the endothelial fenestrae. In: Popper H, ed. Communications of liver cells. Lancaster: MTP Press, 1980: 195–200.
- 15 Fraser R, Clark SA, Bowler LM, et al. Concurrent dilation of hepatic fenestrae and decrease in serum cholesterol in the pantethine/cholesterol-fed rabbit. *NZ Med J* 1987; **100**: 357.
- 16 Van der Smussen P, Van Bossuyt H, Charels K, Wisse E. The structure and function of the cytoskeleton in sinusoidal endothelial cells in the rat liver. In: Kim A, Knook DL, Wisse E, eds. Cells of the hepatic sinusoid, vol 1. Rijswijk, Netherlands. Kupffer Cell Foundation, 1986: 517–22.
- 17 Charels K, De Zanger RB, Van Bossuyt H, Van der Smussen P, Wisse E. Influence of acute alcohol administration on endothelial fenestrae of rat livers. In: Kim A, Knook DL, Wisse E, eds. Cells of the hepatic sinusoid, vol 1. Rijswijk, Netherlands. Kupffer Cell Foundation, 1986: 497–502.
- 18 Steffan AM, Gendreau JL, Kim A. Increase in the number of fenestrae in mouse endothelial liver cells by altering the cytoskeleton with cytochalasin B. *Hepatology* 1987; **7**: 1230–38.
- 19 Fraser R, Day WA, Wright PL. Fatty liver and the sinusoidal cells. In: Garlick D, ed. Festschrift for F. C. Courtice. Sydney: University of New South Wales Press, 1981: 139–44.
- 20 Mak KM, Lieber CS. Alterations in endothelial fenestrations in liver sinusoids of baboons fed alcohol: a scanning electron microscope study. *Hepatology* 1984; **4**: 386–91.
- 21 Horn H, Christoffersen P, Henriksen JH. Alcoholic liver injury: defenestration in noncirrhotic livers—a scanning electron microscopic study. *Hepatology* 1987; **7**: 77–82.
- 22 Bioulac-Sage P, Lamouliatte H, Sarric J, Balabaud C. Perfusion-fixation des biopsies hépatiques à l'aiguille: technique. *Gastroenterol Clin Biol* 1984; **8**: 656–59.
- 23 Lamouliatte H, Dubroca J, Quinton A, Balabaud C, Bioulac-Sage P. Sinusoids in human cirrhotic nodules: better identification with the perfusion fixation technique. *J Submicrosc Cytol* 1985; **17**: 279–86.
- 24 Feigl J. Neue Untersuchungen zur chemie des Blutes bei akuter Alkohol-intoxikation und bei chronischem Alkoholismus mit besonderer Berücksichtigung der Fette und Lipide. *Biochem Ztschr* 1918; **92**: 282–90.
- 25 Fredrickson DS, Levy RI. Familial hyperlipoproteinemia. In: Stanbury JB, Wyngaarden SB, Fredrickson DS, eds. The metabolic basis of inherited disease. New York: McGraw Hill, 1972: 545.
- 26 Lewis B. The hyperlipidaemias. Clinical and laboratory practice. Oxford: Blackwell Scientific, 1976: 304.
- 27 Chait A, Mancini M, February AW, Lewis B. Clinical and metabolic study of alcoholic hyperlipidaemia. *Lancet* 1972; **ii**: 62–64.
- 28 Kudzma DJ, Schonfeld G. Alcoholic hyperlipidemia: induction by alcohol but not by carbohydrate. *J Lab Clin Med* 1971; **77**: 384–95.
- 29 Nilsson-Ehle P, Carlstrom S, Belfrage P. Effects of ethanol intake on lipoprotein lipase activity in adipose tissue of fasting subjects. *Lipids* 1978; **13**: 433–41.
- 30 Nikkila EA, Taskinen MR, Huttunen JK. Effects of acute ethanol load on postheparin plasma lipoprotein lipase and hepatic lipase activities and intravenous fat tolerance. *Horm Metab Res* 1978; **10**: 220–29.
- 31 Mendelson JH, Mello NK. Alcohol-induced hyperlipidemia and beta lipoproteins. *Science* 1973; **180**: 1732–74.
- 32 Avogaro P, Cazzolato G. Changes in the composition and physico-chemical characteristics of serum lipoproteins during ethanol-induced lipaemia in alcoholic subjects. *Metabolism* 1975; **24**: 1231–42.
- 33 Schneider J, Panne E, Braun H, Mordasini R, Kaffarnik H. Ethanol-induced hyperlipoproteinemia. Crucial role of preceding ethanol intake in the removal of chylomicrons. *J Lab Clin Med* 1983; **101**: 114–22.
- 34 Meng MS, Gregg RE, Schaeffer EJ, Hoeg JM, Brewer HB. Presence of 2 forms of apolipoprotein B in patients with dyslipoproteinemia. *J Lipid Res* 1983; **24**: 803–09.
- 35 Nestel PJ, Billington T, Fidge NH. Slower removal of intestinal apolipoprotein B48 than of apolipoprotein B100 in severely hypertriglyceridaemic subjects. *Biochim Biophys Acta* 1983; **751**: 422–27.
- 36 Schneider J, Liesenfeld A, Mordasini R, et al. Lipoprotein fractions, lipoprotein lipase and hepatic triglyceride lipase during short-term and long-term uptake of ethanol in healthy subjects. *Atherosclerosis* 1985; **57**: 281–91.