

Id	Name/Reaction	Information	Kinetics
GLUT2	<p><b>Facilitated glucose transporter member 2</b></p> <p>D-glucose (disse) [<b>glc_dis</b>] ↔ D-glucose (cytosol) [<b>glc</b>]</p> <p>D-galactose (disse) [<b>gal_dis</b>] ↔ D-galactose (cytosol) [<b>gal</b>]</p> <p>“Free galactose is transported into cells through a variety of transporters that are members of the glucose transporter (GLUT) family. Deficient transport of galactose into hepatocytes is evident in human patients with defective GLUT2 transporters (Fanconi-Bickel syndrome) resulting in galactose malabsorption/intolerance (Brown, 2000; Leslie, 2003).”</p> <p>“GLUT2 is not completely specific for glucose and is probably also involved in galactose transport (Brown, 2000).”</p> <p>“Data suggests that all transported sugars are likely to use the same outward-facing sugar-binding site on the transporter molecule. This is further indicated by the observation that each of the alternative substrates for each isoform is a competitive inhibitor of deGlc and that D-glucose is a competitive inhibitor of the transport of the alternative substrates. (Colville,</p>	<p><b>Mechanism</b></p> <p>TCDB:2.A.1.1 (glucose transporter subfamily)</p> <p><b>Protein/Structure</b></p> <p>UniProt:<a href="#">P11168</a> (GTR2_HUMAN)</p> <p><b>Gene</b></p> <p>SLC2A2, GLUT2</p> <p><b>Disease</b></p> <p><a href="#">MIM:227810</a> (Fanconi-Bickel syndrome; FBS)</p>	<p>Km(3-O-MG)=42.3±4.1mM (human liver) (Gould, et al., 1991; Walmsley, et al., 1998)</p> <p>Km(3-O-Methyl glc)=17.3 ± 4.3mM (rat liver) (Ciaraldi, et al., 1986)</p> <p><b>Km(D-glc)=21.7 ± 1.8mM</b> (rat liver) (Ciaraldi, et al., 1986)</p> <p><b>Km(D-glc)=66±14mM</b> (rat? hepatocytes) (Elliott and Craik, 1982)</p> <p><b>Km(D-fruct) = 66mM</b> (Walmsley, et al., 1998)</p> <p><b>Km(D-fruct)=212±32mM</b> (rat? hepatocytes) (Elliott and Craik, 1982)</p> <p><b>Km(D-gal)=174±48mM</b> (rat? hepatocytes) (Elliott and Craik, 1982)</p> <p><b>Km(D-gal) &gt; 50mM</b> (GLUT2 enderocytes) (Walmsley, et al., 1998)</p> <p><b>Km(D-gal) = 85.5 ± 10.7mM</b> (human, liver-type GLUT2) (Colville, et al., 1993)</p> <p><b>Km(D-gal) = 92 ± 8.4mM</b> (human, liver-type GLUT2) (Arbuckle, et al., 1996)</p> <p><b>accumulation rate</b> (human GLUT2):</p> <p>v(deoxy-D-glc) = 4.33±0.15 pmol/min/oocyte</p> <p>v(gal) = 1.68±0.09 pmol/min/oocyte</p> <p>v(fru) = 0.78±0.09 pmol/min/oocyte</p> <p><b>v(D-glc)=220±19mmol/min/l of cell H2O</b> (rat? hepatocytes) (Elliott and Craik, 1982)</p> <p><b>v(D-gal)=288±48 mmol/min/l of cell H2O</b> (rat? hepatocytes) (Elliott</p>

	et al., 1993)”		and Craik, 1982) <b>v(D-fruct)=291±26</b> mmol/min/l of cell H2O (rat? hepatocytes) (Elliott and Craik, 1982)  <b>Competitive inhibitor kinetics</b> (Colville, et al., 1993)
GALK	<b>Galactokinase</b> D-galactose [gal] + ATP [atp] ↔ D-galactose 1-phosphate [gal1p] + ADP [adp] <a href="#">EC:2.7.1.6</a> <a href="#">RHEA:13556</a> <a href="#">KEGG:R01092</a>	<b>Protein</b> <a href="#">UniProt:P51570</a> (GALK1_HUMAN) homodimer P51570*2 <b>Gene</b> GALK, GALK1 <b>Disease</b> <a href="#">MIM:230200</a> (GALCT2 Galactosemia II)	<a href="#">SabioRK:P51570</a> <b>two-substrate ordered, ternary complex reaction</b> (Timson and Reece, 2003) <b>kcat(gal) = 8.7±5 1/s</b> (SABIORK:14785)(Timson and Reece, 2003) <b>km(atp) = 0.034±0.004mM</b> (SABIORK:14792)(Timson and Reece, 2003) <b>km(gal)=0.97±0.22 mM</b> (SABIORK:14785) (Timson and Reece, 2003)  <b>km(gal) = 0.436mM</b> (SABIORK:45367), (Sanguuolo, et al., 2004)  <b>Uncompetitive product inhibition</b> of GALK (adult rat liver) by gal1p with both 1mM and 5mM gal1p altering the Km for galactose from 0.150mM to 0.800mM. (Cuatrecasas and Segal, 1965) <b>ki(gal1p) = 5.3mM (5.0-5.7mM)</b> (Cuatrecasas and Segal, 1965) (1mM gal1p caused 15% inhibition, 5mM gal1p 50% inhibition)  Galactokinase being rate limiting for galactose clearance (Schirmer, et al., 1986) -> [27,28, 50]
IMP	<b>Inositol monophosphatase</b> D-galactose 1-phosphate [gal1p] ↔ D-galactose [gal] + phosphate [pi]	<b>Protein</b> <a href="#">UniProt:P29218</a> (IMPA1_HUMAN) Homodimer P29218*2	<b>Competitive inhibition model:</b> <b>km(ino1p) = 0.320±0.050mM</b> (Slepak, et al., 2007) <b>km(gal1p) = 0.35mM</b> (similar kinetics gal1p to ino1p in vitro)

	EC:3.1.3.25	<b>Gene</b> IMPA1, IMPA	<b>(Parthasarathy, et al., 1997)</b>  “Our kinetic analysis demonstrated that gal1p competitively inhibited human IMPase1 by increasing its Km for inositol-1p (ino1p) from 320±50µM to 980±70µM without changing the Vmax (Slepak, et al., 2007).”
GALT	<b>Galactose-1-phosphate uridyl transferase</b> UDP-D-glucose [ <b>udpglc</b> ] + D-galactose 1-phosphate [ <b>gal1p</b> ] ↔ D-glucose 1-phosphate [ <b>glc1p</b> ] + UDP-D-galactose [ <b>udpgal</b> ]. EC:2.7.7.12 <a href="#">RHEA:13992</a> <a href="#">KEGG:R00955</a>  The catalytic mechanism of GALT is <b>ping-pong kinetics</b> with covalent intermediate UMP-enzyme (Facchiano and Marabotti, 2010).	<b>Protein</b> UniProt: <a href="#">P07902</a> (GALT_HUMAN) homodimer P07902*2  <b>Gene</b> GALT  <b>Disease</b> <a href="#">MIM:230400</a> (GALCT Galactosemia)	<b>Mutation analysis</b> (Quimby, et al., 1996) <b>km(gal1p) = 0.57±0.14mM</b> (human, wildtype) (Quimby, et al., 1996) <b>km(udpglc) = 0.21±0.04mM</b> (human, wildtype) (Quimby, et al., 1996)  <b>Mutation analysis</b> (Tang, et al., 2012) <b>km(gal1p) = 1.25±0.36mM</b> (human, wildtype) (Tang, et al., 2012) <b>km(udpglc) = 0.43±0.09mM</b> (human, wildtype) (Tang, et al., 2012)  (?species, 4°C) (Geeganage and Frey, 1998) <b>km(udpglc) = 0.5±0.1mM</b> <b>v(glc1p) = 281± 18 1/s</b> <b>km(glc1p) = 0.37±0.18mM</b> <b>v(glc1p) = 226± 10 1/s</b> <b>km(gal1p) = 0.061±0.020mM</b> <b>v(glc1p) = 166± 13 1/s</b>  <b>Potent linear competent inhibitors UTP and UDP of UDP-glucose</b> (Segal and Rogers, 1971): <b>Ki(UTP) = 0.13mM</b> (rat, liver)

			<b>Ki(UDP) = 0.35mM</b> (rat, liver) Ki(UMP) = 2.3mM (rat, liver) Ki(UDP-glucuronic acid)=0.40mM (rat, liver)
GALE	<b>UDP-glucose 4-epimerase</b> UDP-D-glucose [ <b>udpglc</b> ] ↔ UDP-D-galactose [ <b>udpgal</b> ] EC:5.1.3.2 <a href="#">RHEA:22171</a> <a href="#">KEGG:R00291</a>  Alternative activity with ! GlcNAc ? UDP-GalNAc <-> UDP-GlcNAc	<b>Protein</b> UniProt: <a href="#">Q14376</a> (GALE_HUMAN) homodimer Q14376*2  <b>Gene</b> GALE  <b>Disease</b> <a href="#">MIM:230350</a>	<b>Mutation analysis</b> <a href="#">Timson, 2005</a> <b>km(udpgal)=0.069±0.012mM</b> (human, wildtype) (Timson, 2005) <b>kcat(udpgal) = 36±1.4 1/s</b> (human, wildtype) (Timson, 2005)  <b>km(udpgal) = 0.15 ± 0.02mM</b> (human, wildtype) (Wohlers and Fridovich-Keil, 2000) km(udpgal, V94M) = 0.27 ± 0.01mM (human, V94M) (Wohlers and Fridovich-Keil, 2000) <b>km(udpgal)=0.140± 0.007mM</b> (human, wildtype) (SABIORK:19823) (Winans and Bertozzi, 2002) <b>km(udpgal)=0.120± 0.04mM</b> (human, wildtype) (SABIORK:46260) (Wasilenko, et al., 2005) <b>kcat= 33.8±11.2</b> (human, wildtype) (SABIORK:16222) (Thoden, et al., 2002) <b>km(udpgal) = 0.230±0.06mM</b> (human, wildtype) (SABIORK:46263) (Quimby, et al., 1997)  “Ethanol treatment increases the NADH/NAD ratio in liver (Keppler, et al., 1970) -> [2-4] and by this inhibits the GALE ->[3,5]. Under these conditions oxidation and elimination ->[6-8] of galactose are impaired. Combined galactose+ethanol treatment results in accumulation of gal1p and udpgal in rat liver. The formation of high amounts of udpgal leads to a change in the distribution of liver uracil nucleotides. A marked decrease of udpglc,

			<p>utp, udp and ump is followed by an increase of the sum of uracil nucleotides. (Keppler, et al., 1970)</p> <p>“The GALE reaction is indicated as the rate-limiting step of galactose metabolism in rat liver by the ratio of galactose metabolites (Keppler, et al., 1970)“.</p> <p>“The almost 4-fold increase of gal1p and updgla and the even stronger drop of the udpglc content in the ethanol treated liver after a galactose load demonstrate the ethanol-induced inhibition of the GALE (Keppler, et al., 1970)”.</p> <p>“Galactose provokes pronounced alterations of the uracil nucleotide contents in the liver, which are intensified by an inhibition of the GALE (Keppler, et al., 1970)”</p>
UGP UGALP	<p><b>UDP-glucose pyrophosphorylase</b></p> <p>D-glucose 1-phosphate [<b>glc1p</b>] + UTP [<b>utp</b>] ↔ UDP-glucose [<b>udglc</b>] + diphosphate [<b>pp</b>]</p> <p>EC:2.7.7.9</p> <p><a href="#">RHEA:19892</a></p> <p><a href="#">KEGG:R00289</a></p> <p>in both reaction directions, the enzyme displays simple Michaelis-Menten kinetics (Chang, et al., 1996)</p> <p>MgUTP is a product inhibitor that shows competitive inhibition with respect to UDP-Glc (Chang, et al., 1996)</p> <p><b>UDP-galactose pyrophosphorylase</b></p>	<p><b>Protein</b></p> <p><a href="#">UniProt:Q16851</a> (UGPA_HUMAN)</p> <p>homooctamer Q16851*8</p> <p><b>Gene</b></p> <p>UGP2, UGP1</p> <p>“Previously we reported that stable transfection of human UDP-glucose pyrophosphorylase (hUGP2) rescued galactose-1-phosphate uridyltransferase (GALT)-deficient yeast from “galactose toxicity.” [Lai2002]</p> <p>“Previously, we showed that galactose-1-phosphate competitively</p>	<p>(human, liver, wildtype) (<a href="#">Chang, et al., 1996</a>)</p> <p><b>km(udpglc) = [0.031 - 0.051]mM</b></p> <p><b>km(pp) = [0.172 - 0.210] mM</b></p> <p><b>km(glc1p) = [0.172 - 0.174] mM</b></p> <p><b>km(utp) = [0.563 - 0.692] mM</b></p> <p><b>ki(utp) = 0.477± 41 mM</b> (competitive inhibition with respect to UDP-glc)</p> <p><b>V<sub> fwd</sub>/V<sub> rev</sub> = 0.260</b></p> <p>(human, liver, wildtype) (<a href="#">Duggleby, et al., 1996</a>)</p> <p><b>km(udpglc) = 0.049±0.004mM</b></p> <p><b>km(pp) = 0.166±0.013 mM</b></p> <p><b>km(glc1p) = 0.172±0.010 mM</b></p> <p><b>km(utp) = 0.563±0.115 mM</b></p> <p><b>ki(utp) = 0.643± 0.047 mM</b> (competitive inhibition with respect to UDP-glc)</p>

	<p>D-galactose-1-phosphate [<b>gal1p</b>] + UTP [<b>utp</b>]  + <math>\leftrightarrow</math> UDP-D-galactose [<b>udpgal</b>]  pyrophosphate [<b>pp</b>]  EC:2.7.7.10  <a href="#">RHEA:14212</a>  <a href="#">KEGG:R00502</a>  Could circumvent GALT deficiency (Isselbacher)  Appears to have very low activity, several magnitudes lower than GALDH [Segal1968]  Activity located in glucose dependent enzyme.  <b>1-2 % activity with gal1p (Km gal ~10-20mM)</b></p>	<p>inhibited UDP-glucose pyrophosphorylase, leading to 66% reduction in UDP-glucose/galactose contents in GALT-deficient cells under galactose challenge [Slepek2007-&gt;Lai2002].”</p> <p>The formation of UDP-glucose is the major physiological function of UGP, however at slow rates, the enzyme also catalyzes the phosphorylation of UDP-galactose. (Knop and Hansen, 1970)  “Under normal physiological conditions, this may not be significant, but, in a galactosemic patient, in which the normal galactose metabolism is impaired, the pyrophosphorylase may participate in an abnormal role.</p>	<p><b>ki(udpglc) = 0.013± 4 mM</b> (competitive inhibition with respect to UTP?)  (human, liver, wildtype) (Knop and Hansen, 1970)  <b>keq([udpglc][pp]/([UTP][glc1p])) = 0.15 – 0.16</b>  <b>km(udpglc) = 50mM</b>  <b>km(utp) = 48 mM</b>  <b>km(glc1p) = 95±10 mM</b></p> <p><b>keq([UTP][glc1p]/([udpglc][pp])) = 4.55±0.1</b> (Guynn, et al., 1974) (0.22)  The saturating concentration for UDP-galactose is 10 times that of UDP-glucose:  <b>km(udpgal) = 10*km(udpglc) ~ 0.5mM</b> (human, liver, wildtype) (Knop and Hansen, 1970)  <b>km(udpgal) = 0.420mM</b> (rabbit, liver, wildtype) (Turnquist, et al., 1974)  udpgal was an adequate substrate at 10 times the concentration ofudpglc, showing 14.3% of udpglc (Calf) and 12.0% (Human). activity with udpgal 2-12% of udpglc (12% with 3mM udpgal) (human liver) (Turnquist, et al., 1974)</p> <p>The activity of UDPG:galactose-1-phosphate uridylyltransferase from rat liver under optimal conditions in vitro is less than 5% of the UDPG pyrophosphorylase activity (Keppler, et al., 1970)[Keppler1970 -&gt;39,40]”</p>
ALDR	<p><b>Aldose reductase (galactitol NAD 1-oxidoreductase)</b></p>	<p><b>Protein</b>  <a href="#">UniProt:P15121</a> (ALDR_HUMAN)</p>	<p><a href="#">SabioRK:P15121</a>  <b>km(gal) = 40.0mM</b> (human brain) (SABIORK: 22893) (Wermuth, et</p>

	<p>D-galactose [<b>gal</b>] + NADPH [<b>nadph</b>] + H <math>\leftrightarrow</math> galactitol [<b>galtol</b>] + NADP [<b>nadp</b>]  EC:1.1.1.21  <a href="#">RHEA:12792</a> -&gt; RHEA:37967  <a href="#">KEGG:R01095</a></p> <p>Aldolase reductase is specific for NADPH as cofactor (NADH ~10% of NADPH-dependent activity) (Wermuth and von Wartburg, 1982).</p> <p>“Aldolase reductase catalyzes the conversion of aldoses and a number of other aldehydes to the corresponding alcohol metabolites. It is one of several cytosolic, monomeric, NADPH-dependent aldehyde and ketone reductases of wide substrate specificity (Wermuth, et al., 1982)”.</p>	<p>monomer P15121*1</p> <p><b>Gene</b>  AKR1B1, ALDR1</p> <p>Tissue distribution: Aldolase reductase activity has been demonstrated in brain, kidney, placenta, testis, lens, lung, heart, and pancreas, but appears to be absent from the liver. The enzyme is localized in the cytoplasm of the cell. [Wermuth1982].</p>	<p><b>al., 1982</b></p> <p><b>kcat(gal) = 0.40 1/s</b> (human brain) (SABIORK: 22893) (Wermuth, et al., 1982)</p> <p><b>km(gal) = 110.0mM</b> (human brain) (SABIORK:15695) (Wermuth and von Wartburg, 1982)</p>
PGM1	<p><b>Phosphoglucomutase-1</b></p> <p>D-glucose 1-phosphate [<b>glc1p</b>] <math>\leftrightarrow</math> D-glucose 6-phosphate [<b>glc6p</b>]  EC:5.4.2.2  <a href="#">KEGG:R00959</a>  <a href="#">RHEA:23539</a></p>	<p><b>Multiple isoforms (PGM1, PGM2)</b></p> <p><b>Protein</b>  <a href="#">UniProt:P36871</a> (PGM1_HUMAN)  monomer P36871*1  main isoform for glc1p <math>\leftrightarrow</math> glc6p reaction</p> <p><b>Gene</b>  PGM1</p> <p><b>Disease</b>  <a href="#">Glycogen storage disease 14</a>  (GSD14) [MIM:<a href="#">612934</a>]</p>	<p>The equilibrium lies strongly toward G6P (K<sub>eq</sub>) and the reaction proceeds through a <b>ping-pong mechanism</b> involving aspartyl-phosphoenzyme and glucose-1,6-bisphosphate intermediates.</p> <p>[glc6p]/[glc1p] ~10-12 (Guynn, et al., 1974)  DeltaG = -7.1 kJ/mol [Koenig2012]</p> <p><b>km(glc1p) = 0.049mM</b> (human, RBC) (<b>Quick, et al., 1974</b>)</p> <p><b>km(glc1p) = 0.045mM</b> (rat, heart) (<b>Kashiwaya, et al., 1994</b>)  <b>km(glc6p) = 0.67mM</b> (rat, heart) (Kashiwaya, et al., 1994)</p>

		<p><a href="#">Congenital disorder of glycosylation 1T</a> (CDG1T) [MIM:<a href="#">614921</a>]: A multisystem disorder caused by a defect in glycoprotein biosynthesis and characterized by under-glycosylated serum glycoproteins.</p> <p><b>Protein</b>  <a href="#">UniProt:Q96G03</a>  (PGM2_HUMAN)</p> <p><b>Gene</b>  PGM2</p>	<p><b>km(glc1p) = 0.083mM</b> (human, RBC, PGM1) (<a href="#">Accorsi, et al., 1989</a>)  ki(fru16bp) = 0.092mM (human, RBC, PGM1) (Accorsi, et al., 1989)</p> <p>The kinetic properties of PGM1 and PGM2 are essentially the same. PGM1 is specific for mutation of glucose, whereas PGM2 also has phosphoribomutase activities. (human, RBC) (Accorsi, et al., 1989)</p>
PPASE	<p><b>Pyrophosphatase</b>  Pyrophosphate [<b>pp</b>] + H<sub>2</sub>O → 2 phosphate [<b>pi</b>]  EC:3.6.1.1  <a href="#">RHEA:24579</a>  <a href="#">KEGG:R00004</a></p>	<p><b>Protein</b>  <a href="#">UniProt:Q15181</a> (IPYR_HUMAN)  homodimer Q15181*2</p> <p><b>Gene</b>  PPA1, IOPPP, PP</p>	<p><b>km(pp) = 0.005mM</b> (rat liver) (<a href="#">Yoshida, et al., 1982</a>)  <b>km(pp) = 0.14mM</b> (human erythrocyte) (Thuillier, 1978)  <b>km(pp) = 0.07mM</b> (rat liver) (<a href="#">Irie, et al., 1970</a>)  <b>Delta G0 = -23.56 kJ/mol</b> (Thuillier, 1978)  <b>Delta G0 = -19.2 kJ/mol</b> (Guynn, et al., 1974)</p>
NDKU	<p><b>Nucleoside diphosphokinase (ATP:UDP phosphotransferase)</b>  ATP [<b>atp</b>] + UDP [<b>udp</b>] ↔ ADP [<b>adp</b>] + UTP [<b>udp</b>]  EC: 2.7.4.6  <a href="#">RHEA:25101</a>  <a href="#">KEGG:R00156</a></p>	<p><b>Multiple isoforms</b></p>	<p>Compulsory-order substituted-enzyme (<b>Ping Pong Bi Bi</b>) mechanism (Lam and Packham, 1986)</p> <p><b>km(atp) = 0.38mM</b> (human, platelets) (<a href="#">Lam and Packham, 1986</a>)  <b>km(adp) = 0.024mM</b> (human, platelets) (Lam and Packham, 1986)  <b>km(gtp) = 0.12mM</b> (human, platelets) (Lam and Packham, 1986)</p> <p><b>km(atp) = 1.33mM</b> (rat, liver) (<a href="#">Kimura and Shimada, 1988</a>)  <b>km(adp) = 0.042mM</b> (rat, liver) (Kimura and Shimada, 1988)  <b>km(udp) = 0.19mM</b>(rat, liver) (Kimura and Shimada, 1988)</p>



			<b>km(atp) = 1.80 mM</b> (rat, liver) (Fukuchi, et al., 1994) <b>km(adp) = 0.066 mM</b> (rat, liver) (Fukuchi, et al., 1994) <b>km(utp) = 27.00mM</b> (rat, liver) (Fukuchi, et al., 1994) km(gtp) = 0.15mM (rat, liver) (Fukuchi, et al., 1994) km(gdp) = 0.049mM (rat, liver) (Fukuchi, et al., 1994)
NADPR	<b>NADP reductase</b> NADP [ <b>nadp</b> ] + H <sub>2</sub> →NADPH [ <b>nadph</b> ]  Modeled via pentose phosphate pathway glucose-6-phosphate dehydrogenase D-glucose 6-phosphate [ <b>glc6p</b> ] + NADP [ <b>nadp</b> ] → 6-phospho-D-glucono-1,5- lactone + NADPH [ <b>nadph</b> ] + H EC: 1.1.1.49 <a href="#">RHEA:15844</a> <a href="#">KEGG:R00835</a>	<b>Protein</b> <a href="#">UniProt:P11413</a> (G6PD_HUMAN) homotetramer (dimer of dimer) P11413*4  <b>Gene</b> G6PD	Delta G0 = -19.6 kJ/mol [Schuster1995]  <b>km(glc6p) = 0.040±0.008 mM</b> (human, placenta) (Ozer, et al., 2001) <b>km(nadp) = 0.020±0.010 mM</b> (human, placenta) (Ozer, et al., 2001) <b>ki(nadph) = 0.0171±0.0032 mM</b> (human, placenta) (Ozer, et al., 2001)  <b>km(glc6p) = 0.072 mM</b> (human, RBC) (Bautista, et al., 1992) <b>km(glc6p) = 0.069±0.003 mM</b> (human, recombinant) (Bautista, et al., 1992) <b>km(nadp) = 0.013 mM</b> (human, RBC) (Bautista, et al., 1992) <b>km(nadp) = 0.012±0.002 mM</b> (human, recombinant) (Bautista, et al., 1992) <b>km(nadph) = 0.015±0.002 mM</b> (human, RBC) (Bautista, et al., 1992) <b>km(nadph) = 0.014±0.003 mM</b> (human, recombinant) (Bautista, et al., 1992)  <b>km(glc6p) = 0.326mM</b> (rat, liver) <b>km(glc6p) = 0.157mM</b> (rat, liver) (Corpas, et al., 1995; Corpas, et al., 1995) <b>km(nadp) = 0.108 mM</b> (rat, liver)

			<b>km(nadp) = 0.258 mM</b> (rat, liver) (Corpas, et al., 1995; Corpas, et al., 1995) <b>ki(nadhp) = 0.010 mM</b> (rat, liver) <b>ki(nadhp) = 0.021 mM</b> (rat, liver) (Corpas, et al., 1995; Corpas, et al., 1995)
--	--	--	---

## References

- Accorsi, A., *et al.* (1989) Isoenzymes of phosphoglucomutase from human red blood cells: isolation and kinetic properties, *Preparative biochemistry*, **19**, 251-271.
- Arbuckle, M.I., *et al.* (1996) Structure-function analysis of liver-type (GLUT2) and brain-type (GLUT3) glucose transporters: expression of chimeric transporters in *Xenopus* oocytes suggests an important role for putative transmembrane helix 7 in determining substrate selectivity, *Biochemistry*, **35**, 16519-16527.
- Bautista, J.M., Mason, P.J. and Luzzatto, L. (1992) Purification and properties of human glucose-6-phosphate dehydrogenase made in *E. coli*, *Biochimica et biophysica acta*, **1119**, 74-80.
- Brown, G.K. (2000) Glucose transporters: structure, function and consequences of deficiency, *Journal of inherited metabolic disease*, **23**, 237-246.
- Chang, H.Y., *et al.* (1996) The importance of conserved residues in human liver UDPglucose pyrophosphorylase, *European journal of biochemistry / FEBS*, **236**, 723-728.
- Ciaraldi, T.P., Horuk, R. and Matthaei, S. (1986) Biochemical and functional characterization of the rat liver glucose-transport system. Comparisons with the adipocyte glucose-transport system, *The Biochemical journal*, **240**, 115-123.
- Colville, C.A., *et al.* (1993) Kinetic analysis of the liver-type (GLUT2) and brain-type (GLUT3) glucose transporters in *Xenopus* oocytes: substrate specificities and effects of transport inhibitors, *The Biochemical journal*, **290 ( Pt 3)**, 701-706.
- Corpas, F.J., *et al.* (1995) Kinetic properties of hexose-monophosphate dehydrogenases. II. Isolation and partial purification of 6-phosphogluconate dehydrogenase from rat liver and kidney cortex, *Molecular and cellular biochemistry*, **144**, 97-104.
- Corpas, F.J., *et al.* (1995) Kinetic properties of hexose-monophosphate dehydrogenases. I. Isolation and partial purification of glucose-6-phosphate dehydrogenase from rat liver and kidney cortex, *Life sciences*, **56**, 179-189.
- Cuatrecasas, P. and Segal, S. (1965) Mammalian Galactokinase. Developmental and Adaptive Characteristics in the Rat Liver, *The Journal of biological chemistry*, **240**, 2382-2388.
- Duggleby, R.G., *et al.* (1996) Sequence differences between human muscle and liver cDNAs for UDPglucose pyrophosphorylase and kinetic properties of the recombinant enzymes expressed in *Escherichia coli*, *European journal of biochemistry / FEBS*, **235**, 173-179.
- Elliott, K.R. and Craik, J.D. (1982) Sugar transport across the hepatocyte plasma membrane, *Biochemical Society transactions*, **10**, 12-13.

- Facchiano, A. and Marabotti, A. (2010) Analysis of galactosemia-linked mutations of GALT enzyme using a computational biology approach, *Protein engineering, design & selection : PEDS*, **23**, 103-113.
- Fukuchi, T., *et al.* (1994) Recombinant rat nucleoside diphosphate kinase isoforms (alpha and beta): purification, properties and application to immunological detection of native isoforms in rat tissues, *Biochimica et biophysica acta*, **1205**, 113-122.
- Geeganage, S. and Frey, P.A. (1998) Transient kinetics of formation and reaction of the uridylyl-enzyme form of galactose-1-P uridylyltransferase and its Q168R-variant: insight into the molecular basis of galactosemia, *Biochemistry*, **37**, 14500-14507.
- Gould, G.W., *et al.* (1991) Expression of human glucose transporters in *Xenopus* oocytes: kinetic characterization and substrate specificities of the erythrocyte, liver, and brain isoforms, *Biochemistry*, **30**, 5139-5145.
- Guynn, R.W., *et al.* (1974) The concentration and control of cytoplasmic free inorganic pyrophosphate in rat liver in vivo, *The Biochemical journal*, **140**, 369-375.
- Irie, M., *et al.* (1970) Distribution and properties of alkaline pyrophosphatases of rat liver, *Journal of biochemistry*, **67**, 47-58.
- Kashiwaya, Y., *et al.* (1994) Control of glucose utilization in working perfused rat heart, *The Journal of biological chemistry*, **269**, 25502-25514.
- Keppler, D., Rudigier, J. and Decker, K. (1970) Trapping of uridine phosphates by D-galactose in ethanol-treated liver, *FEBS letters*, **11**, 193-196.
- Keppler, D.O., *et al.* (1970) The trapping of uridine phosphates by D-galactosamine. D-glucosamine, and 2-deoxy-D-galactose. A study on the mechanism of galactosamine hepatitis, *European journal of biochemistry / FEBS*, **17**, 246-253.
- Kimura, N. and Shimada, N. (1988) Membrane-associated nucleoside diphosphate kinase from rat liver. Purification, characterization, and comparison with cytosolic enzyme, *The Journal of biological chemistry*, **263**, 4647-4653.
- Knop, J.K. and Hansen, R.G. (1970) Uridine diphosphate glucose pyrophosphorylase. IV. Crystallization and properties of the enzyme from human liver, *The Journal of biological chemistry*, **245**, 2499-2504.
- Lam, S.C. and Packham, M.A. (1986) Isolation and kinetic studies of nucleoside diphosphokinase from human platelets and effects of cAMP phosphodiesterase inhibitors, *Biochemical pharmacology*, **35**, 4449-4455.
- Leslie, N.D. (2003) Insights into the pathogenesis of galactosemia, *Annual review of nutrition*, **23**, 59-80.
- Ozer, N., Aksoy, Y. and Ogus, I.H. (2001) Kinetic properties of human placental glucose-6-phosphate dehydrogenase, *The international journal of biochemistry & cell biology*, **33**, 221-226.
- Parthasarathy, R., Parthasarathy, L. and Vadnal, R. (1997) Brain inositol monophosphatase identified as a galactose 1-phosphatase, *Brain research*, **778**, 99-106.
- Quick, C.B., Fisher, R.A. and Harris, H. (1974) A kinetic study of the isozymes determined by the three human phosphoglucomutase loci PGM1, PGM2, and PGM3, *European journal of biochemistry / FEBS*, **42**, 511-517.
- Quimby, B.B., *et al.* (1997) Characterization of two mutations associated with epimerase-deficiency galactosemia, by use of a yeast expression system for human UDP-galactose-4-epimerase, *American journal of human genetics*, **61**, 590-598.
- Quimby, B.B., *et al.* (1996) Functional requirements of the active site position 185 in the human enzyme galactose-1-phosphate uridylyltransferase, *The Journal of biological chemistry*, **271**, 26835-26842.

- Sanguuolo, F., *et al.* (2004) Biochemical characterization of two GALK1 mutations in patients with galactokinase deficiency, *Human mutation*, **23**, 396.
- Schirmer, W.J., *et al.* (1986) Galactose clearance as an estimate of effective hepatic blood flow: validation and limitations, *The Journal of surgical research*, **41**, 543-556.
- Segal, S. and Rogers, S. (1971) Nucleotide inhibition of mammalian liver galactose-1-phosphate uridylyltransferase, *Biochimica et biophysica acta*, **250**, 351-360.
- Slepak, T.I., *et al.* (2007) Involvement of endoplasmic reticulum stress in a novel Classic Galactosemia model, *Molecular genetics and metabolism*, **92**, 78-87.
- Tang, M., *et al.* (2012) Correlation assessment among clinical phenotypes, expression analysis and molecular modeling of 14 novel variations in the human galactose-1-phosphate uridylyltransferase gene, *Human mutation*, **33**, 1107-1115.
- Thoden, J.B., *et al.* (2002) Structural analysis of the Y299C mutant of Escherichia coli UDP-galactose 4-epimerase. Teaching an old dog new tricks, *The Journal of biological chemistry*, **277**, 27528-27534.
- Thuillier, L. (1978) Purification and kinetic properties of human erythrocyte Mg<sup>2+</sup>-dependent inorganic pyrophosphatase, *Biochimica et biophysica acta*, **524**, 198-206.
- Timson, D.J. (2005) Functional analysis of disease-causing mutations in human UDP-galactose 4-epimerase, *The FEBS journal*, **272**, 6170-6177.
- Timson, D.J. and Reece, R.J. (2003) Functional analysis of disease-causing mutations in human galactokinase, *European journal of biochemistry / FEBS*, **270**, 1767-1774.
- Turnquist, R.L., *et al.* (1974) Uridine diphosphate glucose pyrophosphorylase: differential heat inactivation and further characterization of human liver enzyme, *Biochimica et biophysica acta*, **364**, 59-67.
- Walmsley, A.R., *et al.* (1998) Sugar transporters from bacteria, parasites and mammals: structure-activity relationships, *Trends in biochemical sciences*, **23**, 476-481.
- Wasilenko, J., *et al.* (2005) Functional characterization of the K257R and G319E-hGALE alleles found in patients with ostensibly peripheral epimerase deficiency galactosemia, *Molecular genetics and metabolism*, **84**, 32-38.
- Wermuth, B., *et al.* (1982) Purification and characterization of human-brain aldose reductase, *European journal of biochemistry / FEBS*, **127**, 279-284.
- Wermuth, B. and von Wartburg, J.P. (1982) Aldose reductase from human tissues, *Methods in enzymology*, **89 Pt D**, 181-186.
- Winans, K.A. and Bertozzi, C.R. (2002) An inhibitor of the human UDP-GlcNAc 4-epimerase identified from a uridine-based library: a strategy to inhibit O-linked glycosylation, *Chemistry & biology*, **9**, 113-129.
- Wohlers, T.M. and Fridovich-Keil, J.L. (2000) Studies of the V94M-substituted human UDPgalactose-4-epimerase enzyme associated with generalized epimerase-deficiency galactosaemia, *Journal of inherited metabolic disease*, **23**, 713-729.
- Yoshida, C., Shah, H. and Weinhouse, S. (1982) Purification and properties of inorganic pyrophosphatase of rat liver and hepatoma 3924A, *Cancer research*, **42**, 3526-3531.