

From Single Hepatocytes to Whole Liver Function: A Multi-Scale Model of Human Hepatic Galactose Metabolism - Supplementary Information

TABLE OF CONTENT

| | |
|--|------------------------------|
| Table of content..... | Error! Bookmark not defined. |
| Supplementary Tables | Error! Bookmark not defined. |
| Supplementary Table 1 - Reactions and transporters in human galactose metabolism and kinetic parameters. 2 | |
| Supplementary Table 2 - Metabolites in hepatic galactose metabolism. | 10 |
| Supplementary Table 3 - Tissue-and organ parameters..... | 17 |
| Supplementary Table 4 – Parameter for the log-normal distributions..... | 22 |
| Supplementary Table 5 – Diffusion coefficients | 23 |
| Supplementary Table 6 – Organ/Liver parameters..... | 24 |
| Supplementary Figures..... | 27 |
| Supplementary Figure 1 – Validation of single cell model of galactose metabolism Time courses in GALE inhibition via ethanol. | 27 |
| Supplementary Figure 2 – Metabolic Control Analysis (MCA) - Role of the different parameters in controlling hepatic galactose metabolism on a single cell level | 28 |
| Supplementary Figure 2 –Multiple indicator dilution curves in galactosemias | 28 |
| Supplementary Figure 3 – Altered hepatic galactose clearance and dependency on flow in galactosemias analogue to Figure 4. | 28 |
| References | 28 |

ABBREVIATIONS

IVM – in vivo microscopy
IPM – in plastic microscopy
OPS – orthogonal polarization spectral imaging
QSD – quantitative stereological description
SEM – scanning electron microscopy
SE – standard error (measurement)
SD – standard deviation
TEM – transmission electron microscopy

SUPPLEMENTARY TABLES

Supplementary Table 1 - Reactions and transporters in human galactose metabolism and kinetic parameters.

| Id | Information | Kinetics |
|--------------|--|--|
| GLUT2 | <p>Facilitated glucose transporter member 2</p> <p>D-glucose (disse) [glc_dis] ↔ D-glucose (cytosol) [glc] D-galactose (disse) [gal_dis] ↔ D-galactose (cytosol) [gal]</p> <p>Mechanism TCDB:2.A.1.1 (glucose transporter subfamily)</p> <p>Protein/Structure UniProt:P11168 (GTR2_HUMAN)</p> <p>Gene SLC2A2, GLUT2</p> <p>Disease MIM:227810 (Fanconi-Bickel syndrome; FBS)</p> <p>Galactose and glucose transported via GLUT2 (competitive inhibition kinetics) (Brown, 2000; Colville, et al., 1993)</p> <p>Deficient transport of galactose into hepatocytes in human patients with defective GLUT2 transporters (Fanconi-Bickel syndrome) resulting in galactose malabsorption/intolerance (Brown, 2000; Leslie, 2003).</p> | <p>km(D-glc)=21.7 ± 1.8mM (rat liver) (Ciaraldi, et al., 1986) km(D-glc)=66±14mM (rat? hepatocytes) (Elliott and Craik, 1982) km(3-O-MG)=42.3±4.1mM (human liver) (Gould, et al., 1991; Walmsley, et al., 1998) km(3-O-Methyl glc)=17.3 ± 4.3mM (rat liver) (Ciaraldi, et al., 1986)</p> <p>km(D-gal)=174±48mM (rat? hepatocytes) (Elliott and Craik, 1982) km(D-gal) > 50mM (GLUT2 enderocytes) (Walmsley, et al., 1998) km(D-gal) = 85.5 ± 10.7mM (human, liver-type GLUT2) (Colville, et al., 1993) km(D-gal) = 92 ± 8.4mM (human, liver-type GLUT2) (Arbuckle, et al., 1996)</p> <p>Km(D-fru) = 66mM (Walmsley, et al., 1998) Km(D-fru)=212±32mM (rat? hepatocytes) (Elliott and Craik, 1982)</p> <p>Accumulation rate (human GLUT2) v(deoxy-D-glc) = 4.33±0.15 pmol/min/oocyte v(D-gal) = 1.68±0.09 pmol/min/oocyte v(D-fru) = 0.78±0.09 pmol/min/oocyte</p> <p>v(D-glc)=220±19mmol/min/l of cell H2O (rat? hepatocytes) (Elliott and Craik, 1982) v(D-gal)=288±48 mmol/min/l of cell H2O (rat? hepatocytes) (Elliott and Craik, 1982) v(D-fru)=291±26 mmol/min/l of cell H2O (rat? hepatocytes) (Elliott and Craik, 1982)</p> |
| GALK | <p>Galactokinase</p> <p>D-galactose [gal] + ATP [atp] ↔ D-galactose 1-phosphate [gal1p] + ADP [adp]</p> <p>Reaction EC:2.7.1.6 RHEA:13556 KEGG:R01092</p> <p>Protein</p> | <p>Two-substrate ordered, ternary complex reaction (Timson and Reece, 2003)</p> <p>kcat(gal) = 8.7±5 1/s (SABIORK:14785)(Timson and Reece, 2003) km(atp) = 0.034±0.004mM (SABIORK:14792)(Timson and Reece, 2003) km(gal)=0.97±0.22 mM (SABIORK:14785) (Timson and Reece, 2003) km(gal) = 0.436mM (SABIORK:45367), (Sanguuolo, et al., 2004)</p> <p>Uncompetitive product inhibition of GALK (adult rat liver) by gal1p with both 1mM and 5mM gal1p altering the Km for galactose from 0.150mM to 0.800mM</p> |

Commented [WU1]: TODO. Include the missing references
TODO: Update the SABIO-RK entries as soon as available in database
TODO: crosscheck the table with annotations in SBML

| | | |
|-------------|--|--|
| | <p><u>UniProt:P51570</u> (GALK1_HUMAN) homodimer P51570*2</p> <p>Gene GALK, GALK1</p> <p>Disease <u>MIM:230200</u> (GALCT2 Galactosemia II)</p> <p>Galactokinase being rate limiting for galactose clearance (Schirmer, et al., 1986) -> [27,28, 50]</p> | <p>(1mM gal1p caused 15% inhibition, 5mM gal1p 50% inhibition) ki(gal1p) = 5.3mM (5.0-5.7mM) (adult rat liver) (Cuatrecasas and Segal, 1965)</p> |
| IMP | <p>Inositol monophosphatase D-galactose 1-phosphate [gal1p] ↔ D-galactose [gal] + phosphate [pi]</p> <p>Reaction EC:3.1.3.25</p> <p>Protein <u>UniProt:P29218</u> (IMPA1_HUMAN) homodimer P29218*2</p> <p>Gene IMPA1, IMPA</p> <p>Normal substrate inositol-1p (ino1p)</p> | <p>Competitive inhibition model Kinetic analysis demonstrated that gal1p competitively inhibited human IMP1 by increasing Km for inositol-1p (ino1p) from 320±50μM to 980±70μM without changing the Vmax (Slepek, et al., 2007) km(ino1p) = 0.320±0.050mM (Slepek, et al., 2007) km(gal1p) = 0.35mM (similar kinetics gal1p to ino1p in vitro) (Parthasarathy, et al., 1997)</p> |
| GALT | <p>Galactose-1-phosphate uridyl transferase UDP-D-glucose [udpglc] + D-galactose 1-phosphate [gal1p] ↔ D-glucose 1-phosphate [glc1p] + UDP-D-galactose [udpgal].</p> <p>Reaction EC:2.7.7.12 <u>RHEA:13992</u> <u>KEGG:R00955</u></p> <p>Protein <u>UniProt:P07902</u> (GALT_HUMAN) homodimer P07902*2</p> <p>Gene GALT</p> <p>Disease <u>MIM:230400</u> (GALCT Galactosemia)</p> | <p>The catalytic mechanism of GALT is ping-pong kinetics with covalent intermediate UMP-enzyme (Facchiano and Marabotti, 2010).</p> <p>Mutation analysis (Quimby, et al., 1996) km(gal1p) = 0.57±0.14mM (human, wildtype) (Quimby, et al., 1996) km(udpglc) = 0.21±0.04mM (human, wildtype) (Quimby, et al., 1996)</p> <p>Mutation analysis (Tang, et al., 2012) km(gal1p) = 1.25±0.36mM (human, wildtype) (Tang, et al., 2012) km(udpglc) = 0.43±0.09mM (human, wildtype) (Tang, et al., 2012)</p> <p>(?species, 4°C) (Geeganage and Frey, 1998) km(udpglc) = 0.5±0.1mM v(glc1p) = 281± 18 1/s km(glc1p) = 0.37±0.18mM v(glc1p) = 226± 10 1/s km(gal1p) = 0.061±0.020mM v(glc1p) = 166± 13 1/s</p> |

| | |
|------|--|
| | <p>Potent linear competent inhibitors UTP and UDP of UDP-glucose (Segal and Rogers, 1971):</p> <p>Ki(UTP) = 0.13mM (rat, liver)</p> <p>Ki(UDP) = 0.35mM (rat, liver)</p> <p>Ki(UMP) = 2.3mM (rat, liver)</p> <p>Ki(UDP-glucuronic acid)=0.40mM (rat, liver)</p> |
| GALE | <p>UDP-glucose 4-epimerase</p> <p>UDP-D-glucose [udpglc] ↔ UDP-D-galactose [udpgal]</p> <p>Reaction</p> <p>EC:5.1.3.2</p> <p>RHEA:22171</p> <p>KEGG:R00291</p> <p>Protein</p> <p>UniProt:Q14376 (GALE_HUMAN)</p> <p>homodimer Q14376*2</p> <p>Gene</p> <p>GALE</p> <p>Disease</p> <p>MIM:230350 (GALE deficiency)</p> <p>Alternative activity with GlcNAc:</p> <p>UDP-GalNAc ↔ UDP-GlcNAc</p> <p>“Ethanol treatment increases the NADH/NAD ratio in liver (Keppler, et al., 1970) [2-4] and by this inhibits the GALE [3,5]. Under these conditions oxidation and elimination [6-8] of galactose are impaired. Combined galactose+ethanol treatment results in accumulation of gal1p and udpgal in rat liver. The formation of high amounts of udpgal leads to a change in the distribution of liver uracil nucleotides. A marked decrease of udpglc, utp, udp and ump is followed by an increase of the sum of uracil nucleotides.” (Keppler, et al., 1970)</p> <p>“The GALE reaction is indicated as the rate-limiting step of galactose metabolism in rat liver by the ratio of galactose metabolites (Keppler, et al., 1970)“.</p> <p>“The almost 4-fold increase of gal1p and updgal and the even stronger drop of the udpglc content in the ethanol treated liver after a galactose load demonstrates the ethanol-induced</p> |

Potent linear competent inhibitors UTP and UDP of UDP-glucose (Segal and Rogers, 1971):

Ki(UTP) = 0.13mM (rat, liver)

Ki(UDP) = 0.35mM (rat, liver)

Ki(UMP) = 2.3mM (rat, liver)

Ki(UDP-glucuronic acid)=0.40mM (rat, liver)

Mutation analysis(Timson, 2005)

km(udpgal)=0.069±0.012mM (human, wildtype) (Timson, 2005)

kcat(udpgal) = 36±1.4 1/s (human, wildtype) (Timson, 2005)

km(udpgal) = 0.15 ± 0.02mM (human, wildtype) (Wohlers and Fridovich-Keil, 2000)

km(udpgal, V94M) = 0.27 ± 0.01mM (human, V94M) (Wohlers and Fridovich-Keil, 2000)

km(udpgal)=0.140± 0.007mM (human, wildtype) (SABIORK:19823) (Winans and Bertozzi, 2002)

km(udpgal)=0.120± 0.04mM (human, wildtype) (SABIORK:46260) (Wasilenko, et al., 2005)

kcat= 33.8±11.2 (human, wildtype) (SABIORK:16222) (Thoden, et al., 2002)

km(udpgal) = 0.230±0.06mM (human, wildtype) (SABIORK:46263) (Quimby, et al., 1997)

inhibition of the GALE” (Keppler, et al., 1970).
“Galactose provokes pronounced alterations of the uracil nucleotide contents in the liver, which are intensified by an inhibition of the GALE” (Keppler, et al., 1970)

| | | |
|--------------|--|---|
| UGP | UDP-glucose pyrophosphorylase D-glucose 1-phosphate [glc1p] + UTP [utp] ↔ UDP-glucose [udglc]+ diphosphate [pp] Reaction EC:2.7.7.9 RHEA:19892 KEGG:R00289 Protein UniProt:Q16851 (UGPA_HUMAN) homooctamer Q16851*8 Gene UGP2, UGP1 | Enzyme displays simple Michaelis-Menten kinetics in both directions (Chang, et al., 1996) MgUTP is a product inhibitor that shows competitive inhibition with respect to UDP-Glc (Chang, et al., 1996) (human, liver, wildtype) (Chang, et al., 1996) km(udpglc) = [0.031 - 0.051]mM km(pp) = [0.172 - 0.210] mM km(glc1p) = [0.172 - 0.174] mM km(utp) = [0.563 - 0.692] mM ki(utp) = 0.477± 41 mM (competitive inhibition with respect to UDP-glc) V_{fwd}/V_{rev} = 0.260 (human, liver, wildtype) (Duggleby, et al., 1996) km(udpglc) = 0.049±0.004mM km(pp) = 0.166±0.013 mM km(glc1p) = 0.172±0.010 mM km(utp) = 0.563±0.115 mM ki(utp) = 0.643± 0.047 mM (competitive inhibition with respect to UDP-glc) ki(udpglc) = 0.013± 4 mM (competitive inhibition with respect to UTP?) (human, liver, wildtype) (Knop and Hansen, 1970) keq([udpglc][pp]/([UTP][glc1p])) = 0.15 – 0.16 km(udpglc) = 50mM km(utp) = 48 mM km(glc1p) = 95±10 mM keq([UTP][glc1p]/([udpglc][pp])) = 4.55±0.1 (Guynn, et al., 1974) (0.22) The saturating concentration for UDP-galactose is 10 times that of UDP-glucose: km(udpgal) = 10*km(udpglc) ~ 0.5mM (human, liver, wildtype) (Knop and Hansen, 1970) km(udpgal) = 0.420mM (rabbit, liver, wildtype) (Turnquist, et al., 1974) udpgal was an adequate substrate at 10 times the concentration of udpglc, showing 14.3% of udpglc (Calf) and 12.0% (Human). activity with udpgal 2-12% of udpglc (12% with 3mM udpgal) (human liver) (Turnquist, et al., 1974) |
| UGALP | UDP-galactose pyrophosphorylase D-galactose-1-phosphate [gal1p] + UTP [utp] + ↔ UDP-D-galactose [udpgal] pyrophosphate [pp] Reaction EC:2.7.7.10 RHEA:14212 KEGG:R00502 Protein UniProt:Q16851 (UGPA_HUMAN) homooctamer Q16851*8 Gene UGP2, UGP1 The formation of UDP-glucose is the major physiological function of UGP, however at slow rates, the enzyme also catalyzes the phosphorylation of UDP-galactose (Knop and Hansen, 1970) [Segal1968]. Not significant in normal physiological conditions, but in galactosemic patients could circumvent GALT deficiency | |

| | | |
|-------------|--|---|
| | <p>(Isselbacher ?).</p> <p>Stable transfection of human UGP (hUGP2) rescued galactose GALT deficient yeast from “galactose toxicity [Lai2002].</p> | <p>“The activity of UDPG:galactose-1-phosphate uridylyltransferase from rat liver under optimal conditions in vitro is less than 5% of the UDPG pyrophosphorylase activity” (Keppler, et al., 1970)[Keppler1970 ->39,40]</p> <p>gal1p as competitive inhibitor of glc1p</p> <p>“Previously, we showed that galactose-1-phosphate competitively inhibited UDP-glucose pyrophosphorylase, leading to 66% reduction in UDP-glucose/galactose contents in GALT-deficient cells under galactose challenge” [Slepak2007->Lai2002].</p> |
| ALDR | <p>Aldose reductase (galactitol NAD 1-oxidoreductase)</p> <p>D-galactose [gal] + NADPH [nadph] + H ↔ galactitol [galtol] + NADP [nadp]</p> <p>Reaction EC:1.1.1.21 RHEA:12792 -> RHEA:37967 KEGG:R01095</p> <p>Protein UniProt:P15121 (ALDR_HUMAN) monomer P15121*1</p> <p>Gene AKR1B1, ALDR1</p> <p>Aldolase reductase is specific for NADPH as cofactor (NADH ~10% of NADPH-dependent activity) (Wermuth and von Wartburg, 1982). “Aldolase reductase catalyzes the conversion of aldoses and a number of other aldehydes to the corresponding alcohol metabolites. It is one of several cytosolic, monomeric, NADPH-dependent aldehyde and ketone reductases of wide substrate specificity” (Wermuth, et al., 1982)”.</p> | <p>km(gal) = 40.0mM (human brain) (SABIORK:22893) (Wermuth, et al., 1982) kcat(gal) = 0.40 1/s (human brain) (SABIORK:22893) (Wermuth, et al., 1982)</p> <p>km(gal) = 110.0mM (human brain) (SABIORK:15695) (Wermuth and von Wartburg, 1982)</p> |
| PGM1 | <p>Phosphoglucumutase-1</p> <p>D-glucose 1-phosphate [glc1p] ↔ D-glucose 6-phosphate [glc6p]</p> <p>Reaction EC:5.4.2.2 KEGG:R00959 RHEA:23539</p> <p>Protein (multiple isoforms PGM1, PGM2)</p> | <p>The equilibrium lies strongly toward glc6p and reaction proceeds through ping-pong mechanism (Guynn, et al., 1974) The kinetic properties of PGM1 and PGM2 are essentially the same. PGM1 is specific for mutation of glucose, whereas PGM2 also has phosphoribomutase activities. (human, RBC) (Accorsi, et al., 1989)</p> <p>[glc6p]/[glc1p] ~10-12 (Guynn, et al., 1974) DeltaG = -7.1 kJ/mol (König, et al., 2012)</p> |

| | | |
|--------------|--|---|
| | <p>UniProt:P36871 (PGM1_HUMAN) monomer P36871*1 main isoform for glc1p ↔ glc6p reaction</p> <p>Gene PGM1</p> <p>Disease MIM:612934 (Glycogen storage disease 14) MIM:614921 (Congenital disorder of glycosylation 1T CDG1T)</p> <p>Protein UniProt:Q96G03 (PGM2_HUMAN)</p> <p>Gene PGM2</p> <p>CDG1T - A multisystem disorder caused by a defect in glycoprotein biosynthesis and characterized by under-glycosylated serum glycoproteins.</p> | <p>km(glc1p) = 0.049mM (human, RBC) (Quick, et al., 1974)</p> <p>km(glc1p) = 0.045mM (rat, heart) (Kashiwaya, et al., 1994) km(glc6p) = 0.67mM (rat, heart) (Kashiwaya, et al., 1994)</p> <p>km(glc1p) = 0.083mM (human, RBC, PGM1) (Accorsi, et al., 1989) ki(fru16bp) = 0.092mM (human, RBC, PGM1) (Accorsi, et al., 1989)</p> |
| PPASE | <p>Pyrophosphatase Pyrophosphate [pp] + H2O → 2 phosphate [pi]</p> <p>Reaction EC:3.6.1.1 RHEA:24579 KEGG:R00004</p> <p>Protein UniProt:Q15181 (IPYR_HUMAN) homodimer Q15181*2</p> <p>Gene PPA1, IOPPP, PP</p> | <p>km(pp) = 0.005mM (rat liver) (Yoshida, et al., 1982) km(pp) = 0.14mM (human erythrocyte) (Thuillier, 1978) km(pp) = 0.07mM (rat liver) (Irie, et al., 1970) Delta G0 = -23.56 kJ/mol (Thuillier, 1978) Delta G0 = -19.2 kJ/mol (Guynn, et al., 1974)</p> |
| NDKU | <p>Nucleoside diphosphokinase (ATP:UDP phosphotransferase) ATP [atp] + UDP [udp] ↔ ADP [adp] + UTP [udp]</p> <p>Reaction EC: 2.7.4.6 RHEA:25101 KEGG:R00156</p> <p>Multitude of isoforms</p> | <p>Compulsory-order substituted-enzyme (Ping Pong Bi Bi) mechanism (Lam and Packham, 1986)</p> <p>km(atp) = 0.38mM (human, platelets) (Lam and Packham, 1986) km(adp) = 0.024mM (human, platelets) (Lam and Packham, 1986) km(gtp) = 0.12mM (human, platelets) (Lam and Packham, 1986)</p> <p>km(atp) = 1.33mM (rat, liver) (Kimura and Shimada, 1988) km(adp) = 0.042mM (rat, liver) (Kimura and Shimada, 1988) km(udp) = 0.19mM(rat, liver) (Kimura and Shimada, 1988)</p> |

| | | |
|--|--|---|
| | | km(atp) = 1.80 mM (rat, liver) (Fukuchi, et al., 1994) km(adp) = 0.066 mM (rat, liver) (Fukuchi, et al., 1994) km(utp) = 27.00mM (rat, liver) (Fukuchi, et al., 1994) km(gtp) = 0.15mM (rat, liver) (Fukuchi, et al., 1994) km(gdp) = 0.049mM (rat, liver) (Fukuchi, et al., 1994) |
| NADPR | NADP reductase NADP [nadp] + H ₂ →NADPH [nadph] Modeled via glucose-6-phosphate dehydrogenase in pentose phosphate pathway D-glucose 6-phosphate [glc6p] + NADP [nadp] → 6-phospho-D-glucono-1,5-lactone + NADPH [nadph] + H Reaction EC: 1.1.1.49 RHEA:15844 KEGG:R00835 Protein UniProt:P11413 (G6PD_HUMAN) homotetramer (dimer of dimer) P11413*4 Gene G6PD | Delta G0 = -19.6 kJ/mol [Schuster1995] km(glc6p) = 0.040±0.008 mM (human, placenta) (Ozer, et al., 2001) km(nadp) = 0.020±0.010 mM (human, placenta) (Ozer, et al., 2001) ki(nadph) = 0.0171±0.0032 mM (human, placenta) (Ozer, et al., 2001) km(glc6p) = 0.072 mM (human, RBC) (Bautista, et al., 1992) km(glc6p) = 0.069±0.003 mM (human, recombinant) (Bautista, et al., 1992) km(nadp) = 0.013 mM (human, RBC) (Bautista, et al., 1992) km(nadp) = 0.012±0.002 mM (human, recombinant) (Bautista, et al., 1992) km(nadph) = 0.015±0.002 mM (human, RBC) (Bautista, et al., 1992) km(nadph) = 0.014±0.003 mM (human, recombinant) (Bautista, et al., 1992) km(glc6p) = 0.326mM (rat, liver) km(glc6p) = 0.157mM (rat, liver) (Corpas, et al., 1995; Corpas, et al., 1995) km(nadp) = 0.108 mM (rat, liver) km(nadp) = 0.258 mM (rat, liver) (Corpas, et al., 1995; Corpas, et al., 1995) ki(nadhp) = 0.010 mM (rat, liver) ki(nadhp) = 0.021 mM (rat, liver) (Corpas, et al., 1995; Corpas, et al., 1995) |
| ATPS | ATP synthesis ADP [adp] + phosphate [pi] → ATP [atp] Modeled via general ATP producing reaction representative for ATP production via glycolysis and oxidative phosphorylation | |
| GTF GTFGAL GTFGLC | Glycosyltransferase Acceptor [gac] + UDP-glucose [udpglc] -> Acceptor-glucose[gacglc] + UDP [udp] Acceptor [gac] + UDP-glucose [udpgal] -> Acceptor-glucose[gacgal] + UDP [udp] Enzymes that transfer mono- or oligosaccharides from donor molecules to growing oligosaccharide chains or proteins are called glycosyltransferases (Gtfs) | |

GALDH**Galactose 1-dehydrogenase**

D-galactose + NAD⁺ ↔ D-galactono-1,4-lactone + NADH + H⁺

EC.1.1.1.48 (Brenda only bacteria)

D-galactose → galactonate

(first enzyme in oxidative pathway)

[Segal1968 → Cuatrecasas1966,15]

Alternative pathway to xylulose.

D-Galactose + Oxygen + H₂O ⇌ D-Galactonate + Hydrogen
peroxide

EC:1.1.3.9

KEGG:R01098

(only bacteria)

Supplementary Table 2 - Metabolites in hepatic galactose metabolism.

| Id | Name (mass) Annotation | Initial Concentration | Comments |
|--------------|--|---|---|
| glc | D-glucose (M _w 180.2) <u>CHEBI:4167</u> <u>KEGG:C00031</u> | 5.5mM (König, et al., 2012) | [glc] = 5.5mM (König, et al., 2012) 3-10mM (depending on physiological state) |
| gal | D-galactose (M _w 180.2) <u>CHEBI:4139</u> <u>KEGG:C00124</u> | 0.00012mM (no galactose) 0.00144mM (GALT deficient) 0.0013-0.0027mM (GALE deficient) | plasma of post-absorptive humans (data considerable lower (3-18-fold) than conventional enzymatic assay) (Schadewaldt, et al., 2000) [gal] = 0.12±0.03μM (n=16) healthy subjects [gal] = 1.44±0.54μM (n=10) classical galactosemia (GALT deficiency) [gal] = 0.17±0.07μM (n=5) obligate heterozygous parents of classical galactosemia [gal] = 0.11±0.04μM (n=15) diabetic patients GALE deficient patients (blood) (Yamaguchi, et al., 1989) [gal]=24-29mg/L (0.013-0.016mM) [gal]= 48mg/L (0.027mM) Neonatal control (blood): [gal]=13±6 mg/L (0.0072±0.0033mM) (Yamaguchi, et al., 1989) normal values: [gal]= 0.015±0.009mM (range 0-0.044mM) (Orfanos, et al., 1986) Cut-off values for newborn screening blood for galactosemias: “If gal > 60mg/L (0.033mM) or gal1P > 150mg/L (0.058mM).” (Yamaguchi, et al., 1989) |
| glc1p | D-glucose 1-phosphate (M _w 258.1) <u>CHEBI:58601</u> <u>KEGG:C00103</u> | 0.012mM (no galactose) 0.011mM (1h galactose) 0.012mM (1h galactose, GALE inhibition) | [glc1p] = 0.012mM (König, et al., 2012) (Keppler, et al., 1970) [glc1p] = 0.010 ±0.004μmol/g_{ww} (~0.011mM) (starved + galactose 1h, rat, liver) [glc1p] = 0.011 ±0.005μmol/g_{ww} (~0.012mM) (ethanol, starved + galactose 1h, rat, liver) (Guynn, et al., 1974) [glc1p] = 0.0075±0.0010 μmol/g_{ww} (~0.0083mM) (rat liver, starved) [glc1p] = 0.0115±0.008 μmol/g_{ww} (~0.0127mM) (rat liver, fed ad |

| | | | |
|-------|---|--|---|
| | | | libitum) [glc1p] = 0.0132±0.0007 µmol/g _{ww} (~0.0146mM) (rat liver, meal fed) [glc6p]/[glc1p] ~10-12 |
| glc6p | D-glucose 6-phosphate (M _w 258.1) CHEBI:58225 KEGG:C00668 | 0.12mM (no galactose) 0.29mM (1h galactose) 0.30mM (1h galactose, GALE inhibition) | [glc6p] = 0.12mM (König, et al., 2012) (Guynn, et al., 1974) [glc6p] = 0.078±0.011 µmol/g _{ww} (~0.086mM) (rat liver, starved) [glc6p] = 0.147±0.012 µmol/g _{ww} (~0.163mM) (rat liver, fed ad libitum) [glc6p] = 0.157±0.007 µmol/g _{ww} (~0.174mM) (rat liver, meal fed) [glc6p]/[glc1p] ~10-12 (Keppler, et al., 1970) [glc6p] = 0.26 ±0.06µmol/g _{ww} (~0.29mM) (starved + galactose 1h, rat, liver) [glc6p] = 0.30 ±0.13µmol/g _{ww} (~0.33mM) (ethanol, starved + galactose 1h, rat, liver) [glc6p]/[glc1p] = 22.2 ±5.9 (starved + galactose 1h, rat, liver) [glc6p]/[glc1p] = 22.8 ±5.9 (ethanol, starved + galactose 1h, rat, liver) |
| gal1p | D-galactose 1-phosphate (M _w 258.1) CHEBI:58336 KEGG:C00446 | 0.001mM (no galactose) 0.20mM (1h galactose) 0.77mM (1h galactose, GALE inhibition) 1.2mM (GALT deficient, glucose) 5.2mM (GALT deficient, galactose) | (Lai, et al., 2003) (human cells) [gal1p] = ND (not detectable) (Control glucose medium) [gal1p] = 0.2±0.01mM (Control galactose medium) (Keppler, et al., 1970) [gal1p] = 0.18 ±0.04µmol/g _{ww} (~0.2mM)(starved + galactose 1h, rat, liver) [gal1p] = 0.69 ±0.11µmol/g _{ww} (~0.77mM) (ethanol, starved + galactose 1h, rat, liver) (Lai, et al., 2003) (human cells) [gal1p] = 1.2±0.4mM (GALT-deficient glucose medium) [gal1p] = 5.2±0.02mM (GALT-deficient galactose medium) GALT deficiency detected (blood) [gal1p] > 3.0mM (human cells) (Diepenbrock, et al., 1992) GALE deficient patients (blood) (Yamaguchi, et al., 1989) [gal1p]=330-360mg/L (1.28-1.39mM) [gal1p]=474 mg/L (1.84mM) (Yamaguchi, et al., 1989) Neonatal control (blood): |

| | | | |
|---------------|--|--|---|
| | | | gal1P=15±11 mg/L (0.058±0.042mM) (Yamaguchi, et al., 1989) normal values: gal1P = 0.038±0.027 mM (range 0-0.096µM) (Orfanos, et al., 1986) Mean concentration of gal1p (blood) was 0.15mM in cases below the cut-off of 0.74mM (Diepenbrock, et al., 1992) |
| udpglc | UDP-D-glucose (M _w 564.3) <u>CHEBI:58885</u> <u>KEGG:C00029</u> | 0.34mM (no galactose) 0.27mM (1h galactose) 0.17mM (1h galactose, GALE inhibition) | [udpglc] = 0.38mM (König, et al., 2012) [udpglc] = 0.32±0.05 µmol/g_{ww} (~ 0.36mM) (rat liver)(Keppler and Decker, 1969) [udpglc] = 0.26±0.07 µmol/g_{ww} (~ 0.29mM) (rat liver)(Keppler, et al., 1969) (Keppler, et al., 1970) [udpglc] = 0.32 ±0.04µmol/g_{ww} (~ 0.36mM) (fed, rat, liver) [udpglc] = 0.29 ±0.05µmol/g_{ww} (~ 0.32mM) (starved, rat, liver) [udpglc] = 0.24 ±0.09µmol/g_{ww} (~ 0.27mM) (starved + galactose 1h, rat, liver) [udpglc] = 0.15 ±0.03µmol/g_{ww} (~ 0.17mM) (ethanol, starved + galactose 1h, rat, liver) (Guynn, et al., 1974) [udpglc] = 0.342±0.024 µmol/g_{ww} (~ 0.38mM) (rat liver, starved) [udpglc] = 0.433±0.023 µmol/g_{ww} (~ 0.48mM) (rat liver, fed ad libitum) [udpglc] = 0.347±0.027 µmol/g_{ww} (~ 0.39mM) (rat liver, meal fed) (Lai, et al., 2003) (human cells, in µmol/100g(cell protein)) [udpglc] = 236±25 (Control glucose medium) [udpglc] = 179±24 (76% glucose) (Control galactose medium) (Lai, et al., 2003) (human cells, in µmol/100g(cell protein)) [udpglc] = 157±10 (GALT-deficient glucose medium) [udpglc] = 110±10 (70% glucose) (GALT-deficient galactose medium) |
| udpgal | UDP-D-galactose (M _w 564.3) <u>CHEBI:66914</u> <u>KEGG:C00052</u> | 0.11mM (no galactose) 0.36mM (1h galactose) 1.39mM (1h galactose, GALE inhibition) | Both the levels and approximate ratio of 1:3 of udpgal and udpglc are very tightly controlled in normal human cells. (Fridovich-Keil, 2006; Segal, 1995) (1:3 rule udpglc) (Keppler, et al., 1970) [udpgal] = 0.09 ±0.01µmol/g_{ww} (~ 0.10mM) (fed, rat, liver) [udpgal] = 0.09 ±0.01µmol/g_{ww} (~ 0.10mM) (starved, rat, liver) |

| | | | |
|---------------|---|---|--|
| | inhibition) | | <p>[udpgal] = 0.32 ± 0.07 μmol/g_{ww} (~0.36mM) (starved + galactose 1h, rat, liver)</p> <p>[udpgal] = 1.25 ± 0.16 μmol/g_{ww} (~1.39mM) (ethanol, starved + galactose 1h, rat, liver)</p> <p>(Keppler, et al., 1970)</p> <p>[udpgal]/[udpglc] = 3.4 ± 0.3 (fed, rat, liver)</p> <p>[udpgal]/[udpglc] = 3.3 ± 0.3 (starved, rat, liver)</p> <p>[udpgal]/[udpglc] = 0.78 ± 0.39 (starved + galactose 1h, rat, liver)</p> <p>[udpgal]/[udpglc] = 0.11 ± 0.02 (ethanol, starved + galactose 1h, rat, liver)</p> <p>[udpgal]/[gal1p] = 1.94 ± 0.35 (starved + galactose 1h, rat, liver)</p> <p>[udpgal]/[gal1p] = 1.85 ± 0.27 (ethanol, starved + galactose 1h, rat, liver)</p> <p>(Lai, et al., 2003) (human cells, in μmol/100g(cell protein))</p> <p>[udpgal] = 82±10 (Control glucose medium)</p> <p>[udpgal] = 46±4 (56% glucose) (Control galactose medium 24h)</p> <p>(Lai, et al., 2003) (human cells, in μmol/100g(cell protein))</p> <p>[udpgal] = 25±5 (GALT-deficient glucose medium)</p> <p>[udpgal] = 17±3 (68% glucose) (GALT-deficient galactose medium 24h)</p> |
| galtol | D-galactitol (M _w 182.2) <u>CHEBI:16813</u> <u>KEGG:C01697</u> | 0.001mM (no galactose) ~8mM (GALT deficiency) | <p>[galtol] = 4.8-40 μmol/g (~5.3-44mM) (occupational gray matter, human)</p> <p>[galtol] = 17.6 μmol/g (~) (basal ganglia, human) (Wang, et al., 2001)</p> <p>[galtol] = 12.9 μmol/g (~14.3mM) (Wang, et al., 2001) (Wells, et al., 1965)</p> <p>[galtol] = 22.18 μmol/g (~24.6mM) (Wang, et al., 2001) (Quan-Ma, et al., 1966)</p> <p>Galactitol measured directly in GALT-deficient mice are lower (2mM) than levels detected by MRS in human subjects (8mM) (Leslie, 2003; Wang, et al., 2001)</p> |
| atp | ATP (M _w 503.2) <u>CHEBI:30616</u> <u>KEGG:C00002</u> | 2.7mM (no galactose) 2.9mM (1h galactose) 2.9mM (1h galactose, GALE inhibition) | <p>[atp] = 2.8mM (König, et al., 2012)</p> <p>(Guynn, et al., 1974)</p> <p>[atp] = 2.49±0.12 μmol/g_{ww} (~2.77mM) (rat liver, starved)</p> <p>[atp] = 2.56±0.09 μmol/g_{ww} (~2.84mM) (rat liver, fed ad libitum)</p> <p>[atp] = 2.32±0.07 μmol/g_{ww} (~2.58mM) (rat liver, meal fed)</p> <p>[atp] = 2.42±0.50 μmol/g_{ww} (~2.69mM) (rat liver) (Keppler, et al.,</p> |

| | | | |
|-----|---|---|--|
| | | | 1969) (Keppler, et al., 1970) [atp] = 2.60 ± 0.16 μmol/g_{ww} (~2.89mM) (starved + galactose 1h, rat, liver) [atp] = 2.81 ± 0.15 μmol/g_{ww} (~3.12mM) (ethanol, starved + galactose 1h, rat, liver) [atp]/[adp] = 3.14 ± 0.52 (starved + galactose 1h, rat, liver) [atp]/[adp] = 3.10 ± 0.53 (ethanol, starved + galactose 1h, rat, liver) |
| adp | ADP (M _w 424.2) CHEBI:456216 KEGG:C00008 | 1.2mM (no galactose) 1.0mM (1h galactose) 1.0mM (1h galactose, GALE inhibition) | [atp] = 0.8mM (König, et al., 2012) (Guynn, et al., 1974) [adp] = 1.38 ± 0.08 μmol/g_{ww} (~1.53mM) (rat liver, starved) [adp] = 1.06 ± 0.03 μmol/g_{ww} (~1.18mM) (rat liver, fed ad libitum) [adp] = 1.24 ± 0.04 μmol/g_{ww} (~1.38mM) (rat liver, meal fed) [adp] = 1.08 ± 0.12 μmol/g_{ww} (~1.20mM) (rat liver) (Keppler, et al., 1969) (Keppler, et al., 1970) [adp] = 0.88 ± 0.17 μmol/g_{ww} (~0.98mM) (starved + galactose 1h, rat, liver) [adp] = 0.97 ± 0.19 μmol/g_{ww} (~1.08mM) (ethanol, starved + galactose 1h, rat, liver) |
| utp | UTP (M _w 480.1) CHEBI:46398 KEGG:C00075 | 0.27mM (no galactose) | [utp] = 0.27mM (König, et al., 2012) (Guynn, et al., 1974) [utp] = 0.362 ± 0.014 μmol/g_{ww} (~0.40mM) (rat liver, starved) [utp] = 0.494 ± 0.038 μmol/g_{ww} (~0.55mM) (rat liver, fed ad libitum) [utp] = 0.443 ± 0.039 μmol/g_{ww} (~0.49mM) (rat liver, meal fed) |
| udp | UDP (M _w 401.1) CHEBI:58223 KEGG:C00015 | 0.09mM (no galactose) | [udp] = 0.09mM (König, et al., 2012) [utp+udp] = 0.35 ± 0.07 μmol/g_{ww} (~0.39mM) (rat liver) (Keppler, et al., 1969) [utp+udp] = 0.35 ± 0.05 μmol/g_{ww} (~0.39mM) (rat liver) (Keppler and Decker, 1969) (Keppler, et al., 1970) [utp+udp] = 0.34 ± 0.05 μmol/g_{ww} (~0.38mM) (fed, rat, liver) [utp+udp] = 0.23 ± 0.05 μmol/g_{ww} (~0.26mM) (starved, rat, liver) [utp+udp] = 0.15 ± 0.03 μmol/g_{ww} (~0.17mM) (starved + galactose 1h, rat, liver) [utp+udp] = 0.11 ± 0.02 μmol/g_{ww} (~0.39mM) (ethanol, starved + galactose 1h, rat, liver) |

| | | | |
|--------|---|------------------------------|---|
| | | | Marked decrease in [utp+udp] under galactose challenge. |
| phos | Phosphate (M _w 96.0) CHEBI:43474 KEGG:C00009 | 5.0mM (König, et al., 2012) | [pi] = 5.0mM (König, et al., 2012) (Guynn, et al., 1974) [pi] = 4.37±0.16 µmol/g _{ww} (~4.86mM) (rat liver, starved) [pi] = 3.64±0.32 µmol/g _{ww} (~4.04mM) (rat liver, fed ad libitum) [pi] = 4.41±0.10 µmol/g _{ww} (~4.90mM) (rat liver, meal fed) [pi] = 3.18±0.56 µmol/g _{ww} (~3.53mM) (rat liver)(Keppler and Decker, 1969) |
| ppi | Pyrophosphate (M _w 175.0) CHEBI:33019 KEGG:C00013 | 0.008mM(König, et al., 2012) | [pp] = 0.008mM (König, et al., 2012) (Guynn, et al., 1974) [pp] = 0.0023±0.0003 µmol/g _{ww} (~0.0026mM) (rat liver, starved) [pp] = 0.0038±0.0004 µmol/g _{ww} (~0.0042mM) (rat liver, fed ad libitum) [pp] = 0.0049±0.0006 µmol/g _{ww} (~0.0054mM) (rat liver, meal fed) [pp] = 0.0065±0.00086 µmol/g _{ww} (~0.0072mM) (rat total liver) |
| nadp | NADP (M _w 740.4) CHEBI:58349 KEGG:C00006 | 0.1mM | |
| nadph | NADPH (M _w 741.4) CHEBI:57783 KEGG:C00005 | 0.1mM | |
| suc | Sucrose (M _w 342.3) CHEBI:17992 KEGG:C00089 | | |
| h2oM | H2O M CHEBI:15377 KEGG:C00001 | | |
| alb | albumin PR:000003918 | | |
| rbc | red blood cell BTO:0000424 | | |
| galnat | D-galactonate (M _w 195.1) CHEBI:12931 | | |

| KEGG:C00880 | | |
|-------------|---------------|---|
| galn | galactosamine | Uptake of galactosamine by rat liver is a~0.4μmol/g(liver)/min as measured by the disappearance of galactosamine from the medium (Keppler, et al., 1969) Time-dependent decrease in uridine nucleotides in isolated perfused rat livers after galactosamine addition. (Keppler, et al., 1969) |
| amp | AMP | [amp] = 0.28±0.06 μmol/g _{ww} (~0.31mM) (rat liver) (Keppler, et al., 1969) (Keppler, et al., 1970) [amp] =0.15 ±0.09μmol/g _{ww} (~0.167mM) (starved + galactose 1h, rat, liver) [amp] =0.19 ±0.07μmol/g _{ww} (~0.21mM) (ethanol, starved + galactose 1h, rat, liver) |
| ump | UMP | [ump] = 0.04 μmol/g _{ww} (~0.044mM) (rat liver) (Segal and Rogers, 1971) |

Supplementary Table 3 - Tissue-and organ parameters

| Parameter | Symbol | Model value | Human | Dog | Rat |
|---|------------|--|---|-----------------------|--|
| number of hepatocytes along sinusoid | N_c | 20 | 15-25 (human) (Kuntz and Kuntz, 2006) | | 12-20 (rat, from image) (Burkel and Low, 1966) |
| sinusoid length | L_{sin} | 500μm (±125μm) | 500-650μm diameter of hepatic lobules 1.0–1.3mm (Kuntz and Kuntz, 2006)) 500μm (distance between central veins 1000μm) (Lautt, 2009) 350–500μm (Kuntz and Kuntz, 2006) | 500μm (Goresky, 1963) | 400-450μm distance between central veins 809±199μm (SD, n=79, young rat, SEM of corrosion cast)(Warren, et al., 2008) 891±190μm (n=78, old rat, SEM of corrosion cast)(Warren, et al., 2008) |
| Diameter hepatocyte in sinusoidal direction | x_{cell} | L_{sin} / N_c 25μm (calculated from geometry) | 20 – 40μm (Kuntz and Kuntz, 2006) | | No significant difference could be shown in the average size of parenchymal cells among the lobular zones of rat liver. 20.8±0.2μm (SD, n=50, rat, periportal, QSD) (Loud, 1968) 20.8±0.3μm (SD, n=50, rat, midzonal, QSD) (Loud, 1968) 21.0±0.3μm (SD, n=50, rat, perivenious, QSD) (Loud, 1968) Parenchymal cells of normal rat liver are at least 80% homogeneous with respect to the structural parameters measured. 20.1μm (rat, QSD) (Wiener, et al., 1968) |
| sinusoidal radius | y_{sin} | 4.4μm (8.8μm sinusoidal diameter) | Sinusoidal diameter 8.8±0.9μm (human, OPS) (Puhl, et al., 2003) 4-15μm (human) (Kuntz and Kuntz, 2006) 13.23±2.36μm (human, n=100, SEM) (Debbaut, et al., 2014) | | Sinusoidal diameter 5.9±0.17μm (SE, n=545, rat, periportal, IVM) (Wisse, et al., 1985) 7.1±0.29μm (SE, n=498, rat, central, IVM) (Wisse, et al., 1985) 6.42±0.12μm (SE, n=696, rat, periportal, IPM) (Wisse, et al., 1985) 7.62±NDμm (SE, n=696, rat, central, IPM) (Wisse, et al., 1985) 5.9±0.17μm (rat, Zone 1), 7.1±0.29μm (rat, Zone 3) (MacPhee, et al., 1995) 6.4±0.1μm (rat, Zone 1), 8.3±0.2μm (rat, Zone 3) (MacPhee, et al., 1995) 6.6±0.09μm (SEM, n=139, rat, direct sinusoids)(Koo, |

Commented [WU2]: TODO: Update table with final calculated values for model, i.e relative Volumes, and the number of sinusoids (scaling flow and volume)

et al., 1975)
6.3±0.07µm (n=304, rat, branching sinusoids, SEM) (Koo, et al., 1975)
6.3±0.12µm (n=72, rat, direct sinusoids, SEM) (Koo, et al., 1975)
9.4±3.6µm (SD, n=977, young rat, SEM of corrosion cast)(Warren, et al., 2008)
9.7±3.5µm (n=1225, old rat, SEM of corrosion cast)(Warren, et al., 2008)

0.3-1.5µm (rat, SEM micrograph, estimated from image) (Burkel and Low, 1966)
0.2-1µm (rat, SEM micrograph, estimated from image) (Wisse, et al., 1985)
0.3-1.2µm (rat, TEM, estimated from image (Braet and Wisse, 2002)

intersinusoidal distance
~**15-30µm** (SEM, estimated from image)(Wisse, et al., 1985)

16.1±3.9µm (SD, n=567, young rat, SEM of corrosion cast)(Warren, et al., 2008)
15.5±3.8µm (SD, n=558, old rat, SEM of corrosion cast)(Warren, et al., 2008)

width space of Disse y_{dis} **1.2±0.4µm**
0.4-1.5µm (human, SEM, estimated from imaged) (Muto, et al., 1977)
The sinusoidal lining of human liver appeared remarkably similar to that of the rat by both TEM and SEM. (Burwen, et al., 1982)
0.5-1.7µm (human, SEM, estimated from image) (Burwen, et al., 1982)

cell sheet thickness y_{cell} **7.58±1.25µm** (calculated from FSD with geometry)
7.58µm
Calculated from functional sinusoidal density FSD
FSD **391±30** [1/cm] (SD, n=88, human, OPS) (Puhl, et al., 2003)

$$y_{cell} = \frac{1}{2 \cdot FSD} - (y_{sin} + y_{dis})$$

6.1±1.25µm
Calculated from intersinusoidal distance ISD

$$y_{cell} = \frac{ISD}{2} - (y_{sin} + y_{dis})$$

ISD **22.6±2.5µm** (SD, n=88, human, OPS) (Puhl, et al., 2003)

Area between adjacent sinusoid compartments A_{sin} $\pi(y_{sin})^2$

Area between adjacent Disse compartments A_{dis} $\pi(y_{sin} + y_{dis})^2 - A_{sin}$

Area between adjacent sinusoid and Disse $A_{sin\ dis}$ $2\pi \cdot y_{sin} \cdot x_{sin}$

| | | | | |
|---|---------------|--|---|---|
| compartments | | | | |
| Volume sinusoid compartment | V_{sin} | $A_{sin} \cdot x_{sin}$ | | |
| Volume Disse compartment | V_{dis} | $A_{dis} \cdot x_{sin}$ | | |
| Volume cell | V_{cell} | $\pi(y_{sin} + y_{dis} + y_{cell})^2 \cdot x_{cell}$ $-\pi(y_{sin} + y_{dis})^2 x_{cell}$ | | Volumes cytosol calculated per average cell, i.e. per nucleus 5100μm^3 (peripheral, rat, QSD) (Loud, 1968) 5100μm^3 (midzonal, rat, QSD) (Loud, 1968) 5100μm^3 (perivenousl, rat, QSD) (Loud, 1968) No significant difference could be shown in the average size of parenchymal cells among the lobular zones of rat liver. 5100μm^3 (all zones, rat, QSD) (Wiener, et al., 1968) Cell Volumes ~1.4*5100 μm^3 = 7140 μm^3 |
| Volume sinusoidal unit | $V_{sinunit}$ | $L_{sin} \cdot \pi \cdot (y_{sin} + y_{dis} + y_{cell})^2$ 272.9E3 μm^3 (calculated from geometry) | | |
| Volume fraction sinusoidal blood volume, % liver | f_{sin} | $\frac{V_{sin}}{V_{sin} + V_{dis} + V_{cell}}$ 11.1% (calculated from geometry) | 15–25% (percent of liver volume) (Kuntz and Kuntz, 2006) 9-15% (n=6, isolated perfused human liver)(Villeneuve, et al., 1996) | 15.2% (indicator dilution dog) (Goresky, 1963) 15.0% (dog) (Allen and Reeve, 1953; Goresky, 1963) 19.4% (rat) (Everett, et al., 1956; Goresky, 1963) 11.6% (rat) (Brauer, et al., 1959; Goresky, 1963) 10.6% (morphological studies, % volume) (Blouin, et al., 1977; Lutt, 2009) |
| Volume fraction extravascular volumes, % liver | f_{dis} | $\frac{V_{dis}}{V_{sin} + V_{dis} + V_{cell}}$ 6.9% (calculated from geometry) | ~5% (percent of liver volume) (Kuntz and Kuntz, 2006) 5-8% (n=6, isolated perfused human liver)(Villeneuve, et al., 1996) | 6.2% (indicator dilution dog) (Goresky, 1963) 9.5±2.1% (±SD, indicator dilution dog, sucrose volume) (Goresky, 1963) 6.7% (dog) (Allen and Reeve, 1953; Goresky, 1963) 7.3% (rat) (Goresky, 1963) 6.0% (rat) (Brauer, et al., 1959; Goresky, 1963) 4.9% (morphological studies, % volume) (Blouin, et al., 1977; Lutt, 2009) |
| Volume fraction parenchymal cells, % liver | f_{hep} | $\frac{V_{hep}}{V_{sin} + V_{dis} + V_{cell}}$ 81.9% (calculated from geometry) | | 78% (morphological studies, % volume) (Blouin, et al., 1977; Lutt, 2009) |

| | | | | | |
|--|---------------|---|--|---|--|
| RBC velocity | v_{RBC} | 270μm/s\pm58μm/s (mode 180μm/s) | <p>970\pm430μm/s (SD; human, OPS) (Puhl, et al., 2003) Values probably too high due to larger arterial contribution with high flow on organ surface. [TODO: Ref & discussion] 259μm/s (boundary condition, calculated from scaling to tissue sample, human) (Debbaut, et al., 2012) The mean flow velocities of simulations in model of human liver microcirculation based on a 3D image-based geometry were for the r, phi and z directions 100μm/s, 73μm/s and 77μm/s, respectively. (Debbaut, et al., 2012) The sinusoidal circulation is clearly anisotropic. Modeling in human corrosion cast showed that pressure drops significantly through certain sinusoids and stays almost constant through others. This corresponds to the typical presence of preferential pathways in the sinusoidal geometry along tracks with relevant pressure drops (streamlines). (Debbaut, et al., 2012)</p> | <p>93μm/s (dog, calculated from transit time of RBC) (Goresky, 1963) [CHECK, depends on injection time]</p> | <p>180\pm20μm/s (SE, rat) (MacPhee, et al., 1988) 250\pm3μm/s (SE, rat, IVM) (Koo and Liang, 1979) 150\pm6μm/s (SE, rat, stated in (MacPhee, et al., 1988), video flying spot method) 69.2\pm30.6μm/s (\pmSD, mice, IVM) (MacPhee, et al., 1988) 410\pm39μm/s (SEM, n=139, rat, direct sinusoids)(Koo, et al., 1975) 270\pm58μm/s (SEM, n=304, rat, branching sinusoids)(Koo, et al., 1975) 370\pm25μm/s (SEM, n=72, rat, direct sinusoids)(Koo, et al., 1975) Analysis of the blood cell velocity data of Koo as a cumulative distribution showed that the data is fitted better by a log-normal than a normal distribution (Roberts and Rowland, 1985).</p> |
| volumetric blood flow sinusoidal unit | $Q_{sinunit}$ | $\pi(y_{sin})^2 v_{RBC}$ $16.4E3 \frac{\mu m^3}{s}$ (calculated from geometry) | flow through cylinder analogue to (Gross and Aroesty, 1972; Puhl, et al., 2003) | | |
| fenestraction parameter The presence of fenestrae reduces the surface available for free | f_{fen} | 0.09 | <p>107\pm1.5nm (SE, human) (Wisse, et al., 2008)</p> <p>diameter fenestrae</p> | <p>(Wisse, et al., 1985) demonstrated presence of</p> | <p>diameter fenestrae 175nm (Wisse, et al., 1996) 161\pm2.7nm (Sprague-Dawley rats) (Wisse, et al., 2008)</p> |

| | | | |
|---|---|---|--|
| transport, whereas the parenchymal surface available for uptake is 6.0 times enlarged by microvilli (Schaff and Lapis, 1990; Wisse, et al., 1985) | <p>50-300nm (SEM, human)(Braet and Wisse, 2002; Horn, et al., 1987)</p> <p>No difference in ultrastructural morphology was seen between Zones 3 and 1 (Horn, et al., 1987)</p> <p>frequency (SEM, human) (Braet and Wisse, 2002; Horn, et al., 1987) 23.5 [15 – 25] [1/μm²] (n=13, SEM, human, Zone3) (Horn, et al., 1987) 19 [10-24] [1/μm²] (n=10, SEM, human, Zone1) (Horn, et al., 1987)</p> <p>porosity 9.3% [4.8-16.2] (n=13, Zone3, SEM, human) (Horn, et al., 1987) 7.6% [3.8-12.3] (n=10, Zone1, SEM, human) (Horn, et al., 1987)</p> | fenestrae in dog liver with preliminary measurements indicating that the size distribution was almost equal to rat liver fenestrae. | <p>174.6±1.0nm (SE, rat, periportal, TEM) (Wisse, et al., 1985) 147.2±0.9nm (SE, rat, pericentral, TEM) (Wisse, et al., 1985) SEM preparation causes significant shrinkage at the level of fenestrae (approximately 30%)!</p> <p>110.7±0.25nm (SE, rat, periportal, SEM) (Wisse, et al., 1985) 104.8±0.22nm (SE, rat, pericentral, SEM) (Wisse, et al., 1985) 98.0±13.0nm (SD, n=3, SEM, rat) (Fraser, et al., 1988)</p> <p>frequency 9 per μm (SE, rat, periportal, SEM) (Wisse, et al., 1985) 13 [1/μm²] (SE, rat, pericentral, SEM) (Wisse, et al., 1985) 9.08 – 13.3 [1/μm²] (SE, rat) (Wisse, et al., 1985) 20.0±6.3[1/μm²] (SD, n=3, SEM, rat) (Fraser, et al., 1988)</p> <p>porosity 6-8% (Wisse, et al., 1996) A lobular gradient of decreasing fenestrae diameter is compensated by an inverse gradient of fenestrae number. (Wisse, et al., 1996) Only a limited surface of the lining is available for free exchange (~10%) (Wisse, et al., 1996) 17.6±6.9 (SD, n=3, SEM, rat) (Fraser, et al., 1988)</p> |
|---|---|---|--|

Supplementary Table 4 – Parameters for the log-normal distributions

| Parameter | | meanlog | stdlog | mean μ (reported) | standard deviation (reported) | SD references |
|----------------------------|------------|----------------------|------------------------|--------------------------|-------------------------------------|--|
| Sinusoidal length | L_{sin} | 6.184 | 0.2462 | 500 μ m | 125 μ m | based on distance between central veins 809\pm199μm (SD, n=79, young rat, SEM of corrosion cast) (Warren, et al., 2008) scaled to human sinusoidal length Based on distribution of sinusoidal diameter 8.8\pm0.9μm (SD, n=440 in N=11 human, OPS) (Puhl, et al., 2003) |
| Sinusoidal radius | y_{sin} | 1.465 (\pm 0.010) | 0.1017 (\pm 0.0073) | 4.4 μ m | 0.45 μ m | 0.4-1.5μm (human, SEM, estimated from imaged) (Muto, et al., 1977) |
| Width of Disse space | y_{dis} | 0.1296 | 0.3246 | 1.2 μ m | 0.4 μ m | 0.5-1.2μm (human, SEM, estimated from image) (Burwen, et al., 1982) |
| Hepatocyte sheet thickness | y_{cell} | 1.977 (\pm 0.014) | 0.1390 (\pm 0.0099) | 7.58 μ m | 1.25 μ m | 7.58μm Calculated from functional sinusoidal density FSD FSD 391\pm30 [1/cm] (SD, n=88, human, OPS) (Puhl, et al., 2003) $y_{cell} = \frac{1}{2 \cdot FSD} - (y_{sin} + y_{dis})$ 6.1\pm1.25μm Calculated from intersinusoidal distance ISD $y_{cell} = \frac{ISD}{2} - (y_{sin} + y_{dis})$ ISD 22.6\pm2.5μm (SD, n=88, human, OPS) (Puhl, et al., 2003) |
| RBC flow velocity | v_{RBC} | 5.457 (0.0267) | 0.6178 (0.0189) | 270 μ m/s | 58 μ m/s | 270\pm58μm/s (SEM, n=304, rat, branching sinusoids)(Koo, et al., 1975) |

Model fit

| | name | mean | std | unit | meanlog | meanlog_error | sdlog | sdlog_error | scale_fac | scale_unit |
|----------|----------|------------|------------|------|--------------|---------------|--------------|--------------|-----------|------------|
| L | L | 0.0005 | 0.000125 | m | 6.1842957875 | NA | 0.2462206771 | NA | 1000000 | μ m |
| y_sin | y_sin | 0.0000044 | 0.00000045 | m | 1.4652733102 | 0.0102747149 | 0.1017144881 | 0.0072653206 | 1000000 | μ m |
| y_dis | y_dis | 0.0000012 | 0.00000004 | m | 0.129641299 | NA | 0.324592846 | NA | 1000000 | μ m |
| y_cell | y_cell | 0.00000758 | 0.00000125 | m | 1.9769003149 | 0.0140416505 | 0.1390052478 | 0.0099289463 | 1000000 | μ m |
| flow_sin | flow_sin | 0.00027 | 0.000058 | m/s | 5.4572075437 | 0.0267357281 | 0.6178209697 | 0.0189050147 | 1000000 | μ m/s |

Commented [WU3]: TODO: remove
Calculated fit table to make sure the values are correct

Supplementary Table 5 – Diffusion coefficients

| Diffusion coefficients | Symbol | Model value [$\mu\text{m}^2/\text{s}$] | References [$\mu\text{m}^2/\text{s}$] |
|------------------------|-----------|--|---|
| water | D^{h2o} | 2200 | 2100 (water in water) Bionumbers id=104087, ver=7)(Milo, et al., 2010) 2300 (water in water) Bionumbers id=104087, ver=3)(Milo, et al., 2010) |
| sucrose | D^{suc} | 720 | 720 (D37, free diffusion coefficient in water at 37°C)(Renkin, 1977) 520 (sucrose in water) (Bionumbers id=100614, ver=7)(Milo, et al., 2010) |
| glucose | D^{glc} | 910 | 910 (D37, hexose, free diffusion coefficient in water at 37°C)(Renkin, 1977) (Casciari, et al., 1988; Groebe, et al., 1994) 600 (glucose in water) (Bionumbers id=104089, ver=6)(Milo, et al., 2010) 673 (glucose in water) (Bionumbers id=109504, ver=1)(Milo, et al., 2010) |
| galactose | D^{gal} | 910 | 910 (D37, hexose, free diffusion coefficient in water at 37°C)(Renkin, 1977) |
| albumin | D^{alb} | 90 | 90 (D37, free diffusion coefficient in water at 37°C)(Renkin, 1977) |

Supplementary Table 6 – Organ/Liver parameters

| Parameter | Symbol | Model value | Human | Dog | Rat |
|--|--------------|------------------|--|---|---|
| total liver weight | m_{liv} | 1500g | 1500-1800g (man), 1300-1500g (woman) (Kuntz and Kuntz, 2006) 1697±171g (±SD, n=6)(Villeneuve, et al., 1996) 2.5% of body weight (Vollmar and Menger, 2009) | 556g (400 – 800g) (dog) (Goresky, 1963) | 17.1±2.2g (±SD. N=13, in situ perfused rat livers)(Garipey, et al., 1993) |
| density liver tissue | ρ_{liv} | $1 \frac{g}{ml}$ | $1 \frac{g}{ml}$ (Debbaut, et al., 2012) Literature (data from metabolite conversions ~1.2g/ml) | | |
| total liver volume | V_{liv} | 1500ml | 1500ml (calculated by $\frac{m_{liv}}{\rho_{liv}}$) measured by ultrasonography 1524ml (man, 24 years) (Wynne, et al., 1989) 1102ml (man, 91 years, -28%) (Wynne, et al., 1989) 1415ml (woman, 24 years) (Wynne, et al., 1989) 789ml (woman, 91 years, -44%) (Wynne, et al., 1989) 1474ml (all, 24 years) (Wynne, et al., 1989) 934ml (all, 91 years, -37%) (Wynne, et al., 1989) volumes per bodyweight (ultrasonography) 20.7 ml/kgbw (man, 24 years) (Wynne, et al., 1989) 14.5 ml/kgbw (man, 91 years, -30%) (Wynne, et al., 1989) 23 ml/kgbw (woman, 24 years) (Wynne, et al., 1989) 13.6 ml/kgbw (woman, 91 years, -43%) (Wynne, et al., 1989) 23.6ml/kgbw (all, 24 years) (Wynne, et al., 1989) 14.0 ml/kgbw (all, 91 years, -41%) (Wynne, et al., 1989) | 556ml (calculated) | 17.1ml (calculated) |
| parenchymal tissue fraction of liver (due to large vessel, connective tissue, lymphs system, | f_{tissue} | 0.5 | Tissue volume of liver is calculated by $V_{tissue} = f_{tissue} \cdot V_{liv}$ Literature (25 % large vessels, ...) | | |

only part of whole liver
volume is parenchymal
tissue)

| | | | | | |
|--|-----------|-----------------------------------|--|--|--|
| total hepatic blood flow per liver weight (~75-80% portal vein partially deoxygenated, 20-25% hepatic artery well-oxygenated) | q_{liv} | $1.2 \frac{ml}{min \cdot g_{LW}}$ | $\sim 1.0\text{--}1.3 \frac{ml}{min \cdot g_{LW}}$ (Lautt, 2009) $\sim 1 \frac{ml}{min \cdot g_{LW}}$ (Vollmar and Menger, 2009) (Kuntz and Kuntz, 2006) measured by ultrasonography 1.24 ml/min/gLW (man, 24 years) (Wynne, et al., 1989) 1.02 ml/min/gLW (man, 91 years, -18%) (Wynne, et al., 1989) 1.11 ml/min/gLW (woman, 24 years) (Wynne, et al., 1989) 0.88 ml/min/gLW (woman, 91 years, -21%) (Wynne, et al., 1989) 1.18 ml/min/gLW (all, 24 years) (Wynne, et al., 1989) 0.94 ml/min/gLW (all, 91 years, -20%) (Wynne, et al., 1989) | 1.83±0.55 $\frac{ml}{min \cdot g_{LW}}$ (±SD, g liver weight, dog) (Goresky, 1963) | $1.30 \pm 0.13 \frac{ml}{min \cdot g_{LW}}$ (SD, rat 3 month, determined by clearance of albumin)(Warren, et al., 2008) $1.54 \pm 0.19 \frac{ml}{min \cdot g_{LW}}$ (SD, rat 6 month, determined by clearance of albumin)(Warren, et al., 2008) $1.33 \pm 0.28 \frac{ml}{min \cdot g_{LW}}$ (SD, rat 36 month, determined by clearance of albumin)(Warren, et al., 2008) |
| total hepatic blood flow per body weight | | | measured by ultrasonography 25.3 ml/min/kgBW (man, 24 years) (Wynne, et al., 1989) 14.5 ml/min/kgBW (man, 91 years, -43%) (Wynne, et al., 1989) 25.5 ml/min/kgBW (woman, 24 years) (Wynne, et al., 1989) 11.5 ml/min/kgBW (woman, 91 years, -55%) (Wynne, et al., 1989) 25.7 ml/min/kgBW (all, 24 years) (Wynne, et al., 1989) 13.5 ml/min/kgBW (all, 91 years, -47%) (Wynne, et al., 1989) $17.0 \pm 2.72 \frac{ml}{min \cdot kg_{BW}}$ (±SD, n=10 women, Doppler ultrasound)(Carlisle, et al., 1992) $30 \frac{ml}{min \cdot kg_{BW}}$ (Lautt, 2009) | | |

| | | | | | |
|-----------------------------------|-----------|---|---|--|--|
| total hepatic blood flow | Q_{liv} | $q_{liv} \cdot m_{liv}$ $1800 \frac{ml}{min}$ (calculated) | <p>1800 ml/min (man) (Kuntz and Kuntz, 2006)</p> <p>1200 ml/min (woman) (Kuntz and Kuntz, 2006)</p> <p>measured by ultrasonography</p> <p>1864 ml/min (man, 24 years) (Wynne, et al., 1989)</p> <p>1126 ml/min (man, 91 years, -40%) (Wynne, et al., 1989)</p> <p>1546 ml/min (woman, 24 years) (Wynne, et al., 1989)</p> <p>645 ml/min (woman, 91 years, -58%) (Wynne, et al., 1989)</p> <p>1717 ml/min (all, 24 years) (Wynne, et al., 1989)</p> <p>807 ml/min (all, 91 years, -53%) (Wynne, et al., 1989)</p> <p>1067±160 ml/min(±SD, n=6, isolated perfused human liver)(Villeneuve, et al., 1996)</p> <p>992±276 ml/min (n=14)(Jakab, et al., 1995)</p> | 869 ml/min (dog) (Goresky, 1963) | 20.9±1.3 ml/min (±SD, N=13, in situ perfused rat livers, perfusate blood flow)(Garipey, et al., 1993) |
| total number of hepatic sinusoids | N_{sin} | | <p>Calculate based on flow</p> $N_{sin} = \frac{Q_{liv}}{Q_{sin unit}} = \frac{Q_{liv}}{A_{sin} \cdot v_{RBC}} = \frac{Q_{liv}}{\pi (y_{sin})^2 \cdot v_{RBC}}$ <p>$N_{sin} = 2472E6$ (calculated) update</p> <p>Calculate based on volumes</p> $N_{sin} = \frac{V_{tissue}}{V_{sin unit}} = \frac{f_{tissue} \cdot V_{liv}}{\pi \cdot (y_{sin} + y_{dis} + y_{cell})^2 \cdot L_{sin}}$ <p>$N_{sin} = 3250E6$ (calculated) update</p> <p>With the number of hepatic lobuli:</p> <p>$N_{Lob} = 1.0E6-1.5E6$ (Kuntz and Kuntz, 2006)</p> <p>Comes this to an estimated number of sinusoids per lobules of ~2600 sinusoidal units per lobules</p> | | |
| mean transit time RBC | MTT | ~3-4s | Calculate from curves [Villeneuve1996] | Calculate from curves [Goresky1973, Goresky1983] | ~6.3±0.5sec RBC ??(Garipey, et al., 1993) Calculate from curves |
| large vessel transit time | t_0 | ~5-10s | Calculate from curves [Villeneuve1996] | Calculate from curves [Goresky1973, Goresky1983] | Calculate from curves |

Commented [WU4]: Check in Dilution curves what the real transit times are

SUPPLEMENTARY FIGURES

Supplementary Figure 1 – Validation of single cell model of galactose metabolism time courses in GALE inhibition via ethanol.




Fig. 1. Time-dependent changes of galactose metabolite and uracil nucleotide contents in livers of fed rats. D-galactose (2.7 mmole/kg) was injected at zero time, one hr after ethanol administration (130 mmole/kg). Uridine phosphates (UTP + UDP + UMP) are designated as UP, the sum of all acid soluble uracil 5'-nucleotides as ΣUMP. (Keppler, et al., 1970)

Typical effect of galactose challenge on the galactose metabolism of single cell model. Prediction of effects of GALE inhibition. GALE is inhibited by ethanol ingestion, due to altered NAD/NADH quotient resulting in altered GALE activity due to NAD cofactor. Ethanol ingestion was simulated by altering the NAD levels. Experimental data from (Keppler, et al., 1970)

Commented [WU5]: TODO: calculate the GALE inhibition, discuss in context of alcohol effects to the liver

Supplementary Figure 2 – Metabolic Control Analysis (MCA) - Role of the different parameters in controlling hepatic galactose metabolism on a single cell level

Supplementary Figure 2 – Multiple indicator dilution curves in galactosemias

Supplementary Figure 3 – Altered hepatic galactose clearance and dependency on flow in galactosemias analogue to Figure 4.

REFERENCES

- Accorsi, A., *et al.* (1989) Isoenzymes of phosphoglucomutase from human red blood cells: isolation and kinetic properties, *Preparative biochemistry*, **19**, 251-271.
- Allen, T.H. and Reeve, E.B. (1953) Distribution of extra plasma in the blood of some tissues in the dog of measured with P32 and T-1824, *The American journal of physiology*, **175**, 218-223.
- Arbuckle, M.I., *et al.* (1996) Structure-function analysis of liver-type (GLUT2) and brain-type (GLUT3) glucose transporters: expression of chimeric transporters in *Xenopus* oocytes suggests an important role for putative transmembrane helix 7 in determining substrate selectivity, *Biochemistry*, **35**, 16519-16527.
- Bautista, J.M., Mason, P.J. and Luzzatto, L. (1992) Purification and properties of human glucose-6-phosphate dehydrogenase made in *E. coli*, *Biochimica et biophysica acta*, **1119**, 74-80.
- Blouin, A., Bolender, R.P. and Weibel, E.R. (1977) Distribution of organelles and membranes between hepatocytes and nonhepatocytes in the rat liver parenchyma. A stereological study, *The Journal of cell biology*, **72**, 441-455.
- Braet, F. and Wisse, E. (2002) Structural and functional aspects of liver sinusoidal endothelial cell fenestrae: a review, *Comp Hepatol*, **1**, 1.
- Brauer, R.W., Holloway, R.J. and Leong, G.F. (1959) Changes in liver function and structure due to experimental passive congestion under controlled hepatic vein pressures, *The American journal of physiology*, **197**, 681-692.
- Brown, G.K. (2000) Glucose transporters: structure, function and consequences of deficiency, *Journal of inherited metabolic disease*, **23**, 237-246.
- Burkel, W.E. and Low, F.N. (1966) The fine structure of rat liver sinusoids, space of Disse and associated tissue space, *Am J Anat*, **118**, 769-783.
- Burwen, S.J., *et al.* (1982) The perfused human liver wedge biopsy: a new in vitro model for morphological and functional studies, *Hepatology*, **2**, 426-432.
- Carlisle, K.M., *et al.* (1992) Estimation of total hepatic blood flow by duplex ultrasound, *Gut*, **33**, 92-97.
- Casciari, J.J., Sotirchos, S.V. and Sutherland, R.M. (1988) Glucose diffusivity in multicellular tumor spheroids, *Cancer Res*, **48**, 3905-3909.
- Chang, H.Y., *et al.* (1996) The importance of conserved residues in human liver UDPglucose pyrophosphorylase, *European journal of biochemistry / FEBS*, **236**, 723-728.
- Ciaraldi, T.P., Horuk, R. and Matthaei, S. (1986) Biochemical and functional characterization of the rat liver glucose-transport system. Comparisons with the adipocyte glucose-transport system, *The Biochemical journal*, **240**, 115-123.
- Colville, C.A., *et al.* (1993) Kinetic analysis of the liver-type (GLUT2) and brain-type (GLUT3) glucose transporters in *Xenopus* oocytes: substrate specificities and effects of transport inhibitors, *The Biochemical journal*, **290** (Pt 3), 701-706.

Corpas, F.J., *et al.* (1995) Kinetic properties of hexose-monophosphate dehydrogenases. II. Isolation and partial purification of 6-phosphogluconate dehydrogenase from rat liver and kidney cortex, *Molecular and cellular biochemistry*, **144**, 97-104.

Corpas, F.J., *et al.* (1995) Kinetic properties of hexose-monophosphate dehydrogenases. I. Isolation and partial purification of glucose-6-phosphate dehydrogenase from rat liver and kidney cortex, *Life sciences*, **56**, 179-189.

Cuatrecasas, P. and Segal, S. (1965) Mammalian Galactokinase. Developmental and Adaptive Characteristics in the Rat Liver, *The Journal of biological chemistry*, **240**, 2382-2388.

Debbaut, C., *et al.* (2014) Analyzing the human liver vascular architecture by combining vascular corrosion casting and micro-CT scanning: a feasibility study, *J Anat*, **224**, 509-517.

Debbaut, C., *et al.* (2012) Perfusion characteristics of the human hepatic microcirculation based on three-dimensional reconstructions and computational fluid dynamic analysis, *J Biomech Eng*, **134**, 011003.

Diepenbrock, F., *et al.* (1992) Colorimetric determination of galactose and galactose-1-phosphate from dried blood, *Clinical biochemistry*, **25**, 37-39.

Duggleby, R.G., *et al.* (1996) Sequence differences between human muscle and liver cDNAs for UDPglucose pyrophosphorylase and kinetic properties of the recombinant enzymes expressed in *Escherichia coli*, *European journal of biochemistry / FEBS*, **235**, 173-179.

Elliott, K.R. and Craik, J.D. (1982) Sugar transport across the hepatocyte plasma membrane, *Biochemical Society transactions*, **10**, 12-13.

Everett, N.B., Simmons, B. and Lasher, E.P. (1956) Distribution of blood (Fe 59) and plasma (I 131) volumes of rats determined by liquid nitrogen freezing, *Circulation research*, **4**, 419-424.

Facchiano, A. and Marabotti, A. (2010) Analysis of galactosemia-linked mutations of GALT enzyme using a computational biology approach, *Protein engineering, design & selection : PEDS*, **23**, 103-113.

Fraser, R., *et al.* (1988) Nicotine decreases the porosity of the rat liver sieve: a possible mechanism for hypercholesterolaemia, *Br J Exp Pathol*, **69**, 345-350.

Fridovich-Keil, J.L. (2006) Galactosemia: the good, the bad, and the unknown, *Journal of cellular physiology*, **209**, 701-705.

Fukuchi, T., *et al.* (1994) Recombinant rat nucleoside diphosphate kinase isoforms (alpha and beta): purification, properties and application to immunological detection of native isoforms in rat tissues, *Biochimica et biophysica acta*, **1205**, 113-122.

Garipey, L., *et al.* (1993) Clearance by the liver in cirrhosis. II. Characterization of propranolol uptake with the multiple-indicator dilution technique, *Hepatology*, **18**, 823-831.

Geeganage, S. and Frey, P.A. (1998) Transient kinetics of formation and reaction of the uridylyl-enzyme form of galactose-1-P uridylyltransferase and its Q168R-variant: insight into the molecular basis of galactosemia, *Biochemistry*, **37**, 14500-14507.

Goresky, C.A. (1963) A linear method for determining liver sinusoidal and extravascular volumes, *The American journal of physiology*, **204**, 626-640.

Gould, G.W., *et al.* (1991) Expression of human glucose transporters in *Xenopus* oocytes: kinetic characterization and substrate specificities of the erythrocyte, liver, and brain isoforms, *Biochemistry*, **30**, 5139-5145.

Groebe, K., Erz, S. and Mueller-Klieser, W. (1994) Glucose diffusion coefficients determined from concentration profiles in EMT6 tumor spheroids incubated in radioactively labeled L-glucose, *Advances in experimental medicine and biology*, **361**, 619-625.

Gross, J.F. and Aroesty, J. (1972) Mathematical models of capillary flow: a critical review, *Biorheology*, **9**, 225-264.

Guynn, R.W., *et al.* (1974) The concentration and control of cytoplasmic free inorganic

pyrophosphate in rat liver in vivo, *The Biochemical journal*, **140**, 369-375.

Horn, T., Christoffersen, P. and Henriksen, J.H. (1987) Alcoholic liver injury: defenestration in noncirrhotic livers--a scanning electron microscopic study, *Hepatology*, **7**, 77-82.

Irie, M., *et al.* (1970) Distribution and properties of alkaline pyrophosphatases of rat liver, *Journal of biochemistry*, **67**, 47-58.

Jakab, F., *et al.* (1995) The interaction between hepatic arterial and portal venous blood flows; simultaneous measurement by transit time ultrasonic volume flowmetry, *Hepato-gastroenterology*, **42**, 18-21.

Kashiwaya, Y., *et al.* (1994) Control of glucose utilization in working perfused rat heart, *The Journal of biological chemistry*, **269**, 25502-25514.

Keppler, D. and Decker, K. (1969) Studies on the mechanism of galactosamine-1-phosphate and its inhibition of UDP-glucose pyrophosphorylase, *European journal of biochemistry / FEBS*, **10**, 219-225.

Keppler, D., *et al.* (1969) Changes in uridine nucleotides during liver perfusion with D-galactosamine, *FEBS letters*, **4**, 278-280.

Keppler, D., Rudigier, J. and Decker, K. (1970) Trapping of uridine phosphates by D-galactose in ethanol-treated liver, *FEBS letters*, **11**, 193-196.

Keppler, D.O., *et al.* (1970) The trapping of uridine phosphates by D-galactosamine. D-glucosamine, and 2-deoxy-D-galactose. A study on the mechanism of galactosamine hepatitis, *European journal of biochemistry / FEBS*, **17**, 246-253.

Kimura, N. and Shimada, N. (1988) Membrane-associated nucleoside diphosphate kinase from rat liver. Purification, characterization, and comparison with cytosolic enzyme, *The Journal of biological chemistry*, **263**, 4647-4653.

Knop, J.K. and Hansen, R.G. (1970) Uridine diphosphate glucose pyrophosphorylase. IV. Crystallization and properties of the enzyme from human liver, *The Journal of biological chemistry*, **245**, 2499-2504.

König, M., Bulik, S. and Holzhütter, H.G. (2012) Quantifying the contribution of the liver to glucose homeostasis: a detailed kinetic model of human hepatic glucose metabolism, *PLoS computational biology*, **8**, e1002577.

Koo, A. and Liang, I.Y. (1979) Microvascular filling pattern in rat liver sinusoids during vagal stimulation, *J Physiol*, **295**, 191-199.

Koo, A., Liang, I.Y. and Cheng, K.K. (1975) The terminal hepatic microcirculation in the rat, *Q J Exp Physiol Cogn Med Sci*, **60**, 261-266.

Kuntz, E. and Kuntz, H.D. (2006) *Hepatology - Principles and Practice*.

Lai, K., *et al.* (2003) GALT deficiency causes UDP-hexose deficit in human galactosemic cells, *Glycobiology*, **13**, 285-294.

Lam, S.C. and Packham, M.A. (1986) Isolation and kinetic studies of nucleoside diphosphokinase from human platelets and effects of cAMP phosphodiesterase inhibitors, *Biochemical pharmacology*, **35**, 4449-4455.

Lautt, W.W. (2009). In, *Hepatic Circulation: Physiology and Pathophysiology*. San Rafael (CA).

Leslie, N.D. (2003) Insights into the pathogenesis of galactosemia, *Annual review of nutrition*, **23**, 59-80.

Loud, A.V. (1968) A quantitative stereological description of the ultrastructure of normal rat liver parenchymal cells, *The Journal of cell biology*, **37**, 27-46.

MacPhee, P.J., Schmidt, E.E. and Groom, A.C. (1995) Intermittence of blood flow in liver sinusoids, studied by high-resolution in vivo microscopy, *Am J Physiol*, **269**, G692-698.

MacPhee, P.J., *et al.* (1988) Microcirculatory changes in livers of mice infected with murine hepatitis virus. Evidence from microcorrosion casts and measurements of red cell velocity,

Microvasc Res, **36**, 140-149.

Milo, R., *et al.* (2010) BioNumbers--the database of key numbers in molecular and cell biology, *Nucleic Acids Res*, **38**, D750-753.

Muto, M., Nishi, M. and Fujita, T. (1977) Scanning electron microscopy of human liver sinusoids, *Arch Histol Jpn*, **40**, 137-151.

Orfanos, A.P., Jinks, D.C. and Guthrie, R. (1986) Microassay for estimation of galactose and galactose-1-phosphate in dried blood specimens, *Clinical biochemistry*, **19**, 225-228.

Ozer, N., Aksoy, Y. and Ogus, I.H. (2001) Kinetic properties of human placental glucose-6-phosphate dehydrogenase, *The international journal of biochemistry & cell biology*, **33**, 221-226.

Parthasarathy, R., Parthasarathy, L. and Vadnal, R. (1997) Brain inositol monophosphatase identified as a galactose 1-phosphatase, *Brain research*, **778**, 99-106.

Puhl, G., *et al.* (2003) Noninvasive in vivo analysis of the human hepatic microcirculation using orthogonal polarization spectral imaging, *Transplantation*, **75**, 756-761.

Quan-Ma, R., *et al.* (1966) Galactitol in the tissues of a galactosemic child, *Am J Dis Child*, **112**, 477-478.

Quick, C.B., Fisher, R.A. and Harris, H. (1974) A kinetic study of the isozymes determined by the three human phosphoglucomutase loci PGM1, PGM2, and PGM3, *European journal of biochemistry / FEBS*, **42**, 511-517.

Quimby, B.B., *et al.* (1997) Characterization of two mutations associated with epimerase-deficiency galactosemia, by use of a yeast expression system for human UDP-galactose-4-epimerase, *American journal of human genetics*, **61**, 590-598.

Quimby, B.B., *et al.* (1996) Functional requirements of the active site position 185 in the human enzyme galactose-1-phosphate uridylyltransferase, *The Journal of biological chemistry*, **271**, 26835-26842.

Renkin, E.M. (1977) Multiple pathways of capillary permeability, *Circ Res*, **41**, 735-743.

Roberts, M.S. and Rowland, M. (1985) Hepatic elimination--dispersion model, *J Pharm Sci*, **74**, 585-587.

Sangiulo, F., *et al.* (2004) Biochemical characterization of two GALK1 mutations in patients with galactokinase deficiency, *Human mutation*, **23**, 396.

Schadewaldt, P., *et al.* (2000) Analysis of concentration and (13)C enrichment of D-galactose in human plasma, *Clinical chemistry*, **46**, 612-619.

Schaff, Z. and Lapis, K. (1990) Fine structure of hepatocytes during the etiology of several common pathologies, *J Electron Microsc Tech*, **14**, 179-207.

Schirmer, W.J., *et al.* (1986) Galactose clearance as an estimate of effective hepatic blood flow: validation and limitations, *The Journal of surgical research*, **41**, 543-556.

Segal, S. (1995) Defective galactosylation in galactosemia: is low cell UDPgalactose an explanation?, *European journal of pediatrics*, **154**, S65-71.

Segal, S. and Rogers, S. (1971) Nucleotide inhibition of mammalian liver galactose-I-phosphate uridylyltransferase, *Biochimica et biophysica acta*, **250**, 351-360.

Slepek, T.I., *et al.* (2007) Involvement of endoplasmic reticulum stress in a novel Classic Galactosemia model, *Molecular genetics and metabolism*, **92**, 78-87.

Tang, M., *et al.* (2012) Correlation assessment among clinical phenotypes, expression analysis and molecular modeling of 14 novel variations in the human galactose-1-phosphate uridylyltransferase gene, *Human mutation*, **33**, 1107-1115.

Thoden, J.B., *et al.* (2002) Structural analysis of the Y299C mutant of Escherichia coli UDP-galactose 4-epimerase. Teaching an old dog new tricks, *The Journal of biological chemistry*, **277**, 27528-27534.

Thuillier, L. (1978) Purification and kinetic properties of human erythrocyte Mg²⁺-dependent

inorganic pyrophosphatase, *Biochimica et biophysica acta*, **524**, 198-206.

Timson, D.J. (2005) Functional analysis of disease-causing mutations in human UDP-galactose 4-epimerase, *The FEBS journal*, **272**, 6170-6177.

Timson, D.J. and Reece, R.J. (2003) Functional analysis of disease-causing mutations in human galactokinase, *European journal of biochemistry / FEBS*, **270**, 1767-1774.

Turnquist, R.L., *et al.* (1974) Uridine diphosphate glucose pyrophosphorylase: differential heat inactivation and further characterization of human liver enzyme, *Biochimica et biophysica acta*, **364**, 59-67.

Villeneuve, J.P., *et al.* (1996) The hepatic microcirculation in the isolated perfused human liver, *Hepatology*, **23**, 24-31.

Vollmar, B. and Menger, M.D. (2009) The hepatic microcirculation: mechanistic contributions and therapeutic targets in liver injury and repair, *Physiol Rev*, **89**, 1269-1339.

Walmsley, A.R., *et al.* (1998) Sugar transporters from bacteria, parasites and mammals: structure-activity relationships, *Trends in biochemical sciences*, **23**, 476-481.

Wang, Z.J., *et al.* (2001) Proton magnetic resonance spectroscopy of brain metabolites in galactosemia, *Annals of neurology*, **50**, 266-269.

Warren, A., *et al.* (2008) Effects of old age on vascular complexity and dispersion of the hepatic sinusoidal network, *Microcirculation*, **15**, 191-202.

Wasilenko, J., *et al.* (2005) Functional characterization of the K257R and G319E-hGALE alleles found in patients with ostensibly peripheral epimerase deficiency galactosemia, *Molecular genetics and metabolism*, **84**, 32-38.

Wells, W.W., *et al.* (1965) The Isolation and Identification of Galactitol from the Brains of Galactosemia Patients, *The Journal of biological chemistry*, **240**, 1002-1004.

Wermuth, B., *et al.* (1982) Purification and characterization of human-brain aldose reductase, *European journal of biochemistry / FEBS*, **127**, 279-284.

Wermuth, B. and von Wartburg, J.P. (1982) Aldose reductase from human tissues, *Methods in enzymology*, **89 Pt D**, 181-186.

Wiener, J., *et al.* (1968) A quantitative description of cortisone-induced alterations in the ultrastructure of rat liver parenchymal cells, *J Cell Biol*, **37**, 47-61.

Winans, K.A. and Bertozzi, C.R. (2002) An inhibitor of the human UDP-GlcNAc 4-epimerase identified from a uridine-based library: a strategy to inhibit O-linked glycosylation, *Chemistry & biology*, **9**, 113-129.

Wisse, E., *et al.* (1996) Structure and function of sinusoidal lining cells in the liver, *Toxicol Pathol*, **24**, 100-111.

Wisse, E., *et al.* (1985) The liver sieve: considerations concerning the structure and function of endothelial fenestrae, the sinusoidal wall and the space of Disse, *Hepatology*, **5**, 683-692.

Wisse, E., *et al.* (2008) The size of endothelial fenestrae in human liver sinusoids: implications for hepatocyte-directed gene transfer, *Gene Ther*, **15**, 1193-1199.

Wohlert, T.M. and Fridovich-Keil, J.L. (2000) Studies of the V94M-substituted human UDPgalactose-4-epimerase enzyme associated with generalized epimerase-deficiency galactosaemia, *Journal of inherited metabolic disease*, **23**, 713-729.

Wynne, H.A., *et al.* (1989) The effect of age upon liver volume and apparent liver blood flow in healthy man, *Hepatology*, **9**, 297-301.

Yamaguchi, A., *et al.* (1989) Microassay for screening newborns for galactosemia with use of a fluorometric microplate reader, *Clinical chemistry*, **35**, 1962-1964.

Yoshida, C., Shah, H. and Weinhouse, S. (1982) Purification and properties of inorganic pyrophosphatase of rat liver and hepatoma 3924A, *Cancer research*, **42**, 3526-3531.

